

Midwest Fruit Pest Management Guide

2026-2027

Arkansas

University of Arkansas Cooperative Extension Service
AG1304

Illinois

University of Illinois Extension
ICSG-18

Indiana

Purdue Extension
ID-465

Iowa

Iowa State University Extension and Outreach
HORT 3035

Kansas

Kansas State Research and Extension
MF3278

Kentucky

University of Kentucky Cooperative Extension Service
ID-232

Ohio

Ohio State University Extension
Bulletin 506

Wisconsin

University of Wisconsin-Extension
A4104



Midwest Fruit Pest Management Guide 2026-2027

Citation

Long, E., Leach, A., Athey, K., Wimmer, M., Arnoldussen, B., Yadav, R., Meyer, S., Purcell, M., Santiago, J., Holland, L., Joshi, N., Whale, E., Babadoost, M., Vos, R., Slack, S., Cloyd, R., Bessin, R., Ivey, M., Wright, S., Beckerman, J., Leonberger, K., Gauthier, N. 2025. Midwest Fruit Pest Management Guide 2026-2027. Available online at: https://ag.purdue.edu/department/hla/extension/_docs/id-465.pdf

Contributors

University of Arkansas

Entomology: Neelendra Joshi

University of Illinois

Entomology: Kacie Athey*

Horticulture: Elizabeth Wahle*, James Santiago*

Plant Pathology: Mohammed Babadoost*

Iowa State University

Horticulture: Randall Vos*, Suzanne Slack

Kansas State University

Entomology: Raymond Cloyd

University of Kentucky

Entomology: Ric Bessin*

Horticulture: Shawn Wright*, Brent Arnoldussen*, Elayna Stirn

Plant Pathology: Nicole Gauthier*, Kim Leonberger*

University of Minnesota

Horticulture: Madeline Wimmer*

Ohio State University

Entomology: Ashley Leach*

Horticulture: Ram Yadav*

Plant Pathology: Melanie Ivey*

Purdue University

Entomology: Elizabeth Long*

Horticulture: Stephen Meyer, Miranda Purcell

University of Wisconsin

Plant Pathology: Leslie Holland*

*Section Editor

Acknowledgements

Authors would like to thank Janna Beckerman for her years of hard work on this guide.

Foreword

Commercial fruit production has become a highly skilled, technical profession. Concerns about pesticide

Contents

| | |
|--|-----|
| Forward..... | 3 |
| Apple..... | 13 |
| Pear..... | 56 |
| Cherry..... | 76 |
| Peach..... | 90 |
| Plum..... | 116 |
| Grape..... | 134 |
| Blueberry..... | 165 |
| Raspberry and Blackberry..... | 178 |
| Strawberry..... | 194 |
| Vole Control..... | 214 |
| Growth Regulator PHIs and REIs..... | 216 |
| Chemical Weed Control in Fruit Crops..... | 216 |
| Herbicide Recommendations..... | 218 |
| Generic Pesticides..... | 250 |
| Suggested Recordkeeping Form for Restricted Use Pesticide .. | 261 |
| Fruit Grower Newsletters..... | 262 |
| Pesticide Drift Communication Tools..... | 262 |
| Using a Plant Diagnostic Lab..... | 263 |
| Pesticide Applicator Safety Education Programs..... | 265 |
| Pesticide Emergency and Poison Control Centers..... | 265 |
| Conversion Factors for Weights and Measures..... | 266 |

residues, operator risks, and the environment dictate that all fruit growers exercise extreme caution in the use of all pesticides, and indeed, all chemicals. Growers who use these must be certified as private applicators.

Certification requires that applicators understand the following: labels and labeling, safety factors, potential environmental concerns, identification of common pests encountered, pesticides and their use, proper equipment use, application techniques, and applicable state and federal regulations. Training programs are offered to help you in certification. Contact your county Extension office for information.

The pest management recommendations in this guide have been formulated to provide growers with up-to-date information about pesticides and their applicability to problems. We suggest that growers use this information to set up individualized spray programs. Always keep accurate records of materials used, application dates, areas treated, growth stages, and weather conditions. A sample record sheet is on page 285. In case of questions, nothing beats a good set of records. The EPA requires records for restricted use pesticide applications. Some states may require records for general use pesticides (e.g., Kentucky has this requirement).

Handling Pesticides

1. Know the pesticide toxicity and read and follow all label instructions, as well as any websites directed by the label.
2. When mixing pesticides, do not breathe the dust, powder, or vapor. Always mix outdoors.
3. Do not use tobacco, eat, or drink when handling or applying pesticides.
4. Stay out of drift from spray or dust.
5. Rinse liquid containers with water at least three times and pour rinsate into spray tank as it is being filled. Punch holes in metal and plastic containers and crush. Dispose of these and all other pesticide containers where no contamination of crops or water supply can occur. Do not reuse pesticide containers.
6. Use all personal protective equipment as required by the label. If a respirator is required, a medical exam and fit test is needed for some types of respirators.
7. Have a "buddy" around when using acutely toxic organophosphates, just in case.
8. If working with cholinesterase inhibitor insecticides, get an appropriate blood test before the season starts and test periodically during the season.
9. Consult a doctor immediately if you develop unusual symptoms during or after spraying. Symptoms such as blurred vision, nausea, headaches, chest pains, weakness, diarrhea, or cramps indicate possible pesticide poisoning.
10. Wash hands thoroughly before eating, drinking, chewing gum, using tobacco, or using the toilet.
11. Bathe and change clothes daily, and wash contaminated clothing separate from other laundry.
12. Always store pesticides in their original container, never in an unmarked container. Never trust your memory.
13. Always store pesticides under lock and key and keep them away from children.
14. Always use an anti-siphon device when filling the spray tank from a domestic water source.
15. The label is the law. Read and follow all label instructions carefully.
16. Never allow someone under the age of 18 to handle or use pesticides.
17. Train workers according to worker protection standards (WPS). This is required for all workers who will enter areas treated with a pesticide within 30 days after the last application.
18. Suspend applications while people are within the Application Exclusion Zone (AEZ). Application may be resumed when they exit the AEZ.

Management Tips for Safety

1. Maintain accurate spray records. Show application rates, pesticides used, total gallonage, area treated, stage of plant development, and weather data.
2. Be prepared to show your records to the EPA or state regulatory officials if necessary.
3. Do not contaminate forage crops or pastures.
4. Do not allow animals to graze fruit plantings.
5. Prevent excess drift.
6. Maintain equipment in top condition.
7. Protect children, pets, livestock, and the environment from pesticide contamination.
8. Follow all label instructions on re-entry times for pesticides. Regulations mandate re-entry times for all pesticides. Check labels for posting and notification requirements to protect workers. Re-entry times and the required protective clothing for early entry are listed on product.
9. Inform all workers of re-entry restrictions and information on safe pesticide use and/or training to meet WPS requirements.
10. Comply with the Right-To-Know law. Have complete product labels readily available for workers to see. Have the Safety Data Sheet (SDS) for each product you use available for workers to see and for rescue or fire personnel to use in case of emergency.
11. Provide pesticide safety training for pesticide handlers and other workers to comply with Worker Protection Standards (WPS). Training content must comply with WPS requirements.
12. Provide necessary PPE for your workers and regularly inspect and maintain PPE used when applying pesticides.

Pesticide Use and the Law

Pesticides are developed by manufacturers, registered with EPA, and sold to the public with the assumption that users read, understand, and follow instructions on product labels. Pesticide labels include specific information about use, personal protective equipment, environmental precautions, and storage and disposal. The label's purpose is to provide clear directions to allow maximum product benefit while minimizing risks to human health and the environment.

Every pesticide label includes the following statement: "It is a violation of federal law to use this product in a manner inconsistent with its labeling." This language obliges purchasers or users of any pesticide to assume all legal responsibilities for the product's use. Further, courts and regulators recognize that pesticide labels are binding contracts that require those using the products to do so exactly as directed. Terms such as

“must,” “shall,” “do not,” and “shall not” are mandatory statements, users are responsible for specific actions when applying or handling a given product; any departure from such directions is, in the eyes of the law, an illegal use of the pesticide.

“Use” means more than just applying the pesticide. Federal and state regulations define pesticide use to include handling, mixing, loading, storing, transporting, and disposing, as well as human and environmental exposure. This all-encompassing definition covers every activity that involves a pesticide — from purchase to container disposal.

The pesticide label is more than just a piece of paper. It serves a dual function: the label instructs users how to use the product safely and effectively, and it serves as a legal measuring stick. Many statements on the label result from rigorous scientific investigations and governmental regulatory decisions. Pesticide users should read, understand, and follow pesticide label directions to ensure effective pest control, personal safety, environmental protection, and legal compliance.

Pesticide labels include two important statements:

Re-entry or restricted entry interval (REI) statements contain re-entry precautions and state a time interval during which entry into a pesticide-treated site is prohibited. The statement indicates the length of time that must elapse after the pesticide application before individuals may enter the treated area without personal protective clothing and equipment (PPE).

Pre-harvest interval (PHI) statements indicate the time interval that must elapse after the pesticide application before the crop may be harvested. Harvesting prior to the PHI may result in dangerous and illegal pesticide residues on the crop.

Pesticide Use in Greenhouses and High Tunnels

Fruit and vegetable production in greenhouses and high tunnels have increased dramatically in the Midwest in the past few years. Although greenhouse or high tunnel environments may change the composition of the pest complex, using pesticides is often necessary to maintain the adequate levels of control needed to produce a profitable and marketable crop.

Pesticide regulatory agencies in the Midwest vary in their interpretation of whether a high tunnel is a type of greenhouse. For example, most states consider a high tunnel to be a form of greenhouse. That means the pesticides one selects for high tunnel use must be appropriate for greenhouse use. Other states (not covered by this guide) consider high tunnels to be the same as fields when it comes to pesticide use. Still other states, like Missouri, take an intermediate approach: they call a high tunnel a greenhouse when the

sides are closed but call it a field when the sides are open.

It is important to determine how your state views high tunnels. Pesticide labels address greenhouse applications in one of three ways:

- Pesticide labels can clearly state that the products may be used in greenhouses. These products may be used according to label directions. Pesticide labels that have different instructions for greenhouse use. These products also may be used in high tunnels according to label instructions.
- Pesticide labels may clearly prohibit greenhouse use. Obviously, these products cannot be used in a greenhouse under any circumstances.
- Many pesticide labels don't specify whether the product can be used in a greenhouse or not. When labels don't expressly prohibit greenhouse use, most state regulatory agencies interpret that to mean the product can be used in a greenhouse as long as the treated crop is on the label and the product is used according to label directions.

Determining Spray Volume and Rate

Producers spray fruit plantings with insecticides, fungicides, growth regulators, and nutrient solutions in many different formulations and concentrations and at various stages of plant development. The principal targets may be the foliage, flowers, fruit, woody surfaces, or entire plants. The equipment and methods used for such a diverse spray program must be versatile, and the equipment must be properly calibrated for each type of application.

Dilute Spraying

The objective of spraying is to distribute the spray material uniformly over plants or plant parts. Pesticide recommendations are based on the amount of dilute spray needed to wet plants thoroughly, to the point just “prior to runoff.” For example, in typical blueberry, raspberry, or grape plantings with plants 5 to 7 feet tall and 3 to 5 feet wide and set in rows 9 to 10 feet apart, and in most strawberry plantings, 100 gallons of water per acre has been established as a standard dilute spray volume for fungicide and insecticide application. This dilute rate is considered a 1x concentration.

In a standard apple or pear orchard, with trees approximately 20 feet tall, 22 feet wide, and set on rows 35 feet apart, 400 gallons of water per acre is a standard dilute spray for fungicide and insecticide application. Recommendations may be made per 100 gallons or per acre. Dilute is considered 1x concentration. For cherry, peach, and plum, 300 gallons of water per acre is the standard dilute spray volume for full-size trees.

Amount of Dilute Spray per Acre Required for Equivalent Coverage of Plants

The table on page 6 lists the gallons of dilute spray per acre required to provide equivalent coverage for mature trees of different sizes and spacings.

Growth regulators may be applied by high-volume hand-gun or air-blast sprayers, in either dilute or

low-volume applications. Low-volume application may be riskier because any mistakes in concentration are magnified. Read the growth regulator label for suggestions about application methods. Some labels suggest dilute sprays with full coverage, and others suggest a specific amount of chemical in a specific amount of water per acre.

Amount of Dilute Spray Per Acre Required For Equivalent Coverage Of Plants

| Distance Between Rows (Feet) | Plant Height (Feet) | Plant Width (Feet) | Maximum Plant Volume/ Acre (1000 Cu Ft ¹) | Minimum Dilute Spray (Gallons/Acre ²) |
|------------------------------|---------------------|--------------------|---|---|
| 30 | 20 | 15 | 436 | 300 |
| 26 | 16 | 12 | 354 | 225 |
| 24 | 14 | 10 | 254 | 180 |
| 22 | 14 | 10 | 272 | 200 |
| 20 | 12 | 10 | 261 | 185 |
| 18 | 10 | 10 | 242 | 175 |
| 16 | 8 | 8 | 174 | 125 |
| 14 | 6 | 6 | 149 | 105 |
| 12 | 6 | 6 | 131 | 90 |
| 10 | 6 | 4 | 105 | 74 |
| 10 | 4 | 4 | 70 | 49 |

¹Maximum plant volume/acre = plant width x plant height x running feet or row per acre. Running feet of row per acre = 43,560 divided by the distance between rows.

²Minimum dilute gallons per acre = approximately 0.7 gallon /1,000 cubic feet of plant volume.

Low-volume Spraying

Low-volume, or concentrate, spraying is the practice of using less water per acre to apply pesticides. In low-volume spraying, the volume of water applied per acre is reduced in proportion to the increased concentration of pesticide used by 2x, 3x, 4x, or more. Thus, a 3x rate uses a 3x concentration of pesticide in only one-third the water per acre that would be used in dilute spraying.

Growers must apply low-volume sprays with air-assisted sprayers that use a high-velocity airstream to distribute the spray mixture. Most conventional air-assisted sprayers can be used to apply spray mixtures up to 6x concentration. Sprayers specifically designed for ultra-low-volume application should be used for applications up to 6x.

Using low-volume sprays requires less labor, less water, less time, and fewer refills than 1x or dilute mixtures. However, low volume sprays have disadvantages. Savings in gallonage and application costs

decrease most rapidly to about 50 gallons of water per acre (on tree fruit). Below that, the savings may not be worth the additional risk of improper application and wind drift.

When making low-volume pesticide applications:

1. Use extreme care in calibrating the sprayer and maintaining a constant sprayer speed. As gallonage decreases, errors become much more critical.
2. Choose calm winds with good drying conditions for spraying. This may mean spraying at night or early in the morning. Good coverage cannot be achieved in windy conditions (more than 5 mph).
3. Prune plants to create an open canopy for spray penetration. Spray droplets will not penetrate dense foliage.
4. Choose pesticide formulations that will mix satisfactorily. Pay careful attention to increased operator hazards and drift problems.

Gallons Of Spray Per Acre (Approximate) For Various Concentrates

| | 1x | 2x | 3x | 4x | 5x | 6x |
|-----------------------------------|-----|------------------|--------------------|-----|-----|-----|
| Apples | 400 | 200 | 132 | 100 | 80 | 64 |
| Peaches | 300 | 150 | 100 | 75 | 60 | 50 |
| Percent water savings over dilute | | 50% | 67% | 75% | 80% | 84% |
| | | Greatest savings | Diminished savings | | | |

Tree Row Volume Spraying

Tree row volume (TRV) is a method originally used with orchard crops to determine the dilute (1x) volume of spray solution necessary to cover the entire plant surface for any given fruit planting. TRV is an objective method for determining the spray volume required for plants of different sizes, and for changes in canopy size as plants develop during the season.

With the TRV method, growers can easily calculate the volume of dilute spray needed per acre for each planting based on plant size and canopy density. To determine the TRV, growers must accurately measure the between-row spacing, maximum plant height, and cross-row plant spread. See the step-by-step procedure below.

Calculate Tree Row Volume Gallonage

Step 1. Calculate feet of row/acre.

$$\frac{43,560 \text{ sq ft/acre}}{\text{between-row spacing (ft)}} = \text{feet of row/acre}$$

Step 2. Calculate cu ft of TRV/acre.

Feet of row/acre (from Step 1) x plant height (ft) x cross-row plant spread (ft) = cu ft of TRV/acre.

Step 3. Select density factor.

Select one of the following numbers that best indicates the canopy density of each separate planting.

0.70 gal/1,000 cu ft: Plants extremely open, light visible through entire canopy.

0.80 gal/1,000 cu ft: Plants well pruned, with moderate vigor, adequate light penetration into canopy, many holes in foliage where light can be seen through plant.

0.90 gal/1,000 cu ft: Plants pruned minimally, or with high vigor, poor light penetration into canopy, very few holes where light can be seen through plant.

1.00 gal/1,000 cu ft: Plants unpruned, extremely dense, no light visible anywhere through canopy

Step 4. Calculate TRV gallonage/acre.

$$\frac{\text{cu ft of TRV/acre (from Step 2)} \times \text{density (from Step 3)}}{1,000}$$

= gallons of dilute solution to be applied per acre

= TRV gal/acre

Example 1

A vineyard has rows spaced 10 feet apart, the canopy height is 6 feet, and the cross row spread is 4 feet at full canopy. The density factor is 0.90.

Step 1 43,560 sq ft ÷ 10 ft = 4,356 ft of row/acre

Step 2 4,356 x 6 ft x 4 ft = 104,544 cu ft TRV/acre

Step 3 Density has been chosen as 0.90.

Step 4 [104,544 x .90] ÷ 1,000 = 94 TRV gal/acre

Example 2

An apple orchard on dwarfing rootstock has rows spaced 15 feet apart, the canopy height is 12 feet, and the cross row spread is 8 feet at full canopy. The density factor is 0.90.

Step 1 43,560 sq ft ÷ 15 ft = 2,904 ft of row/acre

Step 2 2,904 x 12 ft x 8 ft = 278,784 cu ft TRV/acre

Step 3 Density has been chosen as 0.90.

Step 4 [278,784 x .90] ÷ 1,000 = 251 TRV gal/acre

Spraying Small Volumes

In some cases growers may wish to apply small volumes of pesticides with backpack or hand-held sprayers or wipers. The following table helps convert from the rate per 100 gallons to the rate per gallon. Take care to measure pesticide amounts accurately, because errors are magnified at small volumes. (See Approximate Dilutions for Small Volumes of Spray Mixes table on page 8).

Calibrating Single Nozzle and Boom Sprayers

Calibration is an essential step for using any application equipment. Early spring, when preparing sprayers for early season operations, is a good time to calibrate. Be sure all fittings are tight and there are no leaks. Take the nozzles apart, clean them, and check for worn nozzle tips.

Using wettable powder sprays enlarges nozzle openings, so calibrating each nozzle is essential. Start the season with a calibrated sprayer, and depending on the number of gallons typically sprayed, calibrate the

sprayer again according to intervals specified in the owner's manual (or no later than halfway through the

spray season). Follow the procedure below to calibrate a single nozzle boom sprayer.

Approximate Dilutions for Small Volumes of Spray Mixes

| Equivalent Rates For Different Quantities Of Water | | | | |
|--|-----------------|-----------|-----------|----------|
| Formulation | 100 Gallons | 5 Gallons | 3 Gallons | 1 Gallon |
| Wettable Powder, Dry Flowable, etc. | 5 lb. | 15 tbsp. | 9 tbsp. | 3 tbsp. |
| | 4 lb. | 13 tbsp. | 8 tbsp. | 8 tsp. |
| | 3 lb. | 10 tbsp. | 6 tbsp. | 2 tbsp. |
| | 2 lb. | 8 tbsp. | 4 tbsp. | 4 tsp. |
| | 1 lb. | 3 tbsp. | 6 tsp. | 2 tsp. |
| | 1/2 lb. (8 oz.) | 5 tsp. | 1 tbsp. | 1 tsp. |
| Emulsifiable Concentrate, Liquid | 5 gal. | 1 qt. | 1 1/4 pt. | 13 tbsp. |
| | 4 gal. | 1 1/2 pt. | 1 pt. | 10 tbsp. |
| | 3 gal. | 1 1/4 pt. | 3/4 pt. | 8 tbsp. |
| | 2 gal. | 3/4 pt. | 1/2 pt. | 5 tbsp. |
| | 1 gal. | 1/2 pt. | 8 tbsp. | 3 tbsp. |
| | 1 qt. | 3 tbsp. | 2 tbsp. | 2 tsp. |
| | 1 pt. | 5 tsp. | 1 tbsp. | 1 tsp. |

These approximations are based on average weights of various pesticide products as described in Dry Pesticide Rates for Hand-held Sprayers (University of Kentucky Extension publication HO-83, <https://www.uky.edu/Ag/Horticulture/masabni/Publications/HO-83.pdf>).

Step 1. Check tractor/sprayer speed.

Attach the sprayer to the tractor and make test runs to determine the tractor speeds (mph) in different gears. Run the tractor at the PTO speed used when operating the sprayer. Travel a test course and record time needed to travel a measured distance. Run the test on the same type surface in the planting (sod, not pavement or gravel)

Formula

$$\text{MPH} = \frac{\text{feet traveled}}{\text{seconds}} \times \frac{60}{88}$$

Your tractor sprayer speed

$$\text{MPH} = \frac{\text{feet traveled}}{\text{seconds}} \times \frac{60}{88} = \underline{\hspace{2cm}}$$

Note: The recommended tractor speed for most applications with single nozzle boom sprayers is 2-3 mph. Traveling faster may lead to poor coverage. A convenient method is to set up a calibration course in multiples of 88 feet (88 feet per minute=1 mile per hour). Set markers at 176 feet or 264 feet to correspond to 2 mph and 3 mph when the tractor speed is adjusted (gear and rpm) to cover the distance in 60 seconds (1 minute).

Step 2. Record the sprayer inputs.

| | Your Figures | Example |
|--|--------------|------------------|
| Nozzle type (all nozzles should be identical) | _____ | 110° 04 flat fan |
| Recommended application volume (from manufacturer's label) | _____ | 20 GPA |
| Measured sprayer speed | _____ | 3 MPH |
| Nozzle spacing/band width (in inches) | _____ | 20 inches |

Step 3. Calculate the required nozzle output.

Formula

$$\text{GPM (per nozzle)} = \frac{\text{GPA} \times \text{MPH} \times \text{W}}{5,940 \text{ (constant)}}$$

Where

GPM = required output per nozzle in gallons per minute.
 GPA = desired total carrier volume in gallons per acre.
 MPH = desired ground speed in miles per hour.
 W = inches between nozzles (or band width if making band applications).

Example

$$\text{GPM} = \frac{20 \text{ GPA} \times 3 \text{ MPH} \times 20 \text{ in}}{5940} = \frac{1,200}{5,940} = 0.20 \text{ GPM}$$

Your figures

$$\text{GPM} = \frac{\quad}{5,940} = \frac{\quad}{5,940} = \text{GPM}$$

Step 4. Operate the sprayer.

Set the correct pressure at the gauge using the pressure-regulating valve. Note that recommendations for flat fan nozzles are 15-30 psi (not more than 40 psi for spraying weeds).

Collect and measure the output of each nozzle for one minute.

The output of each nozzle should be the approximately the same as calculated in Step 3 above. There are 128 fluid ounces in one gallon. If you calculate the output at 0.20 GPM, multiply 0.20 by 128, which equals 25.6 fluid ounces in one minute.

If the nozzle output is slightly off from your target, change the pressure. If the nozzle output is significantly off, change the speed or nozzle size.

Compare nozzle output on multiple nozzle booms. Replace all nozzle tips that are more than 10 percent inaccurate. Consistent spray patterns require that outputs from individual nozzles within 10 percent.

Calibration of Air-blast Sprayers

Accurate calibration is the only way to ensure that a sprayer is applying the intended amount of chemical. Applicators must know the amount of water that will be applied per unit of area to make a proper spray mix. Failing to calibrate the sprayer can injure the crop, create a hazardous situation, and waste money. Frequent calibration identifies worn nozzles and keeps applicators aware of factors that can affect the application rate, including travel speed, pressure, and type of nozzle in use.

Pre-calibration Check

Before calibrating, check the sprayer carefully. Be sure the nozzle tips are clean. Replace all worn or damaged nozzles. Check all hoses and fittings for leaks and aging. Make sure the pressure is constant and the tank is free of dirt and debris.

Determining Sprayer Speed

To determine travel speed needed to travel to properly distribute the spray within the canopy by placing water-sensitive spray paper at various locations within the canopy. For proper pesticide application, the air within the canopy must be completely replaced with

spray-laden air from the sprayer. In general, a travel speed of 1 to 3 miles per hour is usually satisfactory, depending on the size and density of the canopy, and capacity of the sprayer.

Before sprayer calibration, applicators must determine the travel speed in miles per hour (mph). To determine travel speed, load the sprayer with clear water and make a test run in the fruit planting. Always make the test run in the fruit planting or on similar ground, because tractor speeds change dramatically from soft to firm surfaces. Set the tractor throttle at a level sufficient to operate the sprayer (PTO speed) and select an appropriate gear. Remember or mark these settings.

Calculate travel speed by measuring the time required to travel any measured distance. A good conversion factor to remember is that 1 mph=88 feet/min. A convenient test length is 176 feet because it is a multiple (2x) of 88. Use the following formula to determine travel speed:

$$\text{Speed (MPH)} = \frac{\text{distance (ft)} \times 60}{\text{time (sec)} \times 88}$$

For example, if it requires 60 seconds to travel a measured distance of 176 feet, the travel speed is:

$$\text{MPH} = \frac{176 \times 60}{60 \times 88} = \frac{10,560}{5,280} = 2 \text{ MPH}$$

Determining Nozzle Flow Rate

To select the correct nozzle and whirlplate sizes, first determine the total gallons per minute (gpm) of output for each particular application.

To determine gpm, you must know the travel speed of the sprayer (mph), the gallons per acre (gpa) to be applied, and the spacing (W) between the rows of plants. These variables will be used, to calculate the gpm. Note: This equation is for one side of the sprayer manifold only; double the calculated answer if using both sides of the sprayer. Use the same size nozzles and whirlplates on both sides of the sprayer.

Step 1. Calculate the total gpm required per side:

$$\text{GPM (per side)} = \frac{\text{GPA} \times \text{MPH} \times \text{W}}{1,000}$$

GPM = gallons per minute (per side)

GPA = gallons per acre

MPH = speed (in miles per hour)

W = spacing between rows (in feet)

Example: To apply 70 gpm while traveling 2 mph, with row spacing of 10 feet apart the gpm per side is:

$$\text{GPM} = \frac{70 \times 2 \times 10}{1,000} = \frac{1,400}{1,000} = 1.4 \text{ GPM}$$

Step 2. Select the correct nozzle-whirlplate combination and operating pressure. Air-blast sprayers normally use disk-core-type cone spray tips. Select the correct size nozzles and whirlplates by using a table that indicates the nozzle size and gallons per minute output at various pressures using specific whirlplates. You can find these tables in the sprayer manufacturer's literature or in nozzle catalogs.

Arrange nozzles in the sprayer manifold so approximately *two-thirds of the total flow comes from nozzles in the upper half* of the manifold, and *one-third of the total flow comes from nozzles in the lower half*. Adjust nozzles this way to provide uniform coverage throughout the canopy. It should adequately penetrate to the top and center of the canopy while avoiding excess application in the lower outside areas.

Step 3. Install the nozzles in their proper outlets. Inspect and clean all nozzles and outlets and determine that the sprayer is operating correctly. Nozzles are an important part of the sprayer; if the nozzles show any defects or wear, replace them.

Step 4. Measure the total gpm from all the nozzles selected in Step 2. Fill the sprayer tank at least half full. Prime the sprayer system and check all the nozzles to make sure none are clogged or partially clogged. Record the exact level of water in the spray tank. Bring the sprayer up to the desired pressure and turn the nozzles on. Use a stopwatch to record how long the sprayer is running. You should operate the sprayer for at least three minutes. Record the new level in the tank or measure the amount of water needed to refill the tank to the original level.

Example: The spray tank is filled to the 100-gallon level. It was predetermined from the manufacturer's tables that the nozzles selected would give a total output of 4 gpm. The sprayer was operated for five minutes at 150 psi on the gauge. After the five minutes, the sight gauge read 75 gal. The actual output was:

$$100 \text{ gal (start)} - 75 \text{ gal (stop)} = 25 \text{ gal per 5 min.} = 5 \text{ gpm}$$

The theoretical output from table information, however, was 4 gpm.

When actual output differs from the calculated output, adjust by changing the pressure (when the difference is small) or changing the nozzle sizes (when the difference is large). Experiment with the pressure to see if the output can be fine-tuned. Refer to manufacturer's tables for recommended operating pressures for nozzles. Never operate above or below recommended pressures.

Repeat these calibration procedures whenever you change the speed, gallons per acre, or row spacing. Periodically check the output from the nozzles during the spraying season. The effectiveness of the spray material directly depends on your skill as an operator.

Field test to confirm calculations:

$$\text{GPA (gallons per acre)} = \frac{\text{gal sprayed} \times 43,560 \text{ ft}^2}{\text{distance traveled (ft)}}$$

Example: A field test is run in which 10 rows, each 200 feet long, with row spacing of 10 feet and it took 35 gallons to refill the sprayer to the original level, the gpa is:

$$\frac{35 \text{ gal} \times 43,560 \text{ ft}^2}{2,000 \text{ ft} \times 10 \text{ ft}} = 76 \text{ GPA}$$

Spray Water pH

Several pesticides break down rapidly in alkaline water (pH above 7.0). Both well and pond water in the Midwest tend to be alkaline. In a matter of hours — or, in extreme instances, only minutes — 50 percent or more of the active ingredient may be hydrolyzed to yield a less active compound. Captan, Dimethoate, Imidan, and Malathion are examples of compounds especially vulnerable to alkaline hydrolysis.

To ensure the maximum effectiveness of pesticide applications, check the pH of spray mixes in the spray tank and add buffering agents if necessary to adjust the pH to neutral (7.0). Many commercial buffering agents are available. Most adjuvants (see definition in the next section) are multipurpose adjuvants, serving as spreaders, activators, etc. Read the labels of both the pesticide and adjuvant before use. Granulated food grade citric acid may be the most convenient and inexpensive acidifying material. Two ounces per 100 gallons has been shown to reduce the pH of tap water from 8.3 to 5.4. Convenient granulated food grade citric acid measures are:

| | |
|-------------|---------------------------|
| per 100 gal | 1/4 cup, slightly rounded |
| per 300 gal | 3/4 cup, rounded |
| per 500 gal | 1 1/3 cups |

Granulated food grade citric acid is available in 50-pound bags from suppliers that handle food grade chemicals. Do not try to acidify solutions containing phosphorous acid, Bordeaux mixture, fixed copper, or other copper compounds.

Spray Adjuvants

Several types of additives are available to improve the effectiveness of spray applications. Collectively, these products are called adjuvants. Here are some adjuvants and their functions:

Activators increase a pesticide's efficacy by increasing the penetration of a spray solution through leaf hairs or waxy cuticle and into a leaf or fruit.

Acidifiers lower the pH of alkaline spray water to reduce the potential breakdown of certain pesticides in the spray tank.

Buffers change the pH of spray water, then hold it at the desired degree of acidity.

De-foamers, when added to the spray tank, break down or prevent the formation of foam.

Elasticizers or drift control agents reduce the breakup of spray droplets into very fine particles and thereby minimize drift.

Surfactants, spreaders, and wetting agents are different names for products that reduce the surface tension around spray droplets, allowing them to spread out more evenly on the surface of leaves or fruit.

Caution: Some surfactants used in combination with certain pesticides can function as activators, which can injure plants. Consult labels or chemical suppliers for more information.

Stickers cause a pesticide to stick to the surface after the spray dries, thereby reducing the potential for loss from rain or overhead irrigation.

Spreader-stickers is a term commonly misused when referring to a surfactant or spreader. A true spreader-sticker combines the characteristics of a surfactant with that of a sticker.

Caution: Do not use an adjuvant with any pesticide without first consulting the specific pesticide label. Improper selection or use can injure crops or reduce effectiveness, particularly when adjuvants are mixed with emulsifiable concentrates.

Pesticide Compatibility

Because of the complex nature of pest management in fruit crops, multiple fungicides and insecticides may need to be tank-mixed together and applied at once. Pesticide compatibility in the spray tank is usually not a problem with newer pesticides. The compatibility of some materials may depend on solvents and emulsifiers within the formulation. Emulsifiable concentrate formulations are more likely to cause compatibility problems than wettable powders. Mixing wettable powders with emulsifiable concentrates may result in incompatibility issues. Compatibility problems are often noted when applicators use lime, copper (Bordeaux), or oil products in a mix. Be aware of spray tank pH as noted above.

Read the comments section in this spray guide for notes about compatibility problems, and read pesticide labels before tank mixing products. Most pesticide

labels give instructions for loading, tank mixes, etc., and we recommend that growers follow the label directions closely to avoid problems.

Making Tank Mixes

Adding the components of a mixture in the order the label specifies is critical; whether some pesticides are physically compatible or not depends on the order in which they are added to the tank. This is especially true for pesticides packaged in water-soluble packets. A mistake in mixing order could prevent the package from dissolving completely, thus preventing uniform distribution of the pesticide in the spray tank. Labels provides mixing instructions for all registered tank mixes. Unless the label states otherwise, never apply any pesticide in the mixture at a rate higher than the label allows for when the pesticide is used alone for the same purpose.

Some pesticide labels do not provide adequate mixing order directions. The usual method for tank mixing is as follows.

First, fill the tank one-quarter to one-half full with the carrier (water) and begin agitation. If a compatibility, buffering or defoaming agent is needed, these products should be added before the other products. If using a drift reduction additive, always consult the label; some are added in the mixing sequence.

Next, slowly add and thoroughly mix the pesticide products, one at a time, beginning with those hardest to mix (such as suspension-forming formulations). Generally, wettable powder (WP) and dry flowable or water-dispersible granule (DF, WDG) products should be added first, followed by flowable (F, FL) and microencapsulated (ME) products. Add emulsifiable concentrates (EC) next, followed by any solution (S) or soluble powder (SP) products. Other spray modifiers (penetrants and surfactants) should be added last. Dry formulations should be preslurried (mixed with a little water) before adding them to the spray tank; this is also a good idea (even with ECs) when using liquid fertilizers as the carrier. Finally, continue adding the carrier to the desired level.

To assure a uniform spray mixture at all times, keep the mixture agitated during the entire application and until the tank is empty. Avoid letting the mixture stand overnight, if possible, without agitation. If you do end up with a physically incompatible spray mix, call the manufacturer of each product to see if you can rejuvenate the mix. Adding a compatibility agent may return the mix to a sprayable form. If you cannot rejuvenate the mix, treat it as pesticide waste.

Tank Mixing Order

1. Fill tank 1/4 to 1/2 full with carrier (water)
2. Begin agitation
3. Utility agents (if needed)
4. Suspension products
 - a. Dry (Pre-mix): WP, DF, WDG,
 - b. Wet F, FL, ME
5. Emulsifiable products (EC)
6. Solution products (S, SP)
7. Spray modifiers (if needed)
8. Finish filling the tank with carrier

From: Illinois Pesticide Applicator Training Manual SP39: General Standards. University of Illinois Extension Pesticide Safety Education Program.

Summary

Pesticide recommendations are confusing because there are so many options for materials to use for certain diseases or insect pests. Additional references may be needed.

With fungicides in particular, a single material may control one or more diseases, but not all. So when several diseases threaten, growers may need to combine materials to achieve control. Insect pests also may appear at the same time, so a tank mix multiple fungicides and insecticides may be applied together. However, not all pesticides are compatible, so applicators should test for compatibility before tank mixing any products.

Certain fungicides and insecticides may be phytotoxic (cause foliar damage) to certain crops and/or cultivars. For example, many grape cultivars are sensitive to sulfur and/or copper. The **Relative Disease Susceptibility and Chemical Sensitivity among Grape Cultivars** table on page 163 lists cultivar sensitivity to these materials. Additionally, some grape cultivars are sensitive to certain strobilurin fungicides, and some strawberry cultivars are sensitive to Sinbar herbicide. Several apple cultivars are sensitive to azoxystrobin, the active ingredient in Abound, Quilt, and Quadris Top fungicides. Always read the comments associated with the materials in this guide.

Pesticide choices can be limited by cultivar, disease or insect pressure, and other factors. Grower preference, experience with materials, and price often influence decisions as well. Pest management in fruit crops is relatively easy as long as growers understand the pests, critical periods for control, proper selection of control materials, and proper application procedures.

Always read the entire pesticide label. If you have any questions about the proper use of a pesticide, refer to other sources, such as the *Midwest Small Fruit Pest Management Handbook* or *Midwest Tree Fruit Pest Management Handbook*. If you still have questions, contact the manufacturer or your state Extension specialist for clarification.

1. APPLE

Apple Insect Pests

The shaded boxes represent the crop stages where common pests in the Midwest are active and action (scouting and preventative sprays) may be necessary/recommended. Weather and degree day accumulation will impact the exact timing of pest appearance in the orchard. **MD signifies timing to set Mating Disruption traps.**

| Stage | | | | | | | | | |
|---------|-------------------|--------------------------------------|---------------|--------------------------------------|-------|--------------------------------------|----------------------------|-------------|-----------------------------|
| Dormant | Green Tip | Half-inch Green | Tight Cluster | Pink | Bloom | Petal Fall | First and Second Cover | Third Cover | Summer Cover |
| | | | | | CM-MD | | codling moth (CM) | | |
| | | | OFM-MD | | | oriental fruit moth (OFM) | | | |
| | | | | | | plum curculio | | | |
| | | | | | | | apple maggot | | |
| | | | | | | | brown marmorated stink bug | | |
| | European red mite | | | | | European red mite | | | |
| | | | | | | | | | Japanese beetle |
| | | rosy apple aphid | | | | rosy apple aphid | | | |
| | San Jose scale | | | | | | San Jose scale | | |
| | | | | | | | | | woolly apple aphid |
| | | | | | | | periodical cicada | | |
| | | | | | | | potato leafhopper | | |
| | | | | | | dogwood borer - MD | dogwood borer | | |
| | | | | | | | green apple aphid | | |
| | | | | | | leafrollers | | | |
| | | | | tarnished plant bug | | | | | |
| | | spotted tentiform leafminer - adults | | spotted tentiform leafminer - larvae | | spotted tentiform leafminer - larvae | | | spotted tentiform leafminer |
| | | ambrosia beetles | | | | | | | |

Apple Spray Schedule

How to read the spray schedule tables

Every apple growth stage has important notes on disease or insect management. In some cases, the reader will be directed to the special problems section at the end of the section or chapter. Please make sure to read thoroughly and contact your state Extension specialist with any specific questions.

Key to tables

E = excellent control

G = good control

F = fair control

[r] = fungicide/Insecticide resistance possible

s = suppression only

i = ineffective

u = unknown efficacy

x = pest not on the label

¹Efficacy data in this publication are based on trials conducted across various regions and does not necessarily reflect local efficacy differences or changes over time. However, results of investigations in local areas of the Midwest Region will be mentioned. Growers should contact their Extension specialist for the most recent or for state-specific information. The information on this publication is only a guide; the authors and their institutions assume no liability for practices implemented based on this information.

Always read and follow pesticide labels. The label is the law. Product registration may vary by state.

²FRAC/IRAC code represents the mode of action of the fungicide/insecticide.

³PHI refers to the pre-harvest interval, which is the number of days before harvest that the product may not be applied.

⁴All fungicides/insecticides have a Restricted-Entry Interval (REI). The restricted-entry interval is the time immediately after a pesticide application when entry into the treated area is limited. Check labels for REI. Restrictions in REI may prohibit the use of certain pesticides during harvest.

Applicators must abide by both maximum amount of product per season AND maximum number of applications.

⁵Max amt refers to the product's maximum amount/acre/year.

⁶Max app refers to the product's maximum number of applications per year.

RUP refers to restricted use pesticide.

Notes on disease management

The fungi that cause apple scab, powdery mildew, and cedar apple rust attack emerged leaves to a greater degree than older leaves. The fungi that cause summer rots attack developing fruit as early as petal fall, even though symptoms may not appear until later in the season.

To protect new growth and fruit, starting fungicide applications early to protect new growth is essential. That said, successful growers understand the limits of what fungicides can do, and they consider pesticide cost and the risk of disease when deciding which fungicide to use and when to use it.

With proper timing and application, captan, mancozeb, or captan plus mancozeb ("cptozeb") can provide very good to excellent scab control from green tip until pink, at a lower cost, and little risk of fungicide resistance. As always, the goal is to keep the number of primary scab lesions low to improve fruit protection later in the season. This is more difficult in cooler, wet years, which may require more frequent spraying.

At tight cluster through first cover (when the risk of powdery mildew, scab, and rust are highest), incorporate the broad-acting, systemic fungicides with the FRAC codes of 3, 7, 11, to improve management and best utilize these fungicides' systemic nature.

Following detection of resistance of apple scab fungus to strobilurins (FRAC 11), DMI (FRC 3), and Syllit fungicides in 2012 in Illinois, using mancozeb, Inspire Super and Fontelis from green tip to 10 days after petal fall provided excellent protection of foliage and fruit against scab, powdery mildew, and rust diseases of apples. Also, spraying crab apple trees at borders of commercial orchards is strongly recommended, as crab apples are very susceptible to scab fungus and they are infected before apple trees, so they provide spore showers (conidial inoculum) to commercial apple trees.

Apple Dormant to Silver Tip - Diseases

Apply before growth starts in spring and when temperatures are above 45°F.

- Make fixed copper applications at silver tip. Do not apply copper after 1/4-inch green leaf stage or when drying conditions are cool and slow, because that may cause severe injury. Many fixed copper fungicides/bactericides are registered for use on apple. Fixed coppers can be mixed with oil. However, never combine copper sulfate alone with dormant oil. Using copper at this stage does not eliminate the need of streptomycin at bloom. Use of copper at this stage has been shown to aid in the management of apple scab, particularly in

orchards that had a high incidence of the disease the previous season.

- Apply Ridomil to soil in early spring before growth starts.
- The suggested urea application rate is 40 lb. of agricultural grade urea (46-0-0) per 100 gal of water. See Sanitation Methods to Aid in Apple Scab Control, page 51.
- Using a forecasting program (e.g., MARYBLIT Program) to predict infection by fire blight bacterium is highly recommended. Such programs usually reduces the use of antibiotics for managing fire blight to one or two sprays.
- Recently, bacterial blister spot disease has been diagnosed in commercial apple orchards in the

Midwest. This disease is effectively managed by applications of streptomycin. So, using streptomycin for control of fire blight disease should be minimized (e.g., by using a fire blight prediction program such as MARYBIT).

- Bitter rot disease is a very serious threat to production of some apple cultivars, especially 'Honeycrisp' in the Midwest. Local research have shown the Merivon, captan, Aprovia, and Ziram provide effective management of this disease. Application of tank-mixed Merivon plus captan has provided the most effective control of the disease. Following recommendations on the labels, particularly on PHI, is very important.

Table 1-1. Apple Diseases - Dormant Through Silver Tip¹

| Product And Formulation Active Ingredient | FRAC Code ² | Fire Blight | Phytophthora Crown And Collar Rot | Scab | REI ³ PHI ⁴ | Max Amt ⁵ Max App ⁶ |
|--|------------------------|-------------------|--|----------------------------|--------------------------------------|--|
| Bordeaux mixture, 8-8-10 copper sulfate | M | 8 lb. s[G] | x | x | NA 24h | NA NA |
| Cuprofix Ultra 40 disperss copper sulfate | M | 5-7.5 lb. s[G] | x | x | NA 48h | 40 lb. NA |
| Kocide 3000 copper hydroxide | M | 3.5-7 lb. s[G] | x | x | NA 48h | 53.3 lb. NA |
| Ridomil Gold SL (SC) mefenoxam | | x | 2 qt./A (2.0) or 1.5 fl. oz./1,000 sq. ft. (see note above) | x | NA 48h | NA NA |
| urea (46-0-0) | NA | x | x | 44 lb. (see note above) | 0 | 50 lb. |
| | | x | x | G | 0 | NA |

Footnotes: See how to read the spray schedule tables section at the beginning of the chapter.

Phytophthora collar, crown and root rot of apple

Ridomil Gold SL is labeled for use on bearing apple trees. Make applications before symptoms appear, especially in areas of the orchard with poor water drainage. Ridomil Gold SL will not revitalize trees showing moderate to severe crown rot symptoms.

Mix 0.5 pint of Ridomil Gold SL with 100 gallons of water. Around the trunk of the tree, apply the amount of diluted mixture indicated in the table below. Make applications in early spring before growth starts and in the fall after harvest and before the ground freezes. On new plantings, delay the first application until two weeks after planting.

To determine trunk diameter, measure the trunk 12 inches above soil line.

Amount of Ridomil Gold SL (Diluted) To Apply For Rot Control

| Trunk Diameter | Quarts of Diluted Mixture/Tree |
|----------------|-----------------------------------|
| < 1 inch | 1 quart |
| 1-3 inches | 3 quarts |
| > 5 inches | 4 quarts |

Do not dip tree roots or spray bare roots with solutions containing Ridomil Gold SL. Do not graze in or feed cover crops from treated orchards. Illegal residues may occur.

Phosphorous acid (phosphonates and phosphites)

A number of phosphorous acid products are registered as fungicides to control root and collar rot

(caused by *Phytophthora* spp.) on apple, pear, and stone fruit. Brand names for these products include but are not limited to Agri-Fos, ProPhyt, Phostrol, and Rampart. Their active ingredient, phosphorous acid, is essentially the same active ingredient as in the fungicide Aliette, which has been registered for use on tree fruit for many years, however these are not generic versions of Aliette or each other. Care must be taken in following the label. In some instances, products are labeled for control of other diseases, although the data is complex and inconsistent.

These materials are applied as foliar sprays. The active ingredient is highly systemic and moves down the tree from the leaves into the crown and roots. See the label for current use recommendations. Although labeled, these products are not recommended for managing apple scab or fire blight in the Midwest.

For spring and summer *Phytophthora* collar, crown and root rot control on tree fruit: Under moderate disease pressure, apply products as indicated on the label on a 30-60-day spray interval. Make the first application in the spring after sufficient foliage is present to absorb the chemical.

Do not apply within two to three weeks of leaf color change in the fall. Foliage must be green and living for the roots to take up and transport Aliette.

Do not apply Aliette if you have applied copper-based fungicides within two weeks to avoid possible phytotoxic reactions.

Apple Green Tip To Half-Inch Green - Diseases

Begin fungicide sprays at green tip and repeat every 5-7 days through the first cover.

Notes on disease management

- **Initiate** applications at green tip or when environmental conditions are favorable for primary scab development. Continue applications through the duration of primary scab on a 7-10 day interval.
- Captan 80WG PLUS Mancozeb 75D F is a highly recommended tank mix (often called “captozebe”) and can be used up to 8 times, limited by the 77-day PHI for Mancozeb. See Note About Mancozeb (EBDC Products), page 50. If you are planning to use captan for control of summer fruit rots, you can consider application of mancozeb alternated with Fontelis and Inspire Super. Resistance of scab fungus to strobilurins (FRAC 11), DMI (FRC 3), and Syllit fungicides in the Midwest has been reported.
- Sulfur is formulated as dusts, liquids, and wettable powders (e.g., wettable sulfur, Microthiol Disperss, Cosavet, Microfine Kumulus, Liquid Sulfur Six, and Dusting Sulfur). Formulations can vary from 80% to 95% elemental sulfur. Formulations with finer particles are more effective. Sulfur is effective against plant-feeding mites but can damage predatory mite populations. Do not use within 10 days of applying oil or captan or when temperatures exceed 85°F. Certain apple varieties are sensitive to sulfur sprays under certain conditions. Do not apply unless the varieties are known to be sulfur tolerant.
- Vangard 75WG is most effective at temperatures below 70°F.

Table 1-2. Apple Diseases - Green Tip Through Half-Inch Green¹

| Product And Formulation | FRAC Code ² | Powdery Mildew | Scab | REI ³ PHI ⁴ | Max Amt ⁵ Max App ⁶ |
|----------------------------|------------------------|----------------|--------------|-----------------------------------|---|
| Active Ingredient | | | | | |
| Captan 80 WDG | M | x | 2.5-5 lb. | 24h | 40 lb. |
| captan | | x | G | 0d | NA |
| Cuprofix Ultra 40 disperss | M | x | 1-2.5 lb. | 12h | NA |
| copper sulfate | | x | F | NA | NA |
| Ferbam Granulfo (76WDG) | M | x | 3.5 lb. | 24h | NA |
| ferbam | | x | F | NA | 3 |
| Kocide 3000 | M | x | 0.75-7.0 lb. | 48h | 53.3 lb. |
| copper hydroxide | | x | F | 0d | NA |
| Microthiol Disperss | M | 10-20 lb. | 10-20 lb. | 24h | NA |
| sulfur | | F-G | i-F | 0d | NA |

(Continued)

Table 1-2. Apple Insects – Green Tip Through Half-Inch Green¹ (continued)

| Product And Formulation Active Ingredient | FRAC Code ² | Powdery Mildew | Scab | REI ³ PHI ⁴ | Max Amt ⁵ Max App ⁶ |
|--|---------------------------|-------------------|--------------|--------------------------------------|--|
| Roper DF Rainshield | M | x | 3 or 6 lb. | 24h | 21 lb. |
| mancozeb | | x | G | 77d | 6 |
| Scala SC | 9 | x | 7-10 fl. oz. | 12h | 40 fl. oz. |
| pyrimethanil | | x | G-E | 72d | NA |
| Vanguard WG | 9 | x | 5 oz. | 12h | 30 oz. |
| cyprodinil | | x | G | 0d | 2 |
| Ziram 76DF | M3 | x | 3 or 6 lb. | 48h | 42.4 lb. |
| ziram | | x | G | 14d | 7 |

Footnotes: See how to read the spray schedule tables section at the beginning of the chapter.

Apple Green Tip To Half-Inch Green – Insects

Notes on insect management

- Superior oil: Apply oil when temperature is above 32°F (for at least 24 hours before and after the application); never during freezing weather. Check label for fungicide/oil compatibility. Oil is most effective when sprayed dilute under calm conditions to ensure thorough coverage of all woody tissue. Delaying oil application until half-inch green controls **European red mite eggs** better than earlier applications. For European red mite, oil should be used at higher rates (1-2%) at this pre-bloom stage but reduced to 0.5% for summer sprays.
- Where **San Jose scale** is a main target of oil sprays, the best application timing is at green tip.
- Wait until half-inch green or pink if your primary target is European red mite or rosy apple aphid. Although Diazinon is labeled for use with oil to increase scale control, trials have shown that oil alone results in greater than 98 percent control of scales if coverage is thorough. Adding an insecticide does improve aphid control.
- Esteem 35WP controls scale anytime between half- inch green and second cover. At half-inch green it also controls **rosy apple aphid**. When used at pink it also controls **leafminer**. The minimum rate is effective when used pre-bloom, but the maximum rate is necessary if application is delayed until the crawler stage in early summer.
- **Oriental fruit moth:** Pheromone traps for oriental fruit moth should be deployed by half-inch green, with the first catch of moths expected at the pink stage.

Table 1-3. Apple Insects – Green Tip Through Half-Inch Green¹

| Product And Formulation Active Ingredient | IRAC Code ² | European Red Mite | Rosy Apple Aphid | San Jose Scale | REI ³ PHI ⁴ | Max Amt ⁵ Max App ⁶ |
|--|---------------------------|-------------------------|---------------------|-------------------|--------------------------------------|--|
| Acramite 50WS | 20D | 1 lb. | x | x | 12h | NA |
| bifenazate | | G | x | x | 7d | 1 |
| Actara (25WDG) | 4A | x | 4.5 oz. | x | 12h | 16.5 oz. |
| thiamethoxam | | x | E | x | See label | NA |
| Agri-Mek SC (0.7SC) (RUP) | 6 | 2.25-4.25 fl. oz. | x | x | 12h | 8.5 fl. oz. |
| abamectin | | G | x | x | 28d | 2 |
| Apollo SC (1SC) | 10A | 4-8 oz. | x | x | 12h | NA |
| clofentezine | | E | x | x | 45d | 1 |
| Asana XL (0.66EC) (RUP) | 3A | x | 4.8-14.5 fl. oz. | 4.8-14.5 fl. oz. | 12h | 101.5 fl. oz. |
| esfenvalerate | | x | G | i | 21d | NA |
| Assail 30SG | 4A | x | 32 fl. oz. | 8 oz. | 12h | 32 oz. |
| acetamiprid | | x | E | s | 7d | 4 |

(Continued)

Table 1-3. Apple Insects - Green Tip Through Half-Inch Green¹ (continued)

| Product And Formulation Active Ingredient | IRAC Code ² | European Red Mite | Rosy Apple Aphid | San Jose Scale | REI ³ PHI ⁴ | Max Amt ⁵ Max App ⁶ |
|--|---------------------------|-------------------------|---------------------|------------------------|--------------------------------------|--|
| Azera 0.21EC | 3A | x | 32 fl. oz. | 32 fl. oz. | 12h | NA |
| azadirachtin + pyrethrins | | x | u | u | 0d | 10 |
| Baythroid XL (1EC) (RUP) | 3A | x | x | 2.4-2.8 fl. oz. | 12h | 2.8 fl. oz. |
| beta-cyfluthrin | | x | x | G | 7d | NA |
| Belay (2.13SC) | 4A | x | 4-6 fl. oz. | 6 fl. oz. | 12h | 12 fl. oz. |
| clothianidin | | x | E | G | 7d | NA |
| Beleaf 50SG | 29 | x | 2-2.8 oz. | x | 12h | 8.4 oz. |
| flonicamid | | x | G | x | 21d | 3 |
| Centaur WDG (70WDG) | 16 | x | x | 34.5 oz. | 12h | 34.5 oz. |
| buprofezin | | x | x | E | 14d | 1 |
| Damoil | UN | 0.5-2% | 0.5-2% | 0.5-2% | NA | NA |
| mineral oil | | u | u | G | NA | NA |
| Danitol 2.4EC (RUP) | 3A | 16-21.3 fl. oz. | 10.6-21.3 fl. oz. | x | 24h | 42.7 fl. oz. |
| fenpropathrin | | F | F | x | 14d | NA |
| Delta Gold (1.5EC) (RUP) | 3A | x | x | 0.9-1.9 fl. oz. | 12h | 3.6 fl. oz. |
| deltamethrin | | x | x | u | 21d | NA |
| Diazinon AG600 | 1B | x | x | 12.75 fl. oz./100 gal. | 4d | 25.5 fl. oz. |
| diazinon | | x | x | x | 21d | 2 |
| Esteem 35WP | 7C | x | 3-5 oz. | 4-5 oz. | 12h | 10 oz. |
| pyriproxyfen | | x | E | E | 45d | 2 |
| Exirel (0.83SE) | 28 | x | 13.5-20.5 fl. oz. | x | 12h | 61.5 fl. oz. |
| cyantraniliprole | | x | E | x | 3d | 3 |
| Imidan 70W | 1B | x | x | 2.125-5.75 lb. | 4 or 7d | 15.5 lb. |
| phosmet | | x | x | F | 7d | NA |
| Kanemite 15SC | 20B | 21-31 fl. oz. | x | x | 12h | 62 fl. oz. |
| acequinocyl | | E | x | x | 14d | 2 |
| Lannate LV (2.4WSL) (RUP) | 1A | x | 1.5-3 pt. | x | 3d | 15 pt. |
| methomyl | | x | G | x | 14d | 5 |
| Mustang Maxx (0.83EC) (RUP) | 3A | x | 1.2-4 fl. oz. | x | 12h | 24 fl. oz. |
| zeta-cypermethrin | | x | u | x | 14d | NA |
| Nealta (1.67SC) | 25 | 13.7 fl. oz. | x | x | 12h | 274 fl. oz. |
| cyflumetofen | | E | x | x | 7d | 2 |
| Neemix 4.5 (0.39L) | UN | x | 5-7 fl. oz. | 7-16 fl. oz. | 12h | NA |
| azadirachtin | | x | F | u | 0d | NA |
| Nexter (SC) | 21A | 11-17 fl. oz. | x | x | 12h | NA |
| pyridaben | | G | x | x | 25d | 1 |
| Onager Optek (1EC) | 10 | 12-24 oz. | x | x | 12h | 24 oz. |
| hexythiazox | | E | x | x | 28d | 1 |

(Continued)

Table 1-3. Apple Insects - Green Tip Through Half-Inch Green' (continued)

| Product And Formulation Active Ingredient | IRAC Code ² | European Red Mite | Rosy Apple Aphid | San Jose Scale | REI ³ PHI ⁴ | Max Amt ⁵ Max App ⁶ |
|--|---------------------------|-------------------------|---------------------|-------------------|--------------------------------------|--|
| Permethrin 3.2EC (RUP) | 3A | x | 4-10 fl. oz. | x | 12h | 20 fl. oz. |
| permethrin | | x | G | x | See label | NA |
| Portal XLO (0.4EC) | 21A | 2 pt. | x | x | 12h | 2 pt. |
| fenpyroximate | | E | x | x | 14d | 1 |
| PQZ (1.87SC) | 9B | x | 2.4-3.2 fl. oz. | x | 12h | 4.8 fl. oz. |
| pyrifluquinazon | | x | E | x | 14d | 2 |
| Savey 50DF | 10A | 3-6 oz. | x | x | 12h | 6 oz. |
| hexythiazox | | E | x | x | 28d | 1 |
| Sevin XLR Plus | 1A | x | 1.5-3 qt. | 1.5-3 qt. | 12h | 15 qt. |
| carbaryl | | x | F | F | 3d | 8 |
| Sivanto Prime (1.67SC) | 4D | x | 7-14 fl. oz. | 10.5-14 fl. oz. | 4h | 28 fl. oz. |
| flupyradifurone | | x | G | G | 14d | NA |
| Vendex 50WP (RUP) | 12B | 1-2 lb. | x | x | 48h | 4 lb. |
| fenbutatin-oxide | | F | x | x | 14d | 2 |
| Versys Inscalis (0.83DC) | 9D | x | 1.5 fl. oz. | x | 12h | 7 fl. oz. |
| afidopyropen | | x | G | x | 7d | NA |
| Vydate L (2L) (RUP) | 1A | 2-4 pt. | 4-8 pt. | x | 48h | 8 pt. |
| oxamyl | | G | G | x | 14d | 4 |
| Warrior II (2.08CS) (RUP) | 3A | x | 1.2-2.5 fl. oz. | 1.2-2.5 fl. oz. | 24h | 12.8 fl. oz. |
| lambda-cyhalothrin | | x | G | i | 21d | NA |
| Zeal (72WP) | 10B | 2-3 oz. | x | x | 12h | 3 oz. |
| etoxazole | | E | x | x | 14d | 1 |

Footnotes: See how to read the spray schedule tables section at the beginning of the chapter.

Apple Tight Cluster To Pink - Diseases

7 days after half-inch green.

Notes on disease management

- Apple pink is a critical time for controlling scab, rust, and powdery mildew.
- See Note About Mancozeb (EBDC Products), page 50.
- Scala 5SC and Vangard are most effective at temperatures below 70°F.
- Topguard Fungicide Specialty Crops: Do not confuse with Topguard EQ, which contains azoxystrobin, and is phytotoxic on many apple varieties.
- Apply prohexidione-calcium (Pro-Ca; Apogee) at 1-3" growth and continue at one- to four-week intervals at the first sign of regrowth, to control shoot growth and reduce the risk of fire blight. See page 47 for more information regarding application of Pro-Ca for management of plant vigor and im-

proved resistance to fire blight. See generics table for a list of Pro-Ca products.

Fungicide resistance alert

- Mix FRAC 3, 7 and/or 11 fungicides with mancozeb or captan for resistance management.
- To limit the potential for fungicide resistance development, do not make more than four (4) applications of any fungicide within each group per season, and delay using them until pink (at the earliest) whenever possible.
- Do not make more than two sequential applications of any fungicide within each group without alternating to a fungicide from a different chemistry group.
- It is recommended that growers alternate between FRAC codes to reduce the risk of fungicide resistance. Rely on the information provided by your local extension specialist.

Table 1-4. Apple Diseases - Tight Cluster Through Pink¹

| Product And Formulation Active Ingredient | IRAC Code ² | Powdery Mildew | Rust | Scab | REI ³ PHI ⁴ | Max Amt ⁵ Max App ⁶ |
|--|---------------------------|-------------------|-----------------|-----------------|--------------------------------------|--|
| Aprovia (EC) | 7 | 5.5-7 fl. oz. | 5.5-7 fl. oz. | 5.5-7 fl. oz. | 12h | 27.6 fl. oz. |
| benzovindiflupyr | | F | u | G-E | 30d | NA |
| Captan 80 WDG | M | 2.5-5 lb. | x | 5 lb. | 24h | 40 lb. |
| captan | | i | x | G | 0d | NA |
| Cevya | 3 | 3-5 fl. oz. | 3-5 fl. oz. | 3-5 fl. oz. | 12h | NA |
| mefentrifluconazole | | G-E | E | E | 0d | NA |
| Cuprofix Ultra 40D | M | x | x | 1-2.5 lb. | 12hr | NA |
| copper sulfate | | x | x | F | NA | NA |
| Excalia (2.84 SC) | 7 | 3-4 fl. oz. | 3-4 fl. oz. | 3-4 fl. oz. | 12h | 8 fl. oz. |
| inpyrfluxam | | G-E | u | E | PF | 2 |
| Ferbam Granuflo (76 WDG) | M3 | x | 3.5 lb. | 3.5 lb. | 24h | NA |
| ferbam | | x | G | F | NA | 3 |
| Flint Extra | 11 | 2.5-2.9 fl. oz. | 2.5-2.9 fl. oz. | 2.5-2.9 fl. oz. | 12h | 10.5 fl. oz. |
| trifloxystrobin | | G [r] | F | G [r] | 14d | NA |
| Fontelis (1.67 SC) | 7 | 16-20 fl. oz. | 16-20 fl. oz. | 16-20 fl. oz. | 12h | 61 fl. oz. |
| penthiopyrad | | G | E | E | 28d | NA |
| Indar 2F | 3 | 6-8 fl. oz. | 6-8 fl. oz. | 6-8 fl. oz. | 12h | 32 fl. oz. |
| fenbuconazole | | E [r] | E | E [r] | 14d | 4 |
| Inspire Super (EW) | 3+9 | 12 fl. oz. | 12 fl. oz. | 12 fl. oz. | 12h | 60 fl. oz. |
| difenoconazole + cyprodinil | | G | E | E | 28d | NA |
| Kenja 400SC | 7 | 12.5 fl. oz. | x | 12.5 fl. oz. | 12h | NA |
| isofetamid | | s | x | F | 20d | NA |
| Luna Privilege (SC) | 7 | 2.4-6.8 fl. oz. | x | 4-6.8 fl. oz. | NA | NA |
| fluopyram | | G | x | G-E | NA | NA |
| Luna Sensation (SC) | 7+11 | 5-5.8 fl. oz. | 4-5.8 fl. oz. | 4-5.8 fl. oz. | 12h | 21 fl. oz. |
| fluopyram + trifloxystrobin | | G [r] | F | E [r] | 14d | 4 |
| Luna Tranquility (SC) | 7+9 | 11.2-16 fl. oz. | x | 11.2-16 fl. oz. | 12h | 54.7 fl. oz. |
| fluopyram + pyrimethanil | | G | x | E | 72d | NA |
| Merivon (2.09SC) | 7+11 | 4-5.5 fl. oz. | 4-5.5 fl. oz. | 4-5.5 fl. oz. | 12h | 22 fl. oz. |
| fluxapyroxad + pyraclostrobin | | G | s | E | 0d | 4 |
| Microthiol Disperss | M | 10-20 lb. | x | 10-20 lb. | 24h | NA |
| sulfur | | G | x | i-F | 0d | NA |
| Miravis (1.67 SC) | 7 | 3.4 fl. oz. | 3.4 fl. oz. | 3.4 fl. oz. | 4h | 13.6 fl. oz. |
| pydiflumetofen | | G | G | G | 30d | 4 |
| Omega 500F | 29 | x | 13.8 fl. oz. | 10-13.8 fl. oz. | 12h | 138 fl. oz. |
| fluazinam | | x | s-G | G | 28d | 10 |
| OSO 5% | 19 | 3.75-13 fl. oz. | x | 3.75-13 fl. oz. | 4h | 78 oz. |
| polyoxin D | | F | x | F | 0d | 6 |
| Pristine | 11+7 | 14.5-18.5 oz. | 14.5-18.5 oz. | 14.5-18.5 oz. | 12h | 74 oz. |
| pyraclostrobin + boscalid | | E [r] | E | E [r] | 0d | 4 |

(Continued)

Table 1-4. Apple Diseases – Tight Cluster Through Pink¹ (continued)

| Product And Formulation Active Ingredient | IRAC Code ² | Powdery Mildew | Rust | Scab | REI ³ PHI ⁴ | Max Amt ⁵ Max App ⁶ |
|--|---------------------------|-------------------|--------------|-----------------|--------------------------------------|--|
| Procure 480 SC | 3 | 8-16 fl. oz. | 8-16 fl. oz. | 8-16 fl. oz. | 12h | 64 fl. oz. |
| triflumizole | | E [r] | G [r] | G [r] | 14d | NA |
| Rally 40WSP | 3 | 5-10 oz. | 5-8 oz. | 5-8 oz. | 24h | 5 lb. |
| myclobutanil | | E [r] | F | E-G[r] | 14d | NA |
| Roper DF rainshield | M | 6 lb. | 6 lb. | 6 lb. | 24h | 21 lb. |
| mancozeb | | i | G | G | 77d | 6 |
| Scala (SC) | 9 | x | x | 7-10 fl. oz. | 12h | 40 fl. oz. |
| pyrimethanil | | x | x | G-E | 72d | NA |
| Sercadis | 7 | 3.5-4.5 fl. oz. | 4.5 fl. oz. | 3.5-4.5 fl. oz. | 12h | 18 fl. oz. |
| fluxapyroxad | | G | s | G-E | 0d | 4 |
| Sovran (50WG) | 11 | 4-6.4 oz. | 3.2-6.4 oz. | 3.2-6.4 oz. | 12h | 25.6 oz. |
| kresoxim-methly | | G [r] | G | G [r] | 30d | 4 |
| Syllit FL | U12 | x | x | 1.5 pt. | 48h | 3 pt. |
| dodine | | x | x | E [r] | pink | 2 |
| Topguard Specialty Crops (SC) | 3 | 8-12 fl. oz. | 8-12 fl. oz. | 13 fl. oz. | 12h | 52 fl. oz. |
| flutriafol | | E | E | G | 14d | 4 |
| Topsin-M WSB | 1 | 0.75-1 lb. | x | 0.75-1 lb. | 24h | 4 lb. |
| thiophanate methyl | | G [r] | x | i | 1d | NA |
| Torino (SC) | U6 | 6.8 oz. | x | x | 4h | 6.8 oz. |
| cyflufenamid | | G | x | x | 14d | 1 |
| Vanguard WG (75WG) | 9 | x | x | 5 oz | 12h | 30 oz. |
| cyprodinil | | x | x | G | 0d | 2 |
| Ziram 76DF | M3 | x | 6 lb. | 6 lb. | 48h | 42.4 lb. |
| ziram | | x | G | G | 14d | 7 |

Footnotes: See how to read the spray schedule tables section at the beginning of the chapter.

Apple Tight Cluster To Pink – Insects

Notes on insect management

- **Rosy apple aphid:** Scout for curled leaves at early pink. Apply aphicide at pink if you find any curled leaves with rosy apple aphid inside.
- **San Jose scale:** Put pheromone traps in place now to monitor adult scales; expect crawlers four to six weeks after adults emerge.
- **Pyrethroids** (Asana, Baythroid, Danitol, Mustang Maxx, Permethrin, and Warrior) kill predatory mites that feed on European red mite and two-spotted spider mite, thereby triggering outbreaks of these pests. Use pyrethroids only if the potential for plant bug and stink bug damage is high.
- **Oriental fruit moth:** If this pest is to be managed by pheromone mating disruption, then dispensers should be deployed at pink. Table 7-1. Products for Pheromone Mating Disruption of Codling Moth, Oriental Fruit Moth, and Dogwood Borer page 26-27.

Table 1-5. Apple Insects – Tight Cluster Through Pink¹

| Product And Formulation Active Ingredient | IRAC Code ² | Rosy Apple Aphid | REI ³ PHI ⁴ | Max Amt ⁵ Max App ⁶ |
|--|---------------------------|-------------------|--------------------------------------|--|
| Actara (25WDG) | 4A | 4.5 oz. | 12h | 16.5 oz. |
| thiamethoxam | | E | 35d | NA |
| Asana XL (0.66EC) (RUP) | 3A | 4.8-14.5 fl. oz. | 12h | 101.5 fl. oz. |
| esfenvalerate | | G | 21d | NA |
| Assail 30SG | 4A | 2.5- 4 oz. | 12h | 32 oz. |
| acetamiprid | | E | 7d | 4 |
| Azera 0.21EC | 3A | 32 fl. oz. | 12h | NA |
| azadirachtin + pyrethrins | | u | 0d | 10 |
| Belay (2.13SC) | 4A | 4-6 fl. oz. | 12h | 12 fl. oz. |
| clothianidin | | E | 7d | NA |
| Beleaf 50SG | 29 | 2-2.8 oz. | 12h | 8.4 oz. |
| flonicamid | | G | 21d | 3 |
| Danitol 2.4EC (RUP) | 3A | 10.6-21.3 fl. oz. | 24h | 42.7 fl. oz. |
| fenpropathrin | | F | 14d | NA |
| Esteem 35WP | 7C | 3-5 oz. | 12h | 10 oz. |
| pyriproxifen | | E | 45d | 2 |
| Exirel (0.83SE) | 28 | 13.5-20.5 fl. oz. | 12h | 61.5 fl. oz. |
| cyantraniliprole | | E | 3d | 3 |
| Lannate LV (2.4WSL) (RUP) | 1A | 1.5-3 pt. | 3d | 15 pt. |
| methomyl | | G | 14d | 5 |
| Mustang Maxx (0.83EC) (RUP) | 3A | 1.2-4 fl. oz. | 12h | 24 fl. oz. |
| zeta-cypermethrin | | u | 14d | NA |
| Neemix 4.5 (0.39L) | UN | 5-7 fl. oz. | 12h | NA |
| azadirachtin | | F | 0d | NA |
| Permethrin 3.2EC (RUP) | 3A | 4-10 fl. oz. | 12h | 20 fl. oz. |
| permethrin | | G | See label | NA |
| PQZ (1.87SC) | 9B | 2.4-3.2 fl. oz. | 12h | 4.8 fl. oz. |
| pyrifluquinazon | | E | 14d | 2 |
| Sevin XLR Plus | 1A | 1.5-3 qt. | 12h | 15 qt. |
| carbaryl | | F | 3d | 8 |
| Sivanto Prime (1.67SC) | 4D | 7-14 fl. oz. | 4h | 28 fl. oz. |
| flupyradifurone | | G | 14d | NA |
| Versys Inscalis (0.83DC) | 9D | 1.5 fl. oz. | 12h | 7 fl. oz. |
| afidopyropen | | G | 7d | NA |
| Vydate L (2L) (RUP) | 1A | 4-8 pt. | 48h | 8 pt. |
| oxamyl | | G | 14d | 4 |
| Warrior II (2.08CS) (RUP) | 3A | 1.2-2.5 fl. oz. | 24h | 12.8 fl. oz. |
| lambda-cyhalothrin | | G | 21d | NA |

Footnotes: See how to read the spray schedule tables section at the beginning of the chapter.

Apple Pink

7-10 days after tight cluster.

| Pest/Problem | Material | Rate/Acre | Comments |
|----------------|---|-----------|---|
| nutrient level | Solubor (boron) | 2 lb. | May add Solubor to pesticide solutions, but check for compatibility, order of mixing, etc. Solubor helps prevent cork spot; see page 31 for more information. Can add urea to pesticide sprays when needed. |
| | AND/OR feed-grade urea (nitrogen) | 3 lb. | |

Apple Bloom - Diseases

7 days after pink. Bloom begins when the first blossom opens (King Bloom).

- Fungicide applications for effective bitter rot control begins now.

Phytotoxicity Alert! From bloom to first cover, be aware of potential phytotoxicity issues with complex tank-mixes involving captan and other fungicides and insecticides.

Streptomycin is the most effective antibiotic for fire blight control. If streptomycin resistance has been confirmed in your orchard, switch to Kasumin 2L in those orchards with a documented streptomycin resistance.

- Fire blight (blossom blight):** Start fire blight sprays at first sign of open blossoms. Repeat sprays at 4- to 5-day intervals through bloom and petal fall on susceptible varieties. Using a disease warning program such as MARYBLYT is highly recommended.
- Do not concentrate Regulaid for fire blight control.
- Fire blight (shoot blight):** Apply the growth regulator prohexidione-calcium (Pro-Ca; Apogee) PLUS Regulaid at petal fall on king blooms for maximum effectiveness. It will take 10 days to two weeks after application for plants to be less susceptible to disease. See comments on page 49. Excessive nitrogen fertilization will make trees more susceptible to fire blight. See generics table for a list of Pro-Ca products.

Table 1-6. Apple Diseases - Bloom¹

| Product And Formulation Active Ingredient | FRAC Code ² | Bitter Rot | Fire Blight | Powdery Mildew | Rust | Scab | Summer Rot (Black And White Rot) | REI ⁴ PHI ³ | Max Amt ⁵ Max App ⁶ |
|--|------------------------|---------------|-------------|----------------|---------------|---------------|----------------------------------|-----------------------------------|---|
| Agri-Mycin 17 | 25 | x | 24-48 oz. | x | x | x | x | 12h | NA |
| streptomycin sulfate | | x | G [r] | x | x | x | x | 50d | NA |
| Aprovia (EC) | 7 | 5.5-7 fl. oz. | x | 5.5-7 fl. oz. | 5.5-7 fl. oz. | 5.5-7 fl. oz. | 5.5-7 fl. oz. | 12h | 276 fl. oz. |
| benzovindiflupyr | | G-E | x | F | u | G | G | 30d | NA |
| Captan 80 WDG | M | 2.5-5 lb. | x | 2.5-5 lb. | x | 5 lb | 2.5-5 lb. | 24h | 40 lb. |
| captan | | E* | x | i | x | G | E | 0d | NA |
| Cevya | 3 | x | x | 3-5 fl. oz. | 3-5 fl. oz. | 3-5 fl. oz. | 3-5 fl. oz. | 12h | NA |
| mefentrifluconazole | | x | x | G-E | E | E | F-G | 0d | NA |
| Cuprofix Ultra 40D | M | x | 5.5-75 lb. | x | x | 1-2.5 lb. | x | 12hr | NA |
| copper sulfate | | x | F-G | x | x | F | x | NA | NA |
| Excalia (2.84 SC) | 7 | 3-4 fl. oz. | x | 3-4 fl. oz. | 3-4 fl. oz. | 3-4 fl. oz. | x | 12h | 8 fl. oz. |
| inpyrfluxam | | u | x | G-E | u | E | x | PF | 2 |

(Continued)

Table 1-6. Apple Diseases - Bloom¹ (continued)

| Product And Formulation Active Ingredient | FRAC Code ² | Bitter Rot | Fire Blight | Powdery Mildew | Rust | Scab | Summer Rot (Black And White Rot) | REI ⁴ PHI ³ | Max Amt ⁵ Max App ⁶ |
|--|---------------------------|------------------|----------------|--------------------|--------------------|--------------------|---|--------------------------------------|--|
| Ferbam Granuflo (76 WDG) | M3 | 3.5 lb. | x | x | 3.5 lb. | 3.5 lb. | 3.5 lb. | 24h | NA |
| ferbam | | F | x | x | G | F | G | NA | 3 |
| Flint Extra | 11 | 2.9 fl. oz. | x | 2.5-2.9 fl. oz. | 2.5-2.9 fl. oz. | 2.5-2.9 fl. oz. | 2.9 fl. oz. | 12h | 10.5 fl. oz. |
| trifloxystrobin | | F-G | x | G [r] | F | E [r] | G | 14d | NA |
| Fontelis (1.67 SC) | 7 | x | x | 16-20 oz. | 16-20 fl. oz. | 16-20 fl. oz. | 16-20 fl. oz. | 12h | 61 fl. oz. |
| penthiopyrad | | x | x | G | E | E | u | 28d | NA |
| Indar 2F | 3 | x | x | 6-8 oz. | 6-8 fl. oz. | 6-8 fl. oz. | x | 12h | 32 fl. oz. |
| fenbuconazole | | x | x | E [r] | E | E [r] | x | 14d | 4 |
| Inspire Super (EW) | 3+9 | x | x | 12 fl. oz. | 12 fl. oz. | 12 fl. oz. | x | 12h | 60 fl. oz. |
| difenoconazole + cyprodinil | | x | x | F | E | E | x | 28d | NA |
| Kasumin 2L | 24 | x | 64 fl. oz. | x | x | x | x | 12h | 256 fl. oz. |
| kasugamycin | | x | G | x | x | x | x | 90d | 4 |
| Kenja 400SC | 7 | x | x | 12.5 oz. | x | 12.5 oz. | x | 12h | NA |
| isofetamid | | x | x | s | x | F | x | 20d | NA |
| kudos 28.5 WDG | PGR | x | See Label | x | x | x | x | 12h | See Label |
| prohexadione calcium | | x | G | x | x | x | x | 45d | NA |
| Luna Privilege (SC) | 7 | x | x | 2.4-6.8 fl. oz. | x | 4-6.8 fl. oz. | x | NA | NA |
| fluopyram | | x | x | G | x | G-E | x | NA | NA |
| Luna Sensation (SC) | 7+11 | 4-5.8 fl. oz. | x | 5-5.8 fl. oz. | 4-5.8 fl. oz. | 4-5.8 fl. oz. | 4-5.8 fl. oz. | 12h | 21 fl. oz. |
| fluopyram + trifloxystrobin | | E | x | G [r] | F | E [r] | E | 14d | 4 |
| Luna Tranquility (SC) | 7+9 | x | x | 11.2-16 fl. oz. | x | 11.2-16 fl. oz. | x | 12h | 54.7 fl. oz. |
| fluopyram + pyrimethanil | | x | x | G | x | E | x | 72d | NA |
| Manzate Pro Stik | M | x | x | 6 lb. | 6 lb. | 6 lb. | 3 lb. | 24h | 21 lb. |
| mancozeb | | x | x | i | G | E | G | 77d | 6 |
| Merivon (2.09SC) | 7+11 | 4-5.5 fl. oz. | x | 4-5.5 fl. oz. | 4-5.5 fl. oz. | 4-5.5 fl. oz. | 4-5.5 fl. oz. | 12h | 22 fl. oz. |
| fluxapyroxad + pyraclostrobin | | E | x | G | s | E | E | 0d | 4 |
| Microthiol Disperss | M | x | x | 10-20 lb. | x | 10-20 lb. | x | 24h | NA |
| sulfur | | x | x | G | x | i-F | x | 0d | NA |
| Miravis (1.67 SC) | 7 | 3.4 fl. oz. | x | 3.4 fl. oz. | 3.4 fl. oz. | 3.4 fl. oz. | 3.4 fl. oz. | 4h | 13.6 fl. oz. |
| pydiflumetofen | | s-F | x | G | G | G | s | 30d | 4 |
| Mycoshield | 41 | x | 1 lb. | x | x | x | x | 12h | 9 lb |
| oxytetracycline | | x | F-G | x | x | x | x | 60d | 6 |

(Continued)

Table 1-6. Apple Diseases – Bloom¹ (continued)

| Product And Formulation Active Ingredient | FRAC Code ² | Bitter Rot | Fire Blight | Powdery Mildew | Rust | Scab | Summer Rot (Black And White Rot) | REI ⁴ PHI ³ | Max Amt ⁵ Max App ⁶ |
|--|---------------------------|----------------|----------------|-------------------|-----------------|-------------------|---|--------------------------------------|--|
| Omega 500F | 29 | 13.8 fl. oz. | x | x | 13.8 fl. oz. | 10-13.8 fl. oz. | 13.8 fl. oz. | 12h | 138 fl. oz. |
| fluazinam | | F | x | x | s-G | G | s-G | 28d | 10 |
| OSO 5% | 19 | x | x | 3.75-13 fl. oz. | x | 3.75-13 fl. oz. | x | 4h | 78 oz. |
| polyoxin D | | x | x | F | x | F | x | 0d | 6 |
| Polyram 80 DF | M3 | x | x | x | 3 lb | 3 lb. | x | 24h | 21 lb. |
| metiran | | x | x | x | G | G | x | 77d | 7 |
| Pristine | 11+7 | 14.5-18.5 oz. | x | 14.5-18.5 oz. | 14.5-18.5 oz. | 14.5-18.5 fl. oz. | 14.5-18.5 oz. | 12h | 74 oz. |
| pyraclostrobin + boscalid | | F-G | x | E [r] | E | E [r] | E | 0d | 4 |
| Procure 480 SC | 3 | x | x | 8-16 fl. oz. | 8-16 fl. oz. | 8-16 fl. oz. | x | 12h | 64 fl. oz. |
| triflumizole | | x | x | E [r] | E [r] | G [r] | x | 14d | NA |
| Rally 40WSP | 3 | x | x | 5-10 oz. | 5-8 oz. | 5-8 oz. | x | 24h | 5 lb. |
| myclobutanil | | x | x | E [r] | F | E-G[r] | x | 14d | NA |
| Roper DF Rainshield | M | 3 lb. | x | 6 lb. | 6 lb. | 6 lb. | 3 lb. | 24h | 21 lb. |
| mancozeb | | see note above | x | i | G | E | see note above | 77d | 6 |
| Scala (SC) | 9 | x | x | x | x | 7-10 fl. oz. | x | 12h | 40 fl. oz. |
| pyrimethanil | | x | x | x | x | G-E | x | 72d | NA |
| Sercadis | 7 | x | x | 3.5-4.5 fl. oz. | 4.5 fl. oz. | 3.5-4.5 fl. oz. | 4.5 fl. oz. | 12h | 18 fl. oz. |
| fluxapyroxad | | x | x | G | s | G-E | F | 0d | 4 |
| Sovran (50WG) | 11 | x | x | 4-6.4 oz. | 3.2-6.4 fl. oz. | 3.2-6.4 oz. | 4-6.4 oz. | 12h | 25.6 oz. |
| kresoxim-methyl | | x | x | G [r] | E | E [r] | G | 30d | 4 |
| Syllit FL | U12 | x | x | x | x | 1.5 pt. | x | 48h | 3 pt. |
| dodine | | x | x | x | x | E [r] | x | pink | 2 |
| Topguard Specialty Crops (SC) | 3 | 13 fl. oz. | x | 8-12 fl. oz. | 8-12 fl. oz. | 13 fl. oz. | 13 fl. oz. | 12h | 52 fl. oz. |
| flutriafol | | u | x | E | E | G | u | 14d | 4 |
| Topsin-M WSB | 1 | x | x | 0.75-1 lb. | x | 0.75-1 lb. | 0.75-1 lb. | 24h | 4 lb |
| thiophanate methyl | | x | x | G [r] | x | i | G | 1d | NA |
| Torino (SC) | U6 | x | x | 6.8 oz. | x | x | x | 4h | 6.8 oz. |
| cyflufenamid | | x | x | E | x | x | x | 14d | 1 |
| Vanguard WG (75WG) | 9 | x | x | x | x | 5 oz. | x | 12h | 30 oz. |
| cyprodinil | | x | x | x | x | G | x | 0d | 2 |
| Ziram 76DF | M3 | 6 lb. | x | x | 6 lb. | 6 lb. | 6 lb. | 48h | 42.4 lb. |
| ziram | | G | x | x | G | G | i | 14d | 7 |

Footnotes: See how to read the spray schedule tables section at the beginning of the chapter.

(Continued)

Apple Bloom - Insects

7-10 days after pink. Bloom begins when the first blossom opens (*king bloom*). Remember to protect pollinators!

- **SAVE THE BEES!** Do not use insecticides or miticides at bloom.
- **Codling moth** (monitoring): Put out pheromone traps now to monitor adult codling moth activity. 2 per 5 acres, minimum of 2 per block. See page 34 for information about how to use traps to determine optimal insecticide timing.
- **Codling moth** (management with mating disruption): Deploy pheromone dispensers at bloom. See details below.

Mating disruption for codling moth, oriental fruit moth, and dogwood borer control

Several products are registered in some states for control of codling moth (CM), oriental fruit moth (OFM),

and dogwood borer (DWB) using the tactic of pheromone mating disruption. These products dispense a species-specific sex attractant that does not kill moths but prevents male moths from locating females for mating, which results in elimination of egg-laying in fruit. Mating disruption is most likely to succeed in blocks of at least 5 acres where initial populations of codling moth are low. If you attempt a mating disruption program in blocks smaller than 5 acres, or where infestation is moderate or high, then you also need to make border sprays or at least one or two insecticide cover sprays. Controlling these moths by mating disruption does not control other insect pests that insecticide applications manage, for example, plum curculio and apple maggot. Many of these products are deployed manually but aerosol emitters and sprayable products are available. The manual products last for several months, while the sprayables last for several weeks.

Table 1-7. Products For Pheromone Mating Disruption Of Codling Moth, Oriental Fruit Moth, And Dogwood Borer

| Target | Type | Product Name | Rate | Duration |
|--|--------------------------|--|--|--------------|
| Codling moth (CM) only | manual dispensers | CheckMate CM-XL 2.0 Dispenser | 120-200 dispensers/A | |
| | | CideTrak CMDA Combo Meso-A | 18-36 dispensers/A | 120-150 days |
| | | Isomate CM Flex | 200-400 dispensers/A | |
| | | Isomate C Plus | 200-400 dispensers/A | |
| | | Isomate C TT | 100-200 dispensers/A | |
| | | NoMate CM Spiral | 300-400 dispensers/A | |
| | sprayable | CheckMate CM 2.0 Flowable | 2.4-4.8 fl. oz./A | |
| | | NoMate CM MEC | 1.34-2.68 fl. oz./A | |
| | aerosol emitters | CheckMate Puffer CM-0 | 1-2 puffer cabinets/A | |
| | | Isomate CM Mist Plus | 1-2 units/A | |
| | | NoMate CM Smart Release | 1-2 units/A | 160 days |
| | amorphous polymer matrix | SPLAT Cydia V2 | 750-2000 grams/A; for 1 kg. as 400 point sources, apply 2.5-gram dollops; for 1 kg. as 1,000 point sources, apply 1-gram dollops | |
| Oriental fruit moth (OFM) only | manual dispensers | CheckMate OFM Dispenser | 100-200 dispensers/A | |
| | | CideTrak OFM-L Meso | 18-35 dispensers/A of one type + 100- 200 dispensers/A of other type | 180 days |
| | | Isomate M Rosso | 100-200 dispensers/A | |
| | | Isomate OFM TT | 100 dispensers/A | |
| | | NoMate OFM Spiral | 100-400 dispensers/A | |
| Oriental fruit moth (OFM) only <i>continued</i> | sprayable | CheckMate OFM-F | 1.32-2.93 fl. oz./A | |
| | aerosol emitters | CheckMate Puffer OFM-0 | 1-2 puffer cabinets/A | |
| | amorphous polymer matrix | SPLAT OFM 30M-1 | 400-1000 grams/A; for 1 kg. as 400 point sources, apply 2.5-gram (1/2 teaspoon) dollops; for 1 kg. as 1,000 point sources, apply 1-gram (1/4 teaspoon) dollops | |
| Both CM and OFM | manual dispensers | CideTrak CM-OFM Combo | 200-440 dispensers/A | 120 days |
| | | CideTrak CMDA+OFM Meso | 30-38 dispensers/A | 150-180 days |
| | | Isomate CM/OFM TT | 200 dispensers/A | |
| | aerosol emitters | CheckMate Puffer CM-OFM Pro | 1-2 puffer cabinets/A | |
| | | Isomate CM/OFM Mist Plus | 1-2 units/A | |
| Dogwood Borer (DWB) | manual dispensers | Isomate® DWB *check state availability | 100-200 dispensers/A | |

Apple Petal Fall - Diseases

7-10 days after bloom.

Notes on disease management

- **Bitter rot and other summer rots:** Applying mancozeb at this time using the label's extended program for control of scab provides collateral control of bitter and summer rot, and saves on applications of captan. Local research in the Midwest has shown that applications of Merivon plus captan are

most effective for control of bitter rot. Also, Aprovia is also effective for control of bitter rot and other summer diseases. Follow label guidelines.

- Excalia may not be applied after petal fall.
- Be aware that captan products may pose a risk of phytotoxicity in complex tank mixes at this stage.
- **Fire blight:** Continue antibiotic sprays on susceptible varieties until all petals have fallen.

Table 1-8. Apple Diseases - Petal Fall¹

| Product And Formulation Active Ingredient | FRAC Code ² | Bitter Rot | Fire Blight | Powdery Mildew | Rust | Scab | Summer Rots | REI ³ PHI ⁴ | Max Amt ⁵ Max App ⁶ |
|--|---------------------------|------------------|----------------|--------------------|--------------------|--------------------|------------------|--------------------------------------|--|
| Agri-Mycin 17 | 25 | x | 24-48 oz. | x | x | x | x | 12h | NA |
| streptomycin sulfate | | x | G [r] | x | x | x | x | 50d | NA |
| Aprovia (EC) | 7 | 5.5-7 fl. oz. | x | 5.5-7 fl. oz. | 5.5-7 fl. oz. | 5.5-7 fl. oz. | 5.5-7 fl. oz. | 12h | 276 fl. oz. |
| benzovindiflupyr | | G-E | x | F | u | G-E | F | 30d | NA |
| Captan 80 WDG | M | 2.5-5 lb. | x | 2.5-5 lb. | x | 5 lb. | 2.5-5 lb. | 24h | 40 lb. |
| captan | | E | x | i | x | G | E | 0d | NA |
| Cevya | 3 | x | x | 3-5 fl. oz. | 3-5 fl. oz. | 3-5 fl. oz. | 3-5 fl. oz. | 12h | NA |
| mefentrifluconazole | | x | x | G-E | E | E | F-G | 0d | NA |
| Cuprofix Ultra 40D | M | x | 5.5-7.5 lb. | x | x | 1-2.5 lb. | x | 12hr | NA |
| copper sulfate | | x | F-G | x | x | F-G | x | NA | NA |
| Excalia (2.84 SC) | 7 | 3-4 fl. oz. | x | 3-4 fl. oz. | 3-4 fl. oz. | 3-4 fl. oz. | x | 12h | 8 fl. oz. |
| inpyrfluxam | | u | x | G-E | u | E | x | PF | 2 |
| Ferbam Granuflo (76 WDG) | M3 | 3.5 lb. | x | x | 3.5 lb. | 3.5 lb. | 3.5 lb. | 24h | NA |
| ferbam | | F | x | x | G | F | G | NA | 3 |
| Flint Extra | 11 | 2.9 fl. oz. | x | 2.5-2.9 fl. oz. | 2.5-2.9 fl. oz. | 2.5-2.9 fl. oz. | 2.9 fl. oz. | 12h | 10.5 fl. oz. |
| trifloxystrobin | | G | x | G [r] | F | E [r] | G | 14d | NA |
| Fontelis (1.67 SC) | 7 | x | x | 16-20 fl. oz. | 16-20 fl. oz. | 16-20 fl. oz. | 16-20 fl. oz. | 12h | 61 fl. oz. |
| penthiopyrad | | x | x | G | E | E | u | 28d | NA |
| Indar 2F | 3 | x | x | 6-8 fl. oz. | 6-8 fl. oz. | 6-8 fl. oz. | x | 12h | 32 fl. oz. |
| fenbuconazole | | x | x | E [r] | E | E [r] | x | 14d | 4 |
| Inspire Super (EW) | 3+9 | x | x | 12 fl. oz. | 12 fl. oz. | 12 fl. oz. | x | 12h | 60 fl. oz. |
| difenoconazole + cyprodinil | | x | x | F | E | E | x | 28d | NA |
| Kasumin 2L | 24 | x | 64 fl. oz. | x | x | x | x | 12h | 256 fl. oz. |
| kasugamycin | | x | G | x | x | x | x | 90d | 4 |
| Kenja 400SC | 7 | x | x | 12.5 oz. | x | 12.5 oz. | x | 12h | NA |
| isofetamid | | x | x | s | x | F | x | 20d | NA |
| kudos 27WDG | PGR | x | See Label | x | x | x | x | 12h | See Label |
| prohexadione calcium | | x | E | x | x | x | x | 45h | NA |

(Continued)

Table 1-8. Apple Diseases - Petal Fall¹ (continued)

| Product And Formulation Active Ingredient | FRAC Code ² | Bitter Rot | Fire Blight | Powdery Mildew | Rust | Scab | Summer Rots | REI ³ PHI ⁴ | Max Amt ⁵ Max App ⁶ |
|--|---------------------------|-------------------|----------------|--------------------|--------------------|----------------------|-------------------|--------------------------------------|--|
| Luna Privilege (SC) | 7 | x | x | 2.4-6.8 fl. oz. | x | 4-6.8 fl. oz. | x | NA | NA |
| fluopyram | | x | x | G | x | G-E | x | NA | NA |
| Luna Sensation (SC) | 7+11 | 4-5.8 fl. oz. | x | 5-5.8 fl. oz. | 4-5.8 fl. oz. | 4-5.8 fl. oz. | 4-5.8 fl. oz. | 12h | 21 fl. oz. |
| fluopyram + trifloxystrobin | | E | x | G [r] | F | E [r] | E | 14d | 4 |
| Luna Tranquility (SC) | 7+9 | x | x | 11.2-16 fl. oz. | x | 11.2-16 fl. oz. | x | 12h | 54.7 fl. oz. |
| fluopyram + pyrimethanil | | x | x | G | x | E | x | 72d | NA |
| Merivon (2.09SC) | 7+11 | 4-5.5 fl. oz. | x | 4-5.5 fl. oz. | 4-5.5 fl. oz. | 4-5.5 fl. oz. | 4-5.5 fl. oz. | 12h | 22 fl. oz. |
| fluxapyroxad + pyraclostrobin | | E | x | G | s | E | E | 0d | 4 |
| Microthiol Disperss | M | x | x | 10-20 lb. | x | 10-20 lb. | x | 24h | NA |
| sulfur | | x | x | G | x | i-F | x | 0d | NA |
| Miravis (1.67 SC) | 7 | 3.4 fl. oz. | x | 3.4 fl. oz. | 3.4 fl. oz. | 3.4 fl. oz. | 3.4 fl. oz. | 4h | 13.6 fl. oz. |
| pydiflumetofen | | s-F | x | G | G | E | s | 30d | 4 |
| Mycoshield | 41 | x | 1 lb. | x | x | x | x | 12h | 9 lb. |
| oxytetracycline | | x | G | x | x | x | x | 60d | 6 |
| Omega 500F | 29 | 13.8 fl. oz. | x | x | 13.8 fl. oz. | 10-13.8 fl. oz. | 13.8 fl. oz. | 12h | 138 fl. oz. |
| fluazinam | | F | x | x | s-G | G | s-G | 28d | 10 |
| OSO 5% | 19 | x | x | 3.75-13 fl. oz. | x | 3.75-13 fl. oz. | x | 4h | 78 oz. |
| polyoxin D | | x | x | F | x | F | x | 0d | 6 |
| Pristine | 11+7 | 14.5-18.5 oz. | x | 14.5-18.5 oz. | 14.5-18.5 oz. | 14.5-18.5 fl. oz. | 14.5-18.5 oz. | 12h | 74 oz. |
| pyraclostrobin + boscalid | | F-E | x | E [r] | E | E [r] | E | 0d | 4 |
| Procure 480 SC | 3 | x | x | 8-16 fl. oz. | 8-16 fl. oz. | 8-16 fl. oz. | x | 12h | 64 fl. oz. |
| triflumizole | | x | x | E [r] | E [r] | G [r] | x | 14d | NA |
| Rally 40WSP | 3 | x | x | 5-10 oz. | 5-8 oz. | 5-8 oz. | x | 24h | 5 lb. |
| myclobutanil | | x | x | G [r] | F | G-E [r] | x | 14d | NA |
| Roper DF Rainshield | M | 3 lb. | x | 6 lb. | 6 lb. | 6 lb. | 3 lb. | 24h | 21 lb. |
| mancozeb | | see note above | x | i | G | G | see note above | 77d | 6 |
| Scala (SC) | 9 | x | x | x | x | 7-10 fl. oz. | x | 12h | 40 fl. oz. |
| pyrimethanil | | x | x | x | x | G-E | x | 72d | NA |
| Sercadis | 7 | x | x | 3.5-4.5 fl. oz. | 4.5 fl. oz. | 3.5-4.5 fl. oz. | 4.5 fl. oz. | 12h | 18 fl. oz. |
| fluxapyroxad | | x | x | G | s | G-E | F | 0d | 4 |
| Sovran (50WG) | 11 | x | x | 4-6.4 oz. | 3.2-6.4 fl. oz. | 3.2-6.4 oz. | 4-6.4 oz. | 12h | 25.6 oz. |
| kresoxim-methyl | | x | x | G [r] | E | E [r] | G | 30d | 4 |

(Continued)

Table 1-8. Apple Diseases – Petal Fall¹ (continued)

| Product And Formulation Active Ingredient | FRAC Code ² | Bitter Rot | Fire Blight | Powdery Mildew | Rust | Scab | Summer Rots | REI ³ PHI ⁴ | Max Amt ⁵ Max App ⁶ |
|--|---------------------------|---------------|----------------|-------------------|--------------|------------|----------------|--------------------------------------|--|
| Topguard Specialty Crops (SC) | 3 | 13 fl. oz. | x | 8-12 fl. oz. | 8-12 fl. oz. | 13 fl. oz. | 13 fl. oz. | 12h | 52 fl. oz. |
| flutriafol | | u | x | E | E | G | u | 14d | 4 |
| Topsin-M WSB | 1 | x | x | 0.75-1 lb. | x | 0.75-1 lb. | 0.75-1 lb. | 24h | 4 lb. |
| thiophanate methyl | | x | x | G [r] | x | i | G | 1d | NA |
| Torino (SC) | U6 | x | x | 6.8 oz. | x | x | x | 4h | 6.8 oz. |
| cyflufenamid | | x | x | E | x | x | x | 14d | 1 |
| Vanguard WG (75WG) | 9 | x | x | x | x | 5 oz. | x | 12h | 30 oz. |
| cyprodinil | | x | x | x | x | G | x | 0d | 2 |
| Ziram 76DF | M3 | 6 lb. | x | x | 6 lb. | 6 lb. | 6 lb. | 48h | 42.4 lb. |
| ziram | | G | x | x | G | G | i | 14d | 7 |

Footnotes: See how to read the spray schedule tables section at the beginning of the chapter.

Apple Petal Fall - Insects

Notes on insect pest management

- The pyrethroids Asana, Baythroid, Danitol, Mustang Maxx, Permethrin, and Warrior are labeled for control of plum curculio, oriental fruit moth and leafrollers. However, pyrethroids are not recommended at this stage, because they kill predatory mites that feed on European red mite and two-spotted spider mite, thereby triggering outbreaks of these pest mites.
- Use insecticides only after petal fall is complete to avoid killing bees and other pollinators.
- Rosy apple aphid** is best treated at pink, but there is some chance to control it at petal fall if infestations develop.
- European red mite:** If not managed pre-bloom, it can be managed at petal fall or later; see products listed at green tip (pages 17-19).
- Although Sevin is listed (page 31) for plum curculio control, there is a risk of fruit thinning when used within 30 days of bloom.

Table 1-9. Apple Insects – Petal Fall¹

| Product And Formulation Active Ingredient | IRAC Code ² | Oriental Fruit Moth | Plum Curculio | REI ³ PHI ⁴ | Max Amt ⁵ Max App ⁶ |
|--|---------------------------|------------------------|------------------|--------------------------------------|--|
| Actara (25WDG) | 4A | x | 4.5-5.5 oz. | 12h | 16.5 oz. |
| thiamethoxam | | x | G | 35d | NA |
| Altacor eVo | 28 | 1.3-2.2 fl. oz. | 1.3-2.2 fl. oz. | 4h | 4.6 fl. oz. |
| chlorantraniliprole | | E | s | 5d | 3 |
| Asana XL (0.66EC) (RUP) | 3A | 4.8-14.5 fl. oz. | 4.8-14.5 fl. oz. | 12h | 101.5 fl. oz. |
| esfenvalerate | | E | G | 21d | NA |
| Assail 30SG | 4A | 5-8 oz. | 8 oz. | 12h | 32 oz. |
| acetamiprid | | E | G | 7d | 4 |
| Avaunt eVo | 22A | 5-6 oz. | 5-6 oz. | 12h | 24 oz. |
| indoxacarb | | G | G | 14d | 4 |
| Azera 0.21EC | 3A | 32 fl. oz. | 32 fl. oz. | 12h | NA |
| azadirachtin + pyrethrins | | u | u | 0d | 10 |

(Continued)

Table 1-9. Apple Insects – Petal Fall¹ (continued)

| Product And Formulation Active Ingredient | IRAC Code ² | Oriental Fruit Moth | Plum Curculio | REI ³ PHI ⁴ | Max Amt ⁵ Max App ⁶ |
|--|---------------------------|------------------------|-------------------|--------------------------------------|--|
| Baythroid XL (1EC) (RUP) | 3A | 2-2.4 fl. oz. | 2.4-2.8 fl. oz. | 12h | 2.8 fl. oz. |
| beta-cyfluthrin | | E | G | 7d | NA |
| Belay (2.13SC) | 4A | 6 fl. oz. | 6 fl. oz. | 12h | 12 fl. oz. |
| clothianidin | | G | G | 7d | NA |
| Danitol 2.4EC (RUP) | 3A | 16-21.3 fl. oz. | 16-21.3 fl. oz. | 24h | 42.7 fl. oz. |
| fenpropathrin | | E | G | 14d | NA |
| Delegate WG (25WG) | 5 | 4.5-7 oz. | x | 4h | 28 oz. |
| spinetoram | | E | x | 7d | 4 |
| Delta Gold (1.5EC) (RUP) | 3A | 0.9-1.9 fl. oz. | 0.9-1.9 fl. oz. | 12h | 3.6 fl. oz. |
| deltamethrin | | u | u | 21d | NA |
| Entrust SC (2SC) | 5 | 6-10 fl. oz. | x | 4h | 29 fl. oz. |
| spinosad | | F | x | 7d | 4 |
| Exirel (0.83SE) | 28 | 10-17 fl. oz. | 13.5-20.5 fl. oz. | 12h | 61.5 fl. oz. |
| cyantraniliprole | | E | G | 3d | 3 |
| Imidan 70W | 1B | 2.1-5.75 lb. | 2.1-5.7 lb. | 4 or 7d | 15.5 lb. |
| phosmet | | E | G | 7d | NA |
| Intrepid 2F | 18 | 12-16 fl. oz. | x | 4h | 64 fl. oz. |
| methoxyfenozide | | G | x | 14d | NA |
| Madex HP | 31 | 0.5-3 fl. oz. | x | 4h | NA |
| <i>Cydia pomonella</i> granulovirus | | G | x | 0d | NA |
| Mustang Maxx (0.83EC) (RUP) | 3A | 1.2-4 fl. oz. | 1.2-4 fl. oz. | 12h | 24 fl. oz. |
| zeta-cypermethrin | | E | G | 14d | NA |
| Neemix 4.5 (0.39L) | UN | 4-10 fl. oz. | x | 12h | NA |
| azadirachtin | | F | x | 0d | NA |
| Permethrin 3.2EC (RUP) | 3A | x | 4-10 fl. oz. | 12h | 20 fl. oz. |
| permethrin | | x | G | See label | NA |
| Proclaim (5SG) (RUP) | 6 | 4.8 oz. | x | 12 or 48h | 14.4 oz. |
| emamectin benzoate | | F | x | 14d | NA |
| Rimon 0.83EC | 15 | 20-40 fl. oz. | x | 12h | 150 fl. oz. |
| novaluron | | G | x | 14d | NA |
| Sevin XLR Plus | 1A | x | 1.5-3 qt. | 12h | 15 qt. |
| carbaryl | | x | G | 3d | 8 |
| Verdepryn 100SL (0.83SL) | 28 | 5.5-11 fl. oz. | 5.5-11 fl. oz. | 4h | 33 fl. oz. |
| cyclaniliprole | | E | G | 7d | 3 |
| Warrior II (2.08CS) (RUP) | 3A | 1.2-2.5 fl. oz. | 1.2-2.5 fl. oz. | 24h | 12.8 fl. oz. |
| lambda-cyhalothrin | | E | E | 21d | NA |

Footnotes: See how to read the spray schedule tables section at the beginning of the chapter.

Apple Petal Fall

| Pest/Problem | Material | Rate/Acre | Comments |
|-------------------------------|---|-----------|---|
| For thinning summer varieties | See Chemical Thinning of Apples, page 44. | | |
| nutrient level | Solubor (boron) AND/OR | 2 lb. | May add to pesticide spray solutions, but check for compatibility, order of mixing, etc. Solubor helps prevent cork spot; see page 49 for more information. |
| | feed-grade urea | 8 lb. | |

Apple First to Second Cover

10 days after petal fall and 10 days later.

Cork spot, bitter pit, Jonathan spot

- Calcium chloride is best applied dilute at 8 lb./A dry formulation (1.5-2 lb. per 100 gal.) in the first or second cover. Do not reapply anytime during the growing season if rain has not washed off residue from previous spray.
- Do not exceed 4 pounds per acre for low volume spray. See Cork Spot and Bitter Pit Management in Apples, page 50.

Excess crop: see Chemical Thinning of Apples, page 44.

Apple First to Second Cover - Diseases

10 days after petal fall and 10 days later

Scab, fruit rots (for orchards with a history of fungicide resistance):

- For early apples (e.g., Lodi, Pristine, Yellow Transparent, Zestar) be aware that Luna Tranquility and Mancozeb and have PHIs of 72 and 77 days, respectively. Fontelis, Inspire Super and Omega have 28d PHI; Aprovia, Miravis and Sovran have a 30d PHI.
- The addition of a spreader or penetrating adjuvant such as organo-silicon blends with either non-ionic surfactants (NIS) or vegetable oils (COC; not mineral); or NIS with 90% concentration is recommended with Aprovia. These include but are not limited to Widespread Max or Bond.
- The addition of LI-700 has been found to improve the efficacy of captan in areas where water has a pH greater than 7.0.

- Topsin-M 70WSB may cause scarf skin on Rome apples if applied within a four-week period following petal fall. Excellent for control of some fruit rots but is no longer effective against scab in commercial orchards.

Scab, rust, powdery mildew, fruit rots, sooty blotch, flyspeck (For orchards with minimal issues of fungicide resistance)

- For **powdery mildew** control after second cover, make applications based on field history and orchard scouting.
- Do not apply Merivon with EC or oil-based products. Apply Merivon mixed with captan.
- Wettable sulfur: Do not apply in hot weather (above 80°F). Do not apply within two weeks of an oil spray or spreader-sticker. Can affect fruit finish of Golden Delicious.
- Bitter rot and summer rots:** Rot control begins now for best results. Applying mancozeb at this time using the label's extended program for control of scab provides collateral control of bitter and summer rot. Bitter rot and other summer diseases are effectively controlled by using Merivon plus captan and Aprovia. Be aware of 77-day PHI for early apples.

Fungicide resistance management

- Rotate mode of action. Do not exceed two sequential applications of the same FRAC group. See comments on page 51.

Table 1-10. Apple Diseases – First And Second Cover¹

| Product And Formulation Active Ingredient | FRAC Code ² | Bitter Rot | Powdery Mildew | Rust | Scab | Sooty Blotch/ Fly Speck | Summer Rots (Black And White Rot) | REI ⁴ PHI ³ | Max Amt ⁵ Max App ⁶ |
|--|---------------------------|------------------|--------------------|--------------------|--------------------|----------------------------------|--|--------------------------------------|--|
| Aprovia (EC) | 7 | 5.5-7 fl. oz. | 5.5-7 fl. oz. | 5.5-7 fl. oz. | 5.5-7 fl. oz. | 5.5-7 fl. oz. | 5.5-7 fl. oz. | 12h | 276 fl. oz. |
| benzovindiflupyr | | G-E | F | u | F | G-E | F | 30d | NA |
| Captan 80 WDG | M | 2.5-5 lb. | 2.5-5 lb. | x | 2.5-5 lb. | 2.5-5 lb. | 2.5-5 lb. | 24h | 40 lb. |
| captan | | E | i | x | G | G-E | E | 0d | NA |
| Cevya | 3 | x | 3-5 fl. oz. | 3-5 fl. oz. | 3-5 fl. oz. | 3-5 fl. oz. | 3-5 fl. oz. | 12h | NA |
| mefentrifluconazole | | x | G-E | E | G-E | E | F-G | 0d | NA |
| Excalia (2.84 SC) | 7 | 3-4 fl. oz. | 3-4 fl. oz. | 3-4 fl. oz. | 3-4 fl. oz. | 3-4 fl. oz. | x | 12h | 8 fl. oz. |
| inpyrfluxam | | u | G | u | G-E | u | x | PF | 2 |
| Ferbam Granuflo (76 WDG) | M3 | 3.5 lb. | x | 3.5 lb. | 3.5 lb. | 3.5 lb. | 3.5 lb. | 24h | NA |
| ferbam | | F | x | G | G | F | G | NA | 3 |
| Flint Extra | 11 | 2.9 fl. oz. | 2.5-2.9 fl. oz. | 2.5-2.9 fl. oz. | 2.5-2.9 fl. oz. | 2.5-2.9 fl. oz. | 2.9 fl. oz. | 12h | 10.5 fl. oz. |
| trifloxystrobin | | F-G | G[r] | F | G[r] | G | G | 14d | NA |
| Fontelis (1.67 SC) | 7 | x | 16-20 fl. oz. | 16-20 fl. oz. | 16-20 fl. oz. | x | 16-20 fl. oz. | 12h | 61 fl. oz. |
| penthiopyrad | | x | G | E | G-E | x | u | 28d | NA |
| Indar 2F | 3 | x | 6-8 fl. oz. | 6-8 fl. oz. | 6-8 fl. oz. | 6-8 fl. oz. | x | 12h | 32 fl. oz. |
| fenbuconazole | | x | E[r] | E | E[r] | G | x | 14d | 4 |
| Inspire Super (EW) | 3+9 | x | 12 fl. oz. | 12 fl. oz. | 12 fl. oz. | 12 fl. oz. | x | 12h | 60 fl. oz. |
| difenoconazole + cyprodinil | | x | G | E | G | E | x | 28d | NA |
| Kenja 400SC | 7 | x | 12.5 fl. oz. | x | 12.5 fl. oz. | x | x | 12h | NA |
| isofetamid | | x | s | x | s | x | x | 20d | NA |
| Luna Privilege (SC) | 7 | x | 2.4-6.8 fl. oz. | x | 2.4-6.8 fl. oz. | 6.8 fl. oz. | x | NA | NA |
| fluopyram | | x | G | x | G | F-G | x | NA | NA |
| Luna Sensation (SC) | 7+11 | 4-5.8 fl. oz. | 5-5.8 fl. oz. | 4-5.8 fl. oz. | 5-5.8 fl. oz. | 4-5.8 fl. oz. | 4-5.8 fl. oz. | 12h | 21 fl. oz. |
| fluopyram + trifloxystrobin | | G-E | G[r] | F | G[r] | G-E | E | 14d | 4 |
| Luna Tranquility (SC) | 7+9 | x | 11.2-16 fl. oz. | x | 11.2-16 fl. oz. | x | x | 12h | 54.7 fl. oz. |
| fluopyram + pyrimethanil | | x | G | x | G | x | x | 72d | NA |
| Manzate Pro Stik | M | x | 6 lb. | 6 lb. | 6 lb. | x | 3 lb. | 24h | 21 lb. |
| mancozeb | | x | i | G | G | x | G | 77d | 6 |
| Merivon (2.09SC) | 7+11 | 4-5.5 fl. oz. | 4-5.5 fl. oz. | 4-5.5 fl. oz. | 4-5.5 fl. oz. | 4-5.5 fl. oz. | 4-5.5 fl. oz. | 12h | 22 fl. oz. |
| fluxapyroxad + pyraclostrobin | | E | G | s | G | E | E | 0d | 4 |
| Microthiol Disperss | M | x | 10-20 lb. | x | 10-20 lb. | x | x | 24h | NA |
| sulfur | | x | G | x | F-i | x | x | 0d | NA |

(Continued)

Table 1-10. Apple Diseases – First And Second Cover¹ (continued)

| Product And Formulation Active Ingredient | FRAC Code ² | Bitter Rot | Powdery Mildew | Rust | Scab | Sooty Blotch/ Fly Speck | Summer Rots (Black And White Rot) | REI ⁴ PHI ³ | Max Amt ⁵ Max App ⁶ |
|--|---------------------------|----------------|-------------------|-----------------|-----------------|----------------------------------|--|--------------------------------------|--|
| Miravis (1.67 SC) | 7 | 3.4 fl. oz. | 3.4 fl. oz. | 3.4 fl. oz. | 3.4 fl. oz. | 3.4 fl. oz. | 3.4 fl. oz. | 4h | 13.6 fl. oz. |
| pydiflumetofen | | s-F | G | G | G | G | s | 30d | 4 |
| Omega 500F | 29 | 13.8 fl. oz. | x | 13.8 fl. oz. | 13.8 fl. oz. | 10-13.8 fl. oz. | 13.8 fl. oz. | 12h | 138 fl. oz. |
| fluazinam | | F | x | s[G] | s[G] | F | s-G | 28d | 10 |
| OSO 5% | 19 | x | 3.75-13 fl. oz. | x | 3.75-13 fl. oz. | x | x | 4h | 4.2 oz. |
| polyoxin D | | x | F | x | F | x | x | 0d | 6 |
| Pristine | 11+7 | 14.5-18.5 oz. | 14.5-18.5 oz. | 14.5-18.5 oz. | 14.5-18.5 oz. | 14.5-18.5 oz. | 14.5-18.5 oz. | 12h | 74 oz |
| pyraclostrobin + boscalid | | F-G | E[r] | E | E[r] | E | E | 0d | 4 |
| Procure 480 SC | 3 | x | 8-16 fl. oz. | 8-16 fl. oz. | 8-16 fl. oz. | x | x | 12h | 64 fl. oz. |
| triflumizole | | x | E[r] | E[r] | E[r] | x | x | 14d | NA |
| Rally 40WSP | 3 | x | 5-10 oz. | 5-8 oz. | 5-10 oz. | x | x | 24h | 5 lb. |
| myclobutanil | | x | E[r] | F | E[r] | x | x | 14d | NA |
| Roper DF rainshield | M | 3 lb. | 6 lb. | 6 lb. | 6 lb. | x | 3 lb. | 24h | 21 lb. |
| mancozeb | | see note above | i | G | G | x | see note above | 77d | 6 |
| Sercadis | 7 | x | 3.5-4.5 fl. oz. | 4.5 fl. oz. | 3.5-4.5 fl. oz. | 4.5 fl. oz. | 4.5 fl. oz. | 12h | 18 fl. oz. |
| fluxapyroxad | | x | G | s | F | F | F | 0d | 4 |
| Sovran (50WG) | 11 | x | 4-6.4 oz. | 3.2-6.4 fl. oz. | 4-6.4 oz. | 4-6.4 oz. | 4-6.4 oz. | 12h | 25.6 oz. |
| kresoxim-methyl | | x | G[r] | E | G[r] | G | G | 30d | 4 |
| Topguard Specialty Crops (SC) | 3 | 13 fl. oz. | 8-12 fl. oz. | 8-12 fl. oz. | 8-12 fl. oz. | x | 13 fl. oz. | 12h | 52 fl. oz. |
| flutriafol | | u | E | E | E[r] | x | u | 14d | 4 |
| Topsin-M WSB | 1 | x | 0.75-1 lb. | x | 0.75-1 lb. | 0.75-1 lb. | 0.75-1 lb. | 24h | 4 lb. |
| thiophanate methyl | | x | G[r] | x | G[r] | G-E | G | 1d | NA |
| Torino (SC) | U6 | x | 6.8 oz. | x | x | x | x | 4h | 6.8 oz. |
| cyflufenamid | | x | E | x | x | x | x | 14d | 1 |
| Ziram 76DF | M3 | 6 lb. | x | 6 lb. | 6 lb. | 6 lb. | 6 lb. | 48h | 42.4 lb. |
| ziram | | G | x | G | G | G | i | 14d | 7 |

Footnotes: See how to read the spray schedule tables section at the beginning of the chapter.

Apple First to Second Cover – Insects

7-10 days after petal fall and 7-10 days later.

- Although Sevin is listed for control of several pests, be aware of a risk of fruit thinning when Sevin is used within 30 days of bloom.

Codling moth

- Initiate codling moth control at first or second cover based on timing of capture in pheromone traps. Insecticide timing ranges from 50-250 degree days after trap-based biofix as shown in table below.
- See Mating Disruption for Codling Moth Control, pages 26-27.
- Virus products (Cyd-X, Madex) are for codling moth. Apply virus products at weekly intervals.

Timing Of First Insecticide Spray For Codling Moth Control On Apple And Pear¹

| Degree-Days (Base 50°F) After Biofix ² | Insecticide Products |
|---|--|
| 50-75 | Dimilin, Rimon |
| 100-200 | Intrepid, Confirm |
| 150-250 | Altacor, Assail, Belay, Delegate, Exirel |
| 250 | Imidan, Avaunt pyrethroids (Asana, Baythroid, Danitol, Mustang Maxx, Warrior) Virus (Cyd-X, Madex) |

¹A second spray should be made 10-14 days later.

²Biofix is defined as the date on which pheromone traps detect sustained flight of moths.

San Jose scale "crawlers"

- May be present by second or third cover. Esteem 35W controls scale anytime between half-inch green and third cover. The minimum rate of Esteem is effective when used pre-bloom, but use the maximum rate if application is delayed until the crawler stage in early summer. One application is sufficient for Esteem, but for other products, use two applications, 10 days apart

Potato leafhopper

- Effectiveness of products for leafhopper control are shown in the table for minor pests on pages 55-58.

Table 1-11. Apple Insects – First And Second Cover¹

| Product And Formulation Active Ingredient | IRAC Code ² | Codling Moth | Oriental Fruit Moth | Plum Curculio | San Jose Scale | REI ³ PHI ⁴ | Max Amt ⁵ Max App ⁶ |
|--|------------------------|------------------|---------------------|------------------|------------------|-----------------------------------|--|
| Actara (25WDG) | 4A | x | x | 4.5-5.5 oz. | x | 12h | 16.5 oz. |
| thiamethoxam | | x | x | G | x | 35d | NA |
| Altacor eVo | 28 | 1.3-2.2 fl. oz. | 1.3-2.2 fl. oz. | 1.3-2.2 fl. oz. | x | 4h | 4.6 fl. oz. |
| chlorantraniliprole | | E | E | s | x | 5d | 3 |
| Asana XL (0.66EC) (RUP) | 3A | 4.8-14.5 fl. oz. | 4.8-14.5 fl. oz. | 4.8-14.5 fl. oz. | 4.8-14.5 fl. oz. | 12h | 101.5 fl. oz. |
| esfenvalerate | | E | E | G | i | 21d | NA |
| Assail 30SG | 4A | 4-8 oz. | 5-8 oz. | 8 oz. | 8 oz. | 12h | 32 oz. |
| acetamiprid | | E | E | G | s | 7d | 4 |
| Avaunt eVo | 22A | 5-6 oz. | 5-6 oz. | 5-6 oz. | x | 12h | 24 oz. |
| indoxacarb | | F | G | G | x | 14d | 4 |

(Continued)

Table 1-11. Apple Insects – First And Second Cover¹ (continued)

| Product And Formulation Active Ingredient | IRAC Code ² | Codling Moth | Oriental Fruit Moth | Plum Curculio | San Jose Scale | REI ³ PHI ⁴ | Max Amt ⁵ Max App ⁶ |
|--|---------------------------|-----------------|------------------------|-------------------|-------------------|--------------------------------------|--|
| Azera 0.21EC | 3A | 32 fl. oz. | 32 fl. oz. | 32 fl. oz. | 32 fl. oz. | 12h | NA |
| azadirachtin + pyrethrins | | x | x | x | x | 0d | 10 |
| Dipel | 11A | 0.5-2 lb. | x | x | x | NA | NA |
| B.t. kurstaki | | F | x | x | x | NA | NA |
| Baythroid XL (1EC) (RUP) | 3A | 2-2.4 fl. oz. | 2-2.4 fl. oz. | 2.4-2.8 fl. oz. | 2.4-2.8 fl. oz. | 12h | 2.8 fl. oz. |
| beta-cyfluthrin | | E | E | G | G | 7d | NA |
| Belay (2.13SC) | 4A | 6 fl. oz. | 6 fl. oz. | 6 fl. oz. | 6 fl. oz. | 12h | 12 fl. oz. |
| clothianidin | | G | G | G | G | 7d | NA |
| Centaur WDG (70WDG) | 16 | x | x | x | 34.5 oz. | 12h | 34.5 oz. |
| buprofezin | | x | x | x | E | 14d | 1 |
| Confirm 2F | 18 | 20 fl. oz. | x | x | x | 4h | 120 fl. oz. |
| tebufenozide | | F | x | x | x | 14d | NA |
| Cyd-X HP | 0 | 0.5-3 fl. oz. | x | x | x | 4h | NA |
| <i>Cydia pomonella</i> granulovirus | | F | x | x | x | 0d | NA |
| Danitol 2.4EC (RUP) | 3A | 16-21.3 fl. oz. | 16-21.3 fl. oz. | 16-21.3 fl. oz. | x | 24h | 42.7 fl. oz. |
| fenpropathrin | | E | E | G | x | 14d | NA |
| Delegate WG (25WG) | 5 | 4.5-7 oz. | 4.5-7 oz. | s | x | 4h | 28 oz. |
| spinetoram | | E | E | s | x | 7d | 4 |
| Delta Gold (1.5EC) (RUP) | 3A | 0.9-1.9 fl. oz. | 0.9-1.9 fl. oz. | 0.9-1.9 fl. oz. | 0.9-1.9 fl. oz. | 12h | 3.6 fl. oz. |
| deltamethrin | | u | u | u | u | 21d | NA |
| Dipel | 11A | 0.5-2 lb. | x | x | x | NA | NA |
| B.t. kurstaki | | F | x | x | x | NA | NA |
| Entrust SC (2SC) | 5 | 6-10 fl. oz. | 6-10 fl. oz. | x | x | 4h | 29 fl. oz. |
| spinosad | | G | F | x | x | 7d | 4 |
| Esteem 35WP | 7C | 4-5 oz. | x | x | 4-5 oz. | 12h | 10 oz. |
| pyriproxifen | | F | x | x | E | 45d | 2 |
| Exirel (0.83SE) | 28 | 8.5-17 fl. oz. | 10-17 fl. oz. | 13.5-20.5 fl. oz. | x | 12h | 61.5 fl. oz. |
| cyantraniliprole | | E | E | G | x | 3d | 3 |
| Imidan 70W | 1B | 2.1-5.75 lb. | 2.1-5.75 lb. | 2.1-5.7 lb. | 2.1-5.75 lb. | 4 or 7d | 15.5 lb. |
| phosmet | | G | E | G | F | 7d | NA |
| Intrepid 2F | 18 | 16 fl. oz. | 12-16 fl. oz. | x | x | 4h | 64 fl. oz. |
| methoxyfenozide | | s | G | x | x | 14d | NA |
| Lannate LV (2.4WSL) (RUP) | 1A | 3 pt. | x | x | x | 3d | 15 pt. |
| methomyl | | G | x | x | x | 14d | 5 |
| Madex HP | 3I | 0.5-3 fl. oz. | 0.5-3 fl. oz. | x | x | 4h | NA |
| <i>Cydia pomonella</i> granulovirus | | G | G | x | x | 0d | NA |
| Movento (2SC) | 23 | 6-9 fl. oz. | x | x | 6-9 fl. oz. | 24h | 25 fl. oz. |
| spirotetramat | | s | x | x | G | 7d | NA |
| Mustang Maxx (0.83EC) (RUP) | 3A | 1.2-4 fl. oz. | 1.2-4 fl. oz. | 1.2-4 fl. oz. | x | 12h | 24 fl. oz. |
| zeta-cypermethrin | | E | E | G | x | 14d | NA |

(Continued)

Table 1-11. Insecticides for control of apple insects – first and second cover¹ (continued)

| Product And Formulation Active Ingredient | IRAC Code ² | Codling Moth | Oriental Fruit Moth | Plum Curculio | San Jose Scale | REI ³ PHI ⁴ | Max Amt ⁵ Max App ⁶ |
|--|---------------------------|-----------------|------------------------|------------------|-------------------|--------------------------------------|--|
| Neemix 4.5 (0.39L) | UN | 4-10 fl. oz. | 4-10 fl. oz. | x | 7-16 fl. oz. | 12h | NA |
| azadirachtin | | F | F | i | u | 0d | NA |
| Permethrin 3.2EC (RUP) | 3A | x | x | 4-10 fl. oz. | x | 12h | 20 fl. oz. |
| permethrin | | x | x | G | x | See Label | NA |
| Proclaim (5SG) (RUP) | 6 | 4.8 oz. | 4.8 oz. | x | x | 12 or 48h | 14.4 oz. |
| emamectin benzoate | | F | F | x | x | 14d | NA |
| Rimon 0.83EC | 15 | 20-40 fl. oz. | 20-40 fl. oz. | x | x | 12h | 150 fl. oz. |
| novaluron | | E | G | x | x | 14d | NA |
| Sevin XLR Plus | 1A | 1-3 qt. | x | 1.5-3 qt. | 1.5-3 qt. | 12h | 15 qt. |
| carbaryl | | G | x | G | F | 3d | 8 |
| Sivanto Prime (1.67SC) | 4D | x | x | x | 10.5-14 fl. oz. | 4h | 28 fl. oz. |
| flupyradifurone | | x | x | x | G | 14d | NA |
| Verdepryn 100SL (0.83SL) | 28 | 5.5-11 fl. oz. | 5.5-11 fl. oz. | 5.5-11 fl. oz. | x | 4h | 33 fl. oz. |
| cyclanilprole | | E | E | G | x | 7d | 3 |
| Warrior II (2.08CS) (RUP) | 3A | 1.2-2.5 fl. oz. | 1.2-2.5 fl. oz. | 1.2-2.5 fl. oz. | 1.2-2.5 fl. oz. | 24h | 12.8 fl. oz. |
| lambda-cyhalothrin | | E | E | E | i | 21d | NA |

Footnotes: See how to read the spray schedule tables section at the beginning of the chapter.

Apple Third to Summer Covers – Diseases

Third cover: 10-14 days after second cover.

Apple summer cover sprays: Depending on rainfall, apply at intervals of 10 to 14 days.

- Roper (mancozeb) has a 77-day PHI and should not be used on early apples or after third cover.
- Fontelis, Inspire Super and Omega have 28d PHI.
- Aprovia, Miravis and Sovran have a 30d PHI.
- ProPhyt plus captan has provided control of sooty blotch and flyspeck equal to captan plus Topsin-M. ProPhyt plus captan tank-mix is also effective against bitter rot and other summer diseases if applied alternatively with Merivon plus captan.
- Captan 80WDG 4 lb. plus the use of an acidifier may be equally effective. Other formulations are available, such as 4L and 50WP. See Use of Captan Fungicide on Tree Fruit-Restricted Entry Intervals (REI), page 50.

Table 1-12. Apple Diseases – Third And Summer Cover¹

| Product And Formulation Active Ingredient | FRAC Code ² | Bitter Rot | Powdery Mildew | Scab | Sooty Blotch/Fly Speck | Summer Rots (Black And White Rot) | REI ³ PHI ⁴ | Max Amt ⁵ Max App ⁶ |
|--|---------------------------|---------------|-------------------|---------------|------------------------------|---|--------------------------------------|--|
| Aprovia (EC) | 7 | 5.5-7 fl. oz. | 5.5-7 fl. oz. | 5.5-7 fl. oz. | 5.5-7 fl. oz. | 5.5-7 fl. oz. | 12h | 27.6 fl. oz. |
| benzovindiflupyr | | G-E | F | G-E | G-E | G | 30d | NA |
| Captan 80 WDG | M | 2.5-5 lb. | 2.5-5 lb. | 5 lb. | 2.5-5 lb. | 2.5-5 lb. | 24h | 40 lb. |
| captan | | E | i | G | G-E | E | 0d | NA |
| Cevya | 3 | x | 3-5 fl. oz. | 3-5 fl. oz. | 3-5 fl. oz. | 3-5 fl. oz. | 12h | NA |
| mefentrifluconazole | | x | G-E | E | E | F-G | 0d | NA |

(Continued)

Table 1-12. Apple Diseases - Third And Summer Cover¹ (continued)

| Product And Formulation Active Ingredient | FRAC Code ² | Bitter Rot | Powdery Mildew | Scab | Sooty Blotch/Fly Speck | Summer Rots (Black And White Rot) | REI ³ PHI ⁴ | Max Amt ⁵ Max App ⁶ |
|--|---------------------------|---------------|--------------------|--------------------|------------------------------|---|--------------------------------------|--|
| Cuprofix Ultra 40D | M | x | x | 1-2.5 lb. | x | x | 12hr | NA |
| copper sulfate | | x | x | F | x | x | NA | NA |
| Excalia (2.84 SC) | 7 | 3-4 fl. oz. | 3-4 fl. oz. | 3-4 fl. oz. | 3-4 fl. oz. | x | 12h | 8 fl. oz. |
| inpyrfluxam | | u | G | E | u | x | PF | 2 |
| Ferbam Granuflo (76 WDG) | M3 | 3.5 lb. | x | 3.5 lb. | 3.5 lb. | 3.5 lb. | 24h | NA |
| ferbam | | F | x | F | F | G | NA | 3 |
| Flint Extra | 11 | 2.9 fl. oz. | 2.5-2.9 fl. oz. | 2.5-2.9 fl. oz. | 2.5-2.9 fl. oz. | 2.9 fl. oz. | 12h | 10.5 fl. oz. |
| trifloxystrobin | | F-G | G [r] | E [r] | G | G | 14d | NA |
| Fontelis (1.67 SC) | 7 | x | 16-20 fl. oz. | 16-20 fl. oz. | x | 16-20 fl. oz. | 12h | 61 fl. oz. |
| penthiopyrad | | x | G | E | x | u | 28d | NA |
| Indar 2F | 3 | x | 6-8 fl. oz. | 6-8 fl. oz. | 6-8 fl. oz. | x | 12h | 32 fl. oz. |
| fenbuconazole | | x | E [r] | E [r] | G | x | 14d | 4 |
| Inspire Super (EW) | 3+9 | x | 12 fl. oz. | 12 fl. oz. | 12 fl. oz. | x | 12h | 60 fl. oz. |
| difenoconazole + cyprodinil | | x | F | E | E | x | 28d | NA |
| Kenja 400SC | 7 | x | 12.5 fl. oz. | 12.5 fl. oz. | x | x | 12h | NA |
| isofetamid | | x | s | F | x | x | 20d | NA |
| Luna Privilege (SC) | 7 | x | 2.4-6.8 fl. oz. | 4-6.8 fl. oz. | 6.8 fl. oz. | x | NA | NA |
| fluopyram | | x | G | G-E | F-G | x | NA | NA |
| Luna Sensation (SC) | 7+11 | 4-5.8 fl. oz. | 5-5.8 fl. oz. | 4-5.8 fl. oz. | 4-5.8 fl. oz. | 4-5.8 fl. oz. | 12h | 21 fl. oz. |
| fluopyram + trifloxystrobin | | G | G [r] | E [r] | G-E | E | 14d | 4 |
| Luna Tranquility (SC) | 7+9 | x | 11.2-16 fl. oz. | 11.2-16 fl. oz. | x | x | 12h | 54.7 fl. oz. |
| fluopyram + pyrimethanil | | x | G | E | x | x | 72d | NA |
| Merivon (2.09SC) | 7+11 | 4-5.5 fl. oz. | 4-5.5 fl. oz. | 4-5.5 fl. oz. | 4-5.5 fl. oz. | 4-5.5 fl. oz. | 12h | 22 fl. oz. |
| fluxapyroxad + pyraclostrobin | | E | G | E | E | E | 0d | 4 |
| Microthiol Disperss | M | x | 10-20 lb. | 10-20 lb. | x | x | 24h | NA |
| sulfur | | x | G | i-F | x | x | 0d | NA |
| Miravis (1.67 SC) | 7 | 3.4 fl. oz. | 3.4 fl. oz. | 3.4 fl. oz. | 3.4 fl. oz. | 3.4 fl. oz. | 4h | 13.6 fl. oz. |
| pydiflumetofen | | s-F | G | E | G | s | 30d | 4 |
| Omega 500F | 29 | 13.8 fl. oz. | x | 10-13.8 fl. oz. | 10-13.8 fl. oz. | 13.8 fl. oz. | 12h | 138 fl. oz. |
| fluazinam | | F | x | G | F | s-G | 28d | 10 |
| OSO 5% | 19 | x | 3.75-13 fl. oz. | 3.75-13 fl. oz. | x | x | 4h | 78 oz. |
| polyoxin D | | x | F | F | x | x | 0d | 6 |

(Continued)

Table 1-12. Apple Diseases - Third And Summer Cover¹ (continued)

| Product And Formulation Active Ingredient | FRAC Code ² | Bitter Rot | Powdery Mildew | Scab | Sooty Blotch/Fly Speck | Summer Rots (Black And White Rot) | REI ³ PHI ⁴ | Max Amt ⁵ Max App ⁶ |
|--|---------------------------|-------------------|--------------------|--------------------|------------------------------|---|--------------------------------------|--|
| Pristine | 11+7 | 14.5-18.5 oz. | 14.5-18.5 oz. | 14.5-18.5 oz. | 14.5-18.5 oz. | 14.5-18.5 oz. | 12h | 74 oz. |
| pyraclostrobin + boscalid | | F-G | E [r] | E [r] | E | G | 0d | 4 |
| Procure 480 SC | 3 | x | 8-16 fl. oz. | 8-16 fl. oz. | x | x | 12h | 64 fl. oz. |
| triflumizole | | x | E [r] | G [r] | x | x | 14d | NA |
| Rally 40WSP | 3 | x | 5-10 oz. | 5-8 oz. | x | x | 24h | 5 lb. |
| myclobutanil | | x | E [r] | G-E [r] | x | x | 14d | NA |
| Roper DF Rainshield | M | 3 lb. | 6 lb. | 6 lb. | x | 3 lb. | 24h | 21 lb. |
| mancozeb | | see note above | i | G | x | see note above | 77d | 6 |
| Sercadis | 7 | x | 3.5-4.5 fl. oz. | 3.5-4.5 fl. oz. | 4.5 fl. oz. | 4.5 fl. oz. | 12h | 18 fl. oz. |
| fluxapyroxad | | x | G | G-E | F | F | 0d | 4 |
| Sovran (50WG) | 11 | x | 4-6.4 oz. | 3.2-6.4 oz. | 4-6.4 oz. | 4-6.4 oz. | 12h | 25.6 oz. |
| kresoxim-methyl | | x | G [r] | E [r] | G | G | 30d | 4 |
| Topguard Specialty Crops (SC) | 3 | 13 fl. oz. | 8-12 fl. oz. | 13 fl. oz. | x | 13 fl. oz. | 12h | 52 fl. oz. |
| flutriafol | | u | E | G | x | u | 14d | 4 |
| Topsin-M WSB | 1 | x | 0.75-1 lb. | 0.75-1 lb. | 0.75-1 lb. | 0.75-1 lb. | 24h | 4 lb. |
| thiophanate methyl | | x | G [r] | i | G-E | G | 1d | NA |
| Vanguard WG (75WG) | 9 | x | x | 5 oz. | x | x | 12h | 30 oz. |
| cyprodinil | | x | x | G | x | x | 0d | 2 |
| Ziram 76DF | M3 | 6 lb. | x | 6 lb. | 6 lb. | 6 lb. | 48h | 42.4 lb. |
| ziram | | G | x | G | G | i | 14d | 7 |

Footnotes: See how to read the spray schedule tables section at the beginning of the chapter.

Apple Third to Summer Covers - Insects

Third cover: 10 days after second cover.

Apple summer cover sprays: Depending on rainfall, apply at intervals of 10 (if rainy) to 14 days (if dry).

Apple maggot flies

- Flies begin emerging from the soil about mid-June. Monitor for the first appearance of flies each year by examining fruit and leaves in the center of trees in detail, using yellow sticky board traps baited with an attractant, hanging red or green spheres coated with a sticky substance in trees, or combining all three methods. Continue applications until late September or as long as flies are present.

Codling moth

- Timing for sprays depends on which insecticides will be used. See table on page 52 for target timing

for each product. Because codling moths have multiple generations per year, additional applications may be necessary later in the season to maintain control.

Japanese beetles

- Begin treatments as soon as observed. Multiple applications may be necessary.

Brown marmorated stink bug (BMSB)

- The products below may be needed despite their harm to beneficial arthropods that help control San Jose scale, woolly apple aphid, and mites.
- Baythroid XL 1EC, Belay 2.13SC, Danitol 2.4EC, Mustang Maxx, Lannate SP, Lannate LV, and Warrior II are only labeled for general stink bug control, not specifically for brown marmorated stink bug.

Plum curculio

- A second generation of plum curculio has been observed in some areas of the Midwest, which may require additional management to maintain control.

San Jose scale crawlers

- Crawlers may be present by second or third cover. The minimum rate of Esteem 35W is effective when used pre-bloom, but use the maximum rate if application is delayed until the crawler stage in early summer. One application is sufficient for Esteem, but for other products, use two applications, 10 days apart.

Table 1-13. Apple Insects - Third And Summer Cover¹

| Product And Formulation Active Ingredient | IRAC Code ² | Apple Maggot | Brown Marmorated Stink Bug | Codling Moth | Japanese Beetle | Oriental Fruit Moth | San Jose Scale | Woolly Apple Aphid | REI ³ PHI ⁴ | Max Amt ⁵ Max App ⁶ |
|--|------------------------|------------------|----------------------------|------------------|-----------------|---------------------|------------------|--------------------|-----------------------------------|--|
| Admire Pro (4.6F) | 4A | 2.8 fl. oz. | x | x | x | x | 2.8-8 fl. oz. | 7-10.5 fl. oz. | 12h | 10.5 fl. oz. |
| imidacloprid | | G | x | x | x | x | F | G | 21d | NA |
| Altacor eVo | 28 | 1.3-2.2 fl. oz. | x | 1.3-2.2 fl. oz. | x | 1.3-2.2 fl. oz. | x | x | 4h | 4.6 fl. oz. |
| chlorantraniliprole | | s | x | E | x | E | x | x | 5d | 3 |
| Asana XL (0.66EC) (RUP) | 3A | 4.8-14.5 fl. oz. | 4.8-14.5 fl. oz. | 4.8-14.5 fl. oz. | x | 4.8-14.5 fl. oz. | 4.8-14.5 fl. oz. | x | 12h | 101.5 fl. oz. |
| esfenvalerate | | G | E | E | x | E | i | x | 21d | NA |
| Assail 30SG | 4A | 8 oz. | x | 4-8 oz. | 5-8 oz. | 5-8 oz. | 8 oz. | 2.5-4 oz. | 12h | 32 oz. |
| acetamiprid | | G | x | E | G | E | s | F | 7d | 4 |
| Avaunt eVo | 22 | 6 fl. oz. | x | 5-6 oz. | x | 5-6 oz. | x | x | 12h | 24 oz. |
| indoxacarb | | F | x | F | x | G | x | x | 14d | 4 |
| Azera 0.21EC | 3A | 32 fl. oz. | 32 fl. oz. | 32 fl. oz. | 32-56 fl. oz. | 32 fl. oz. | 32 fl. oz. | 32 fl. oz. | 12h | NA |
| azadirachtin + pyrethrins | | u | u | u | G | x | u | u | 0d | 10 |
| Baythroid XL (1EC) (RUP) | 3A | 2.4-2.8 fl. oz. | 2-2.4 fl. oz. | 2-2.4 fl. oz. | x | 2-2.4 fl. oz. | 2.4-2.8 fl. oz. | x | 12h | 2.8 fl. oz. |
| beta-cyfluthrin | | G | E | E | x | E | G | x | 7d | NA |
| Belay (2.13SC) | 4A | 6 fl. oz. | 4-6 fl. oz. | 6 fl. oz. | x | 6 fl. oz. | 6 fl. oz. | 4-6 fl. oz. | 12h | 12 fl. oz. |
| clothianidin | | G | G | G | x | G | G | u | 7d | NA |
| Beleaf 50SG | 29 | x | x | x | x | x | x | 2-2.8 oz. | 12h | 8.4 oz. |
| flonicamid | | x | x | x | x | x | x | F | 21d | 3 |
| Centaur WDG (70WDG) | 16 | x | x | x | x | x | 34.5 oz. | x | 12h | 34.5 oz. |
| buprofezin | | x | x | x | x | x | E | x | 14d | 1 |
| Confirm 2F | 18 | x | x | 20 fl. oz. | x | x | x | x | 4h | 120 fl. oz. |
| tebufenozide | | x | x | F | x | x | x | x | 14d | NA |
| Cyd-X HP | 0 | x | x | 0.5-3 fl. oz. | x | x | x | x | 4h | NA |
| <i>Cydia pomonella</i> granulovirus | | x | x | F | x | x | x | x | 0d | NA |

(Continued)

Table 1-13. Apple Insects - Third And Summer Cover¹ (continued)

| Product And Formulation Active Ingredient | IRAC Code ² | Apple Maggot | Brown Marmorated Stink Bug | Codling Moth | Japanese Beetle | Oriental Fruit Moth | San Jose Scale | Woolly Apple Aphid | REI ³ PHI ⁴ | Max Amt ⁵ Max App ⁶ |
|--|------------------------|-------------------|----------------------------|-----------------|-----------------|---------------------|------------------------|------------------------|--------------------------------------|--|
| Danitol 2.4EC (RUP) | 3A | 16-21.3 fl. oz. | x | 16-21.3 fl. oz. | 16-21.3 fl. oz. | 16-21.3 fl. oz. | x | x | 24h | 42.7 fl. oz. |
| fenpropathrin | | G | x | E | E | E | x | x | 14d | NA |
| Delegate WG (25WG) | 5 | 6-7 oz. | x | 4.5-7 oz. | x | 4.5-7 oz. | x | x | 4h | 28 oz. |
| spinetoram | | s | x | E | x | E | x | x | 7d | 4 |
| Delta Gold (1.5EC) (RUP) | 3A | 0.9-1.9 fl. oz. | 1.9 fl. oz. | 0.9-1.9 fl. oz. | x | 0.9-1.9 fl. oz. | 0.9-1.9 fl. oz. | x | 12h | 3.6 fl. oz. |
| deltamethrin | | u | u | u | x | u | u | x | 21d | NA |
| Diazinon AG 600 (RUP) | 1B | x | x | x | x | x | 12.75 fl. oz./100 gal. | 12.75 fl. oz./100 gal. | 4h | 25.5 fl. oz. |
| diazinon | | x | x | x | x | x | u | u | 21d | 2 |
| Dipel | 11A | x | x | 0.5-2 lb. | x | x | x | x | NA | NA |
| B.t. kurstaki | | x | x | F | x | x | x | x | NA | NA |
| Entrust SC (2SC) | 5 | 6-10 fl. oz. | x | 6-10 fl. oz. | x | 6-10 fl. oz. | x | x | 4h | 29 fl. oz. |
| spinosad | | s | x | G | x | F | x | x | 7d | 4 |
| Esteem 35WP | 7C | x | x | 4-5 oz. | x | x | 4-5 oz. | x | 12h | 10 oz. |
| pyriproxyfen | | x | x | F | x | x | E | x | 45d | 2 |
| Exirel (0.83SE) | 28 | 13.5-20.5 fl. oz. | x | 8.5-17 fl. oz. | x | 10-17 fl. oz. | x | x | 12h | 61.5 fl. oz. |
| cyantraniliprole | | s | x | E | x | E | x | x | 3d | 3 |
| Imidan 70W | 1B | 2.1-5.75 lb. | x | 2.1-5.75 lb. | 2.1-5.75 lb. | 2.1-5.75 lb. | 2.125-5.75 lb. | x | 4 or 7d | 15.5 lb. |
| phosmet | | E | x | G | G | E | F | x | 7d | NA |
| Intrepid 2F | 18 | x | x | 16 fl. oz. | x | 12-16 fl. oz. | x | x | 4h | 64 fl. oz. |
| methoxyfenozide | | x | x | s | x | G | x | x | 14d | NA |
| Lannate LV (2.4WSL) (RUP) | 1A | x | 1.5-3 pt. | 3 pt. | x | x | x | x | 3d | 15 pt. |
| methomyl | | x | i | G | x | x | x | x | 14d | 5 |
| Madex HP | 31 | x | x | 0.5-3 fl. oz. | x | 0.5-3 fl. oz. | x | x | 4h | NA |
| <i>Cydia pomonella</i> granulovirus | | x | x | G | x | G | x | x | 0d | NA |
| Movento (2SC) | 23 | x | x | 6-9 fl. oz. | x | x | 6-9 fl. oz. | 6-9 fl. oz. | 24h | 25 fl. oz. |
| spirotetramat | | x | x | s | x | x | G | G | 7d | NA |
| Mustang Maxx (0.83EC) (RUP) | 3A | 1.2-4 fl. oz. | 1.2-4 fl. oz. | 1.2-4 fl. oz. | 1.2-4 fl. oz. | 1.2-4 fl. oz. | x | x | 12h | 24 fl. oz. |
| zeta-cypermethrin | | G | E | E | E | E | x | x | 14d | NA |
| Neemix 4.5 (0.39L) | UN | x | 7-16 fl. oz. | 4-10 fl. oz. | 4-16 fl. oz. | 4-10 fl. oz. | 7-16 fl. oz. | 5-7 fl. oz. | 12h | NA |
| azadirachtin | | x | u | F | F | F | u | u | 0d | NA |

(Continued)

Table 1-13. Apple Insects - Third And Summer Cover¹ (continued)

| Product And Formulation Active Ingredient | IRAC Code ² | Apple Maggot | Brown Marmorated Stink Bug | Codling Moth | Japanese Beetle | Oriental Fruit Moth | San Jose Scale | Woolly Apple Aphid | REI ³ PHI ⁴ | Max Amt ⁵ Max App ⁶ |
|--|------------------------|-----------------|----------------------------|-----------------|-----------------|---------------------|-----------------|--------------------|-----------------------------------|---|
| Proclaim (5SG) (RUP) | 6 | x | x | 4.8 oz. | x | 4.8 oz. | x | x | 12 or 48h | 14.4 oz. |
| emamectin benzoate | | x | x | F | x | F | x | x | 14d | NA |
| Rimon 0.83EC | 15 | x | 20-30 fl. oz. | 20-40 fl. oz. | x | 20-40 fl. oz. | x | x | 12h | 150 fl. oz. |
| novaluron | | x | i | E | x | G | x | x | 14d | NA |
| Sevin XLR Plus | 1A | 1.5-3 qt. | x | 1-3 qt. | 1.5-3 qt. | x | 1.5-3 qt. | 1.5-3 qt. | 12h | 15 qt. |
| carbaryl | | G | x | G | E | x | F | i | 3d | 8 |
| Sivanto Prime (1.67SC) | 4D | x | x | x | x | x | 10.5-14 fl. oz. | 12-14 fl. oz. | 4h | 28 fl. oz. |
| flupyradifurone | | x | x | x | x | x | G | s | 14d | NA |
| Surround WP (95WP) | UN | x | x | x | 25-50 lb. | x | x | x | 4h | NA |
| kaolin | | x | x | x | u | x | x | x | 0d | NA |
| Verdepryn 100SL (0.83SL) | 28 | x | 5.5-11 fl. oz. | 5.5-11 fl. oz. | x | 5.5-11 fl. oz. | x | x | 4h | 33 fl. oz. |
| cyclaniliprole | | x | s | E | x | E | x | x | 7d | 3 |
| Versys Inscalis (0.83DC) | 9D | x | x | x | x | x | x | 3.5 fl. oz. | 12h | 7 fl. oz. |
| afidopyropen | | x | x | x | x | x | x | s | 7d | NA |
| Vydate L (2L) (RUP) | 1A | x | 1.5-4 pt. | x | x | x | x | x | 48h | 8 pt. |
| oxamyl | | x | i | x | x | x | x | x | 14d | 4 |
| Warrior II (2.08CS) (RUP) | 3A | 1.2-2.5 fl. oz. | 1.2-2.5 fl. oz. | 1.2-2.5 fl. oz. | 1.2-2.5 fl. oz. | 1.2-2.5 fl. oz. | 1.2-2.5 fl. oz. | x | 24h | 12.8 fl. oz. |
| lambda-cyhalothrin | | F | E | E | E | E | i | x | 21d | NA |

Footnotes: See how to read the spray schedule tables section at the beginning of the chapter.

Special Problems and Pests of Apple

Ambrosia Beetles

- Ambrosia beetles are an emerging pest of apple, particularly in young or high-density plantings. Outbreaks are often associated with tree stress caused by the orchard site, including poor drainage or winter injury. Activity begins in early spring as females target stressed trees for colonization. Damage appears as small, round holes in the trunk, often surrounded by fine sawdust. Ambrosia beetles can be monitored using ethanol-baited traps placed at or below canopy height. Preventive trunk-applied pyrethroid sprays may provide limited protection if applied before attack but management should focus on improving drainage and reducing stress to maintain tree health.

For more detailed information about disease and insect control and integrated pest management (IPM), consult the *Midwest Tree Fruit Pest Management Handbook* and use it in conjunction with this guide. Contact your state's Cooperative Extension service to get a copy.

Horticultural Management

Special notes for growth regulator use in apples

Blush on apples

Blush and Blush 2x contain 5.25% and 10% Prohydrojasmon propyl-3-oxo-2-pentylcyclopentylacetate,

respectively. Blush is used to promote early color change in red apples that have difficulty developing color. Effectiveness depends on cultivar and environmental conditions. Cultivars or strains in environments where color development is normally satisfactory may not develop significant additional red color.

Make 1-2 applications of Blush at 26-52 fl. oz. or Blush 2X at 13-52 fl. oz. per acre per year at 7-14 day intervals, 7-42 days prior to anticipated harvest. Use the higher rate if environmental conditions are not as favorable for color development. Avoid applications during the hottest part of the day. Apply under slow-drying conditions in early morning or at night. Do not apply to stressed or injured trees.

Ethephon on apples

Ethephon, which is available as a 21.3% formulation of 2-chloroethylphosphonic acid (Ethrel or Motivate) may:

1. Promote early color development and maturity.
2. Loosen fruit for easier harvesting by hand or machine.
3. Increase fruit bud formation and early bearing on young trees.
4. Promote fruit thinning and return bloom.
5. Cause premature fruit drop, particularly on spur-type trees.

Promoting early color development and maturity

To increase red coloration and early maturity, apply Ethephon 2 14 to 21 days prior to anticipated harvest and 1-2 weeks before the preferred harvest time at 1-4 pt. per acre for early to mid-season maturing varieties and 2-4 pt. per acre for later maturing (maturing after McIntosh) varieties. Apply in a dilute spray with plenty of water to ensure thorough coverage.

Color development should be apparent in about 7 days. Ethephon 2 is most effective under weather conditions that favor color development.

Do not apply ethephon during hot weather or when hot weather is forecasted in the next 14 days. Apply ethephon between 60°F and 90°F. Most red apple varieties do not develop red color during hot weather with or without ethephon. Ethephon speeds up ripening. Do not use ethephon on Golden Delicious.

Add a fruit drop inhibitor to control pre-harvest drop of the fruit. You can add naphthaleneacetic acid (NAA) to the same spray as ethephon. NAA is effective for 7 to 10 days, and a second application might be necessary if harvest is delayed.

No spreader-sticker is necessary. Ethephon does not overcome poor management practices. Trees of mod-

erate vigor, well-pruned, and thoroughly sprayed, respond most favorably with well-colored fruit of uniform maturity. For dense trees, harvest outer fruit first and then apply ethephon. Harvest at proper maturity; do not delay harvest to obtain additional red coloration. Treat only the acreage that you can harvest and market in a timely manner. You should promptly market fruit treated with ethephon because it may have short shelf life.

Increasing early bearing on young trees

To increase fruit bud development on young, nonbearing trees, apply a foliar spray of Ethephon 2 two to four weeks after full bloom. Spray trees thoroughly almost to runoff at 2-8 pt per acre.

Increasing flower bud development on bearing trees

Delay treatment until after June drop and six weeks following full bloom to help prevent fruit thinning. Apply 0.5-3 pt. per acre. Yield and fruit size reduction may occur; however, flowering should increase the following spring.

Do not use on low vigor trees, as excessive growth reduction may occur.

Stop-drop sprays

If used properly, stop-drop sprays of naphthaleneacetic acid (NAA) can significantly reduce pre-harvest apple drop. Use knowledge of orchard conditions when applying stop-drop sprays and keep notes on the responses in your orchard.

Concentration And Timing Of Stop-Drop Application

| Variety | Application Time Before Picking (Days) | NAA Concentration (ppm) |
|------------------|--|-------------------------|
| Red Delicious | 7-10 | 10-15 |
| Jonathan | 7-10 | 10 |
| Golden Delicious | 7 | 10 |
| Rome Beauty | 7 | 10 |
| Winesap | 7 | 15 |

Apply NAA (Fruitone N, Amid-Thin W, K-salt Fruit Fix 800, K-salt Fruit Fix 200, and PoMaxa) before the beginning of fruit drop (7 to 14 days before harvest) at the rate of 5 ppm for summer varieties and 10 ppm for late varieties. This application should normally prevent fruit drop for 7 to 10 days.

Make a second NAA application within 7 to 10 days of the first application if fruits were not harvested. Do not exceed two NAA applications. Do not apply within two days of harvest. NAA works best as a dilute spray.

Using NAA too early, or in greater than recommended concentrations, may accelerate fruit maturity and decrease storage life. Apply stop-drop sprays at concentrations no higher than 3x. You may apply stop-drop sprays with pesticides. Do not use stop-drop sprays on trees in low states of vigor; healthy leaves are essential for these sprays to be effective.

ReTain

For harvest management

ReTain is labeled on apple, pear, peach, nectarine, plum, prune, and apricot. The active ingredient in ReTain is aminotheoxy-vinylglycine (AVG), a natural inhibitor of ethylene synthesis. Ripening fruit normally produces ethylene gas, which promotes further ripening and pre-harvest drop in some varieties. After treatment with ReTain, fruit produce less ethylene, which slows the ripening process and reduces pre-harvest drop.

Growers who have large plantings of a variety may consider applying ReTain to some of the planting as a harvest management tool to allow a later harvest of treated trees.

Timing. Best results are obtained when ReTain is applied before the first visible signs of ripening. Research has shown that ReTain should be applied four weeks before the optimum harvest date on apples and one or two weeks before the anticipated beginning of the normal harvest period for untreated fruit for other fruit types. Do not apply ReTain to plants of fruit under stress from heat, water, disease or insects as these plants may not respond to Retain. The PHI for ReTain is 7 days.

Application rate. Apply ReTain as a single or double application. The label rate is one pouch (11.7 oz.) or two pouches per acre. For apples apply one pouch per acre 28 days prior to the anticipated beginning of the normal harvest period for untreated fruit. A second application may be made to enhance the effects of ReTain. Apply one pouch of ReTain per acre 14 to 21 days after the first application, but not less than 7 days before harvest. The second application helps fruit retain their firmness during storage. Maintain spray solution pH between 6 and 8.

Dilute ReTain in at least 100 gal. of water per acre. Best results are obtained when applied under slow drying conditions in the morning or evening.

Thorough wetting and coverage are essential for optimum effectiveness. Although ReTain seems to be compatible with other materials, it should only be applied with other products if compatibility has been verified. Do not apply if rain is expected within eight hours of application.

Additives. Include a 100% organosilicone surfactant at 0.05 to 0.1 percent (v/v) (e.g., Widespread Max).

For increasing fruit set

ReTain is labeled to increase fruit set on apples, European pears, peaches, and cherries. Make a single application from pink to full bloom on apple, from white bud to full bloom on pear, and from balloon stage to first bloom on cherries. Never apply ReTain earlier or later than these stages.

Comments. ReTain is expensive; therefore, use it only on high-value, productive blocks with good fruit quality. Store treated fruit separately. Do not use an adjuvant for bloom applications. Do not apply more than two pouches of ReTain per acre per year.

ProVide 10SG on apples

Russetting

Applying ProVide, a mixture of gibberellins A4 and A7, reduces (but does not eliminate) russetting on Golden Delicious.

Apply ProVide two to four times with the first spray beginning at petal fall and continuing at 7- to 10-day intervals. The rate is 2.1 to 3.5 oz. (60-100) grams of ProVide applied in 100 gal of solution per application per acre. Spray additives are not required and have the potential to cause russetting.

Under conditions of high humidity and rain, you will obtain best russet control with four 3.5 oz. (100-gram) per-acre applications. Do not use excessive spray volumes, because excess moisture can induce russet. Direct 85 percent of the spray volume to the upper two-thirds of the tree.

Stayman cracking

Start applying ProVide 10SG three to six weeks before cracking begins (normally by mid-June to mid-July). Apply three to six consecutive sprays at 14- to 21-day intervals at an application rate of 1.8-3.5 oz. (50-100 grams) of ProVide per 50 gal. per half acre, per application. The use of a nonionic surfactant enhances spray coverage and improves absorption for cracking suppression. Do not use ProVide for cracking suppression if ProVide has been used in the same season on the same fruit for russet suppression. (REI is 4 hours.) Because weather changes influence cracking, and because cracking can occur over extended periods, multiple applications have given the best response. Apply ProVide early in the morning or late in the evening under slow drying conditions to maximize absorption.

Promalin on apples

Promalin contains 1.8% N-(phenylmethyl)-1H-purine 6-amine and 1.8 percent gibberellins A4 and A7. A single application to Delicious from early king bloom to the early stages of petal fall of the side blossoms elongate the fruit and encourages the development of more prominent calyx lobes.

The application rate is 1 to 2 pt. in 75 to 200 gal. of spray mixture per acre. If the bloom period is prolonged, two lower rate applications provide better results. Make the first application of 0.5 to 1 pt. of Promalin per acre at the beginning of the bloom period as above. Make the second application of 0.5 to 1 pt. of Promalin per acre three to 21 days later when the remainder of the canopy comes into bloom.

Do not exceed 2 pt. per acre for the combined sprays. Do not apply Promalin when air temperatures are below freezing or higher than 90°F.

Promalin for branching

Promalin may be used as a single application alone or in a Promalin-latex paint spot application to apples, non-bearing pears, and non-bearing sweet cherries. This treatment increases lateral bud break and shoot growth and improves branch angles to produce a stronger, better-shaped tree for early production. You must apply to non-bearing pear and sweet cherry one year before harvest.

You can make foliar Promalin applications on bearing and nursery apples and non-bearing pears at 1-3 inches of new terminal growth. The applications rate is 125-500 ppm (0.25-1 pt. of Promalin per 5 gal. of spray solution). On apple, non-bearing pears, and non-bearing sweet cherries, you may treat trees when they have reached a terminal height at which lateral branching is desired. For this treatment, apply Promalin at 250-1,000 ppm (0.5-2 pt.) per 5 gal. of spray solution.

You must make Promalin-latex paint applications before bud break or you may injure new shoot tips, causing shoot growth failure. Apply uniformly to cover the bark surface with a brush or sponge only on one-year-old wood. The application rate is 5,000 to 7,500 ppm (0.2-0.33 pt. or 3.2-5.3 fl. oz.) of Promalin per pint of latex paint.

Chemical thinning of apples

Chemical sprays can reduce fruit set on apples and promote larger fruit size at harvest and increase return bloom. These have become standard practices in most commercial orchards. Proper use is vital to the success of chemical thinning.

NAA (naphthalene acetic acid), NAD (naphthalene acetamide), Sevin (1-naphthyl-N-methyl-carbamate), and MaxCel (6-benzyladenine) are suggested.

Apply NAA to fall and winter varieties when king fruit are 11 to 13 mm in diameter. Sevin is more effective than NAA for fruit larger than 13 mm. Sevin gives uniform results from petal fall to 21 days later. NAD is most effective when applied from late bloom to petal fall. NAD is milder than NAA and is less likely to cause over-thinning.

You can combine NAA and Sevin, and apply it on fall and winter varieties when king fruit are 11 to 13 mm in diameter, and on summer varieties (Wealthy and Earliblaze) at petal fall.

Applying NAA to early summer varieties may result in excessive foliage injury, fruit cracking, and premature ripening.

In the warmer parts of the Midwest, concentrations of NAA that successfully thin frequently cause pygmy apples on spur-type Red Delicious. These small seedless apples persist through harvest and are a nuisance. Sevin is preferred for thinning spur-type Red Delicious. In some experiments, Sevin has over-thinned Rome and Gallia Beauty, so do not use it on these varieties. NAA is not successful in thinning Fuji, as this often results in pygmy apples. Honeycrisp is easy to overthin, so you should use combinations. In general, overapplication of NAA or use during warm temperatures can result in pygmy fruit regardless of variety.

You can avoid the variability of results and excessive foliage injury often experienced with NAA by using it at one-third and one-half of the rates recommended on the label in combination with 0.75 pt. of Tween 20 (also sold as Scattics, Akest TW 20) per 100 gal. Adding Tween 20 increases the rate of foliar absorption and decreases the effects of seasonal factors, such as temperature, relative humidity, and wind, on the drying rate and amount of material entering the leaf. Eliminating foliage wilting and tree "shock" results in better fruit size at harvest than the same amount of fruit thinning obtained by the full dosage of NAA alone.

Wetting agents that have been used successfully in tests in Illinois and Indiana include Regulaid and Nu Film 17.

MaxCel, Rite Way, Exilis 9.5 SC, Exilis Plus for thinning

Fuji thinning

Fuji has been notoriously difficult to thin in the past. On young trees MaxCel at 100 ppm (2 qt.) + 1 qt. Sevin XLR per 100 gal has worked well.

There are two options on mature trees with a heavy set:

1. 150 ppm (3 qt.) MaxCel + 1 qt. Sevin XLR
2. If **captan has not been used** prior to this during the current season: 100 ppm (2 qt) MaxCel + 1 qt. Sevin XLR + 1 qt. horticultural spray oil per 100 gal.

Accede® for cropload management in apple and peaches

Accede is a relatively new PGR containing the active ingredient 1-aminocyclopropanecarboxylic acid (ACC) from Valent Biosciences. This product has been shown to be especially useful in apples as a “rescue” thinner during the 15-20-mm fruitlet size window, which can also encourage greater return bloom. Apply 23 to 46 fl. oz. or 200 to 400 ppm of Accede in 100 gallons of water.

Accede can also be used as a chemical thinner in peaches and nectarines. This product is the first registered chemical thinner for Peaches. For peaches and nectarines, applications should be made from pink bud to full bloom.

Many conditions can affect the efficacy of this product such as cultivar, prevailing and anticipated climactic conditions, tree vigor, fruit set potential and orchard history. Research is still being conducted to determine the ideal use cases for this product. Use only as directed by the label.

Accede is not registered for use in the following states (IL, IA, IN, KY, NE, KS, OK).

Important reminders about chemical thinning

NAA generally gives best results under fast drying conditions and when the temperature is between 70°F and 75°F. Amid-Thin gives best results under slow drying conditions and is often applied in the evening.

Thorough spraying and uniform coverage are necessary for satisfactory results. However, to reduce the degree of thinning or avoid over-thinning, reduce the concentration (but not amount) per tree.

Lower limbs are easier to thin. Reduce spray application on lower limbs by shutting off one or more nozzles.

Some spray applied to the tree tops will fall on lower limbs.

Concentrate chemical thinner sprays have been satisfactory. Calibration allows the right amount of material to reach all parts of the tree and row. Avoid double applications to row ends, etc. Miscalibrating the sprayer manifold is magnified in concentrate application. Concentrating more than 4x has provided variable results, and should be avoided.

Applying chemical thinning sprays after frost or freezing temperatures is risky. Foliage exposed to such conditions absorbs chemicals more readily, and over-thinning may result. If you must spray under such conditions, reduce the concentration from 25 to 30 percent. Chemical thinners are generally more effective under the following conditions:

1. Low vigor trees
2. Light pruning
3. Heavy bloom
4. Poor pollination
5. High humidity before spraying
6. Slow drying of spray
7. Poor air drainage
8. Cloudy, cool weather preceding or following the bloom period

Keep records of the prevailing conditions when you make applications and leave several trees unsprayed to evaluate the results of thinning. This allows you to work out the concentrations best suited for your orchard. Using the carbohydrate balance model housed in NEWA or the Malusim app is an important tool which relates weather conditions to increases or decrease in thinner efficacy. This is an important tool to reduce over or under thinning when using chemical fruitlet thinners.

Defruiting young apple trees

It is often desirable to remove all the fruit from young trees when they have not reached a profitable bearing size. NAA at 15 ppm + Sevin XLR at 1 qt./100 gal applied at petal fall effectively defruits Jonathan, Red Delicious, and McIntosh. For other cultivars, use NAA at the recommended rate + Sevin XLR at 1 qt./100 gal. These sprays may not completely defruit the trees, but higher NAA rates may cause leaf damage.

Promoting return bloom

NAA may be used to promote return bloom for the following season in young trees that are not initiating enough flower buds and on older trees prone to biennial bearing that produce few flowers in off years such as Fuji, Jonagold, Mutsu, Braeburn and Golden Delicious. An application at a spray concentration of 3 to 5 ppm five to six weeks after full bloom in enough water to provide good coverage will increase bloom the following season. An additional one to two applications at 7- to 10-day intervals may be used for certain varieties that do not respond sufficiently based on previous experience.

Recommended Chemical Thinners For Apples¹

| Cultivars | NAD ^{2,3} (PPM) | NAA ^{2,4} (PPM) | NAA2 + WA ⁵ (PPM) | Sevin XLR Plus ^{2,6,7} (Qt./100 Gal.) | MaxCel ^{2,8} | Combinations ^{2,3,7,8} (PPM + Qt./100 Gal.) |
|------------------------|-----------------------------|-----------------------------|---------------------------------|---|-----------------------|---|
| Summer Varieties | 35-50 | | | | E | NAA 5-10 + Sevin 1/2-1 |
| Paulared | | 5-10 | 3-5 | 1/2-1 | M | |
| Gala | | 5-10 | 3-5 | 1/2-1 | M | |
| Jonamac | | 5-10 | 3-5 | 1/2-1 | M | |
| McIntosh | 35-50 | 7 1/2-12 | 3-5 | 1/4-1/2 | E | |
| Jonathan | 35-50 | 7 1/2-12 | 3-5 | 1/4-1/2 | E | |
| Spartan | | 10-15 | 5-7 1/2 | 1/2-1 | ? | |
| Cortland | 35-50 | 7 1/2-12 | 3-5 | 1/4 | E | |
| Grimes Golden | 35-50 | 5-10 | 5-7 1/2 | | ? | NAD 25-50 + Sevin 1/2-1 |
| Red Delicious/non-spur | | 5-10 | 3-5 | 1/2-1 | E | |
| Red Delicious/spur | | 5-10 | 5-7 1/2 | 1/2-1 | M | |
| Honeycrisp | | 3-5 | | 1/4-1/2 | ? | N.R. ⁹ |
| Empire | | 10-15 | 5-7 1/2 | 1/2-1 | E | |
| Golden Delicious | | 10-20 | 5-10 | 1/2-1 | M | NAA 5-10 + Sevin 1/2-1 |
| Blushing Golden | | | | 1/4-1/2 | ? | |
| Firmgold | | | | 1/4-1/2 | ? | |
| Idared | | | | 1/2-1 | E | |
| Winesap | 35-50 | 7 1/2-10 | 3-5 | 1/2-1 | E | |
| Stayman and Turley | 35-50 | 7 1/2-10 | 3-5 | 1/2-1 | M | |
| Braeburn | | 7 1/2 | 7 1/2 | | | NAA 7 1/2 + Sevin 1 |
| Rome | 50-60 | 15-20 | 7 1/2-10 | N.R. ⁹ | E | |
| Fuji ⁸ | | N.R. ⁹ | | | H | MaxCel 150 ppm + Sevin 1 |

¹ Thinning Strategies for 2022 by Anna Wallis, Philip Schwallier, and Amy Irish-Brown. Available online at: <https://www.canr.msu.edu/apples/uploads/files/Thinning%20Guide%202022%20updated.pdf>

² Lower concentrations suggested when conditions are favorable for thinning.

³ Apply NAD (Amid-Thin) from late bloom to petal fall.

⁴ Apply NAA, Sevin, or combinations to fall and winter varieties when king fruits are 11-13 mm in diameter. On summer varieties (such as Wealthy and Earliblaze) apply the combination at petal fall.

⁵ WA = wetting agent: Regulaid at 0.5 pt per 100 gal.

⁶ Adding NAA at 2.5-4 ppm to Sevin stimulates the initiation of fruit buds for return bloom. This low-NAA rate should not thin fruit or cause pygmy apples on Red Delicious.

⁷ The Sevin XLR Plus formulation is most commonly used for thinning and is the only formulation labeled for early use (80% petal fall to 6 mm fruit diameter). Reduce spray application to lower portion of tree to avoid overthinning. Consult the label if you use other Sevin formulations.

⁸ Variety ease of thinning with MaxCel: **E** = easy. **M** = moderate. **H** = hard. See MaxCel Recommendation Tables for suggested rates of Maxcel and Sevin for thinning.

⁹ N.R. = not recommended.

MaxCel For Apples And Pears

| Use | For Fruit Thinning, Sizing, And Enhanced Return Bloom |
|--------------|--|
| Application | Apply 75 to 200 ppm spray concentration. Refer to dilution table on label for assistance. |
| Spray Volume | Use sufficient volume to ensure complete tree coverage. |
| Spray Timing | Apply when average king fruit diameter is 5-15 mm. 10 mm is optimal. Do not apply more than twice in a season. |

MaxCel For Thinning Apples Only¹

| Thinning Difficulty | Aggressive | Moderate | Slight |
|---------------------|---------------------------|--------------------|-----------|
| Hard to thin | 100-150 ppm + Sevin + oil | 100 ppm + Sevin | 100 ppm |
| Moderate to thin | 100 ppm + Sevin | 75-100 ppm + Sevin | 75 ppm |
| Easy to thin | 75-100 ppm + Sevin | 75 ppm | 50-75 ppm |

¹ See Recommended Chemical Thinners for Apples for variety thinning difficulty rating.

MaxCel Dilution Table¹

Fluid ounces of MaxCel per 100 gallons of spray required to obtain given ppm concentrations.

| 10 PPM | 25 PPM | 50 PPM | 75 PPM | 100 PPM | 125 PPM | 150 PPM | 175 PPM | 200 PPM |
|--------|--------|--------|--------|---------|---------|---------|---------|---------|
| 6 | 16 | 32 | 48 | 64 | 80 | 96 | 112 | 128 |

¹MaxCel contains 75 grams active ingredient per 128 fl. oz. (1 gal.).

NAA Formulations

Not all NAA formulations have the same amount of active ingredients. Because calculating ppm can be difficult, this table describes materials and amounts of formulation per 100 gal of water required to make a 10 ppm solution (table developed by R. Marini, VPI).

NAD And NAA Formulations For Chemical Thinning Of Apples

| Trade Name | Chemical | Formulation | Acid Equivalent (% Of Active Ingredient) | Amount Of Formulation Per 100 Gal. To Make 10 PPM |
|----------------------|---|-------------|--|---|
| Amid-Thin W | 1-Napthaleneacetamide | WP | 8.4 | 1.6 oz. |
| Fruitone N | 1-Napthaleneacetic acid, sodium salt (3.5%) | WP | 3.1 | 4 oz. |
| PoMaxa | 1-Napthaleneacetic acid, sodium salt (3.5%) | liquid | 3.1 | 4 fl. oz. |
| K-salt Fruit Fix 800 | 1-Napthaleneacetic acid, potassium salt (24.2%) | liquid | 20.2 | 0.63 fl. oz. |
| K-salt Fruit Fix 200 | 1-Napthaleneacetic acid, potassium salt (6.25%) | liquid | 5.18 | 2.47 fl. oz. |

Promoting branching on high-density dwarf fruit trees

Branch inducing growth regulators contain cytokinins, gibberellins or both. Cytokinins stimulate cell division and differentiation and promote shoot initiation and release of lateral buds from apical dominance; they also play an indirect role in overcoming dormancy.

Gibberellins, like cytokinins, promote cell division while also stimulating elongation to increase shoot length. However, at rates high enough to stimulate growth, gibberellins will entirely de-fruit trees and inhibit return bloom. Growers who wish to promote lateral branching, but who also desire a potential crop the following year to manage vigor, should apply a cytokinin only bioregulator. Lower rates of active ingredient used to moderately thin fruit, reduce russetting, improve size or shape, or to increase fruit set after frost will not diminish return bloom.

The following are some chemical plant growth regulators labeled for lateral branch promotion on tree fruits. Be sure to check the product registration as some are not available for distribution in all states. Cytokinins: Maxcel® – Valent Biosciences, LLC (not registered in KS, NE, OK); Cytokin® – Miller Chemical & Fertilizer, LLC (not registered in KY, KS, MN, NE); Exilis® 9.5 SC – Fine Americas, Inc. (not registered in AR, IN, KY, KS, MN, NE, OK). Cytokinins and gibberellins: Promalin® – Valent BioSciences, LLC; Typy® – Nufarm Americas,

Inc.; Cytoplex® HMS® – Miller Chemical & Fertilizer, LLC (not registered in IA, KY, KS, MN, NE, OK, WV); Perlan® – Fine Americas, Inc. (not registered in AR, IA, IN, IL, KY, KS, MO, NE, OH, OK).

Foliar sprays are effective starting when terminal growth is 1-3 inches long at the time of application.

Rates for non-bearing trees are generally between 125 and 500 ppm for apples and 250 and 1,000 ppm for pears and sweet cherries. For first-leaf apple trees received from the nursery as whips, apply 500 to 1,000 ppm of MaxCell 20 days after bud break followed about a week or so later by removal of the 2nd through 4th apical buds on the leader. For second and third-year apple trees, apply 200-300 ppm (non-bearing) or 200 ppm (bearing) of MaxCel. Spreading of some rapidly growing shoots is usually necessary, but for most, the crotch angles that develop are wider compared to other branch-promoting techniques and require less manual labor for correction.

For further information on branch promotion see <http://www.uky.edu/hort/sites/www.uky.edu/hort/files/documents/HortFact3011-4.pdf>

Managing the shoot blight phase of fire blight with prohexidione-calcium (Pro-Ca)

Prohexidione-calcium (Pro-Ca, Apogee) is used on apple trees for three reasons:

- To reduce shoot growth because trees are overly vigorous because of crop loss, inappropriate root-stock, tree spacing, or excessive nitrogen fertilization.
- To reduce the risk and reduce susceptibility to fire blight.
- Pro-Ca is available as Apogee and various other products. See generics table for a listing of Pro-Ca products.

Pro-Ca inhibits gibberellin biosynthesis, which stops terminal growth early.

Pro-Ca can decrease the length of shoots by 30 to 60 percent. Pro-Ca does not affect blossom blight occurrence, but when used effectively, reduces the occurrence of shoot blight. Shoots with inhibited growth are less susceptible to fire blight. The decrease in blight susceptibility does not occur until about 10 to 14 days after application. Note that apple varieties differ in their susceptibility to damage from Pro-Ca. Do not apply Pro-Ca to Empire or Stayman varieties, because it causes cracking on these varieties.

Consider using Pro-Ca to reduce the threat of shoot blight on vigorous trees of susceptible varieties that have nearly filled their tree space. Pro-Ca only decreases host susceptibility; it does not affect the pathogen directly. Pro-Ca is not a substitute for streptomycin for blossom blight control during bloom. If needed, you can combine Pro-Ca with streptomycin in one of the bloom sprays.

In general, the less shoots have developed before the application of Pro-Ca is made the greater effect it will have on shoot extension. Follow the label of the particular Pro-Ca product, however aiming for applications at the pink stage of floral development is a good benchmark.

Split applications have been shown to provide longer shoot suppression during the summer. Vegetative suppression lasts two to five weeks only during the

current growing season. When fire blight is a concern, increase the first Pro-Ca application to help control vigor early and reduce the risk of fire blight.

Pro-Ca is locally systemic. This means if you spray the tops of trees, Pro-Ca has an effect only on the tops where it was applied. This allows growers to use Pro-Ca in problematic areas of trees that need localized control. For example, if the bottoms of trees had frost damage that resulted in a low crop load, applying Pro-Ca to the bottoms will control the excessive growth. Tree vigor, variety (see table below), crop load, and application timing affect the efficacy of Pro-Ca and the rate needed.

Prohexidione-calcium (Pro-Ca) application rates for vegetative growth control in apples

Pro-Ca can cause a tree to retain more fruit and thinning programs will need to be adjusted for this. To decrease June drop on trees with light bloom, apply Pro-Ca at 1-3 inches of new shoot growth. Consult label of your specific product for rates and application instructions. Applying Pro-Ca in the same season as ProVide reduces the effectiveness of both growth regulators.

Timing. Pro-Ca is considerably less effective if applied too late. Apply at the earliest labeled application window for your specific product for optimum results. The decrease in blight susceptibility does not occur until about 10 to 14 days after application.

Additives. Use a nonionic surfactant with Pro-Ca. Follow the manufacturer's rate recommendations. If you mix Pro-Ca in hard water (water that contains high levels of calcium carbonate), add 1 lb. of spray-grade ammonium sulfate for each pound of Pro-Ca.

Comments. Pro-Ca's ability to control growth does not depend on concentration. There is no difference in shoot growth control between dilute and concentrate sprays, provided the total amount of chemical per acre

Selected Apple Cultivars And Their Sensitivity To P Pro-Ca (Apogee)

| Pro-Ca Sensitivity | Cultivars | Recommendation |
|--------------------|--|--|
| Very sensitive | Cortland, Gala, Ginger Gold, Northern Spy, Paula Red, Rome | Consider reducing spray rates. |
| Sensitive | Fuji, Golden Delicious, Jonamac, Spartan, | |
| Less sensitive | Golden Supreme, Jonagold, Jonathan, IdaRed, McIntosh | Consider using an additional 1 oz. per acre |
| Special | Empire, Stayman, Winesap | On spur type, begin first application 2 weeks after bloom, followed by an application 2 weeks later, and then 2 weeks after for medium-size trees. |
| Phytotoxic | Empire, Stayman, Winesap | Do not use. |

is the same. Pro-Ca's level of growth control is rate dependent. The recommended rate provides the greatest and quickest reduction, and the effect on growth declines as the rate is reduced. Do not tank mix Pro-Ca with boron, calcium chloride, or calcium nitrate. Pro-Ca is rainfast after 8 hours. (REI is 12 hours; PHI is 45 days.)

Cork spot and bitter pit management in apples

Cork spot and bitter pit are related to low levels of calcium and high levels of nitrogen in the fruit. However, low calcium is not the only cause of these disorders. Excessive tree vigor and a light fruit crop increase cork spot and bitter pit. Bitter pit is primarily a storage disorder, and calcium treatments before and after harvest can reduce this.

No one cultural practice eliminates these disorders and you need a multifaceted approach for control. For example:

- Apply lime to raise the soil pH to around 6.5 to make calcium more available for tree uptake.
- Balance tree nutritional levels by analyzing soil and tissue. Keep nitrogen, potassium, and magnesium levels from becoming excessive and avoid low levels of calcium, boron, and zinc. If tissue analysis boron levels are low, apply Solubor at 2 lb./acre at pink and again at petal fall.
- Work to moderate tree vigor. Avoid excessive pruning and tree overcrowding and make moderate nitrogen applications. Summer pruning of water sprouts between mid-July and mid-August also helps minimize cork spot.
- Adjust fruit density by chemically thinning fruit in heavy cropping years to avoid a light crop and calcium disorders the following year.
- Apply calcium in 6-8 sprays starting at first cover. Calcium chloride is most commonly used; it is inexpensive but can be hard on pumps. Calcium nitrate (Nutrical) is an alternative. Calcium chloride is sold as dry formulations (such as DowFlake Xtra 83-87% and Cor-Clear 28%) and as liquid formulations with 10 percent calcium (such as EezyCal 8-0-0-10 and Loveland 10% Calcium). Applying calcium at a low rate every 7 days is more effective than a higher rate every 14 days. Application between first cover and third cover is most critical, but later sprays also help. You can apply higher rates after mid-July.

Managing cork spot in Honeycrisp

Honeycrisp is one of the cultivars most susceptible to corking. Affected cells start to show damage around two weeks after petal fall, but visual symptoms usually show up in mid to late June. When the disorder starts,

damaged cells usually have higher rates of protein synthesis, respiration, and cell division, but these cells become brown and die shortly thereafter. As fruit development progresses, severely damaged fruit become cracked and deformed with deep brown, cork-like areas scattered throughout the flesh.

Calcium and boron deficiencies are suspected as the main reasons for cork spot development. The flesh and peels of Honeycrisp apples have less calcium than some other varieties, so keeping up with the calcium sprays is essential for managing cork spot. Calcium moves very slowly into the tissue, so more water and good coverage are essential for better uptake.

The recommended application rate is about 1.5-2 pounds of calcium chloride dissolved in 400 gallons or more of water per acre. Begin applying calcium in the first cover spray. Apply this spray to runoff.

Disease Management Notes

Note About Mancozeb (EBDC Products)

EBDC products have two rate recommendations, depending on how you use the fungicides.

Mancozeb cannot be used past bloom at the 6 lb. per acre rate, but is permitted past bloom at the lower rate of no more than 3 lb. per acre. However, this lower rate may be insufficient under heavy scab pressure without a tank-mix partner. The application of mancozeb from bloom through first cover for the control of apple scab heavily contributes to the prevention of bitter rot in highly susceptible varieties, like Honeycrisp.

- Do not apply mancozeb within 77 days of harvest.
- Do not apply more than 24 lb. of mancozeb (Roper and other generic formulations) or more than 25.6 lb. of penncozeb per acre per year if using for pre-bloom (6 lb.) applications.
- Do not apply more than 21 lb. of mancozeb or more than 22.4 lb. of Penncozeb per year if using the 3 lb. per acre rate.
- Label recommendations for mancozeb are identical for apples and pears.

Use of Captan fungicide on tree fruit-restricted entry intervals (REI)

Most captan formulations (Captan 50W, Captan 80WDG, Captan 4L) are currently available with a 24-hour REI. The REI was reduced from 4 days to 24 hours a few years ago for apples, cherries, plums/fresh prunes, and peaches. However, some formulations produced by certain companies still may have the 4-day REI. Check the label of the captan product you plan to purchase to be sure it has a 24-hour REI.

Fungicide resistance management

Many of our “best” fungicides may lose their efficacy due to fungicide resistance evolving in the pathogens we are trying to manage. These fungicides include Topsin-M, Scala, Vangard, the sterol-inhibiting fungicides (Rally, Rhyme, Rubigan, Indar, Inspire Super, Procure, Topguard); the strobilurin fungicides (Sovran, Pristine, Merivon, and Flint); and the succinase dehydrogenase inhibiting fungicides (Aprovia, Fontelis, Luna Sensation, Luna Tranquility, Pristine and Merivon, and Syllit (dodine)). Because they all have specific modes of action, fungi such as the apple scab, bitter rot and powdery mildew pathogens can rapidly evolve resistance to them. Fungicide resistance, or at least reduced sensitivity, has been observed for apple scab, bitter rot and powdery mildew to both the sterol-inhibitor and strobilurin fungicides in the United States and throughout the Midwest.

To delay resistance development, never use these fungicides alone in a season-long program and use them as little as possible. Most of the newer fungicides limit the number of applications that can be made per season (generally no more than four), and labels state that no more than two sequential applications should be made without alternating with another fungicide with a different mode of action. The sterol-inhibiting fungicides, strobilurin fungicides, and succinase dehydrogenase inhibiting fungicides have different modes of action and can be alternated with each other in a fungicide resistance management program.

A good approach is to alternate one to two spray blocks of these materials. For example: a spray of Sovran (a strobilurin fungicide) alternated with a spray that contains Rally (a sterol-inhibiting fungicide) mixed with a broad-spectrum protectant fungicide such as captan or mancozeb.

Use of pre-mixes

Many chemical manufacturers provide pre-mixes (pre-formulated tank mixtures). Take care when using these pre-mixes so that your rotation partners are not in the same chemical family as the pre-mix. For example, if you use Pristine or Merivon (a pre-mix of a strobilurin and SDHI fungicide; FRAC11+7), avoid using Sovran or Flint (FRAC11), or the Luna series (FRAC 7+11; FRAC 7;FRAC 7+9), all of which contain either a strobilurin (FRAC 11) or an SDHI (FRAC 7). Do not rotate with Aprovia, Fontelis, Kenja or Miravis, all of which are SDHI fungicides (FRAC 7).

Blister spot on Mutsu, Cortland, Fuji

Blister spot is a bacterial disease of susceptible apple varieties — most notably Mutsu (Crispin). It is caused by a bacterium in the genus *Pseudomonas*. New blister

spot outbreaks have been identified on Cortland and Fuji, and outbreaks have been reported on other varieties interplanted with Mutsu.

Due to resistance issues, streptomycin is no longer suggested for disease management. Due to the severity and lack of control options for blister spot on Mutsu, the variety we recommend replacing Mutsu with Shizuka, which is similar in color and quality to Mutsu.

Sanitation methods to aid in apple scab and bitter rot control

Especially in years after a high incidence of apple scab developed in the orchard, sanitation is important because apple scab overwinters only in fallen leaves. Similarly, bitter rot pathogens overwinter in mummified apples and dead woods.

The sanitation methods described below can reduce the amount of apple scab inoculum (ascospores) by as much as 50 percent.

Applying 5% urea to the orchard floor (40 lb. per acre in 100 gal. of water) provides nitrogen to help microorganisms decompose leaves, killing the overwintering apple scab fungus.

Flail mowing the orchard also has been reported to reduce apple scab inoculum by as much as 50 percent.

You can flail mow or apply nitrogen in the fall and/ or spring. Each method has been reported to reduce the number of scab ascospores by as much as 50 percent; however, the combined effects do not provide complete control. Using both methods probably does not reduce ascospore more than 50 percent. Be sure to recognize that urea provides nitrogen, and modify your fertilization program appropriately.

Removing mummified fruits and dead woods before bloom and destroying (e.g., burning) them reduce initial inoculum for fruit infection.

Insect Management Notes

Insecticide resistance in codling moth populations

Several states (including those covered by this guide) have reported codling moth populations that are suspected or confirmed to be resistant to certain insecticides. The resistance traits of populations differ among orchards and regions, so resistance may account for control failures in some orchards, even though the same insecticides may provide effective control in other locations.

Resistance is not the only cause for control failures, so always consider whether the cause of poor control was due to other issues, including inadequate rates,

inadequate spray volumes, spray timing, or wash-off due to rainfall. Where these factors do not appear to explain poor control, resistance — particularly to the organophosphates (Imidan, Diazinon) — may be the reason, and switching to other insecticides is recommended. Where control programs have been effective and resistance does not seem to be a problem, rotating among insecticides with different modes of action is recommended to delay resistance development.

Effectiveness ratings of insecticides for control of codling moth are shown in the table for first and second cover sprays, on pages 35-37.

Insect populations resistant to the organophosphates exhibit resistance to all the organophosphates that are labeled for codling moth control in apples (Diazinon, Imidan), so switching among these insecticides offers no benefit. Laboratory research and field observations have shown that organophosphate-resistant codling moth populations also are less susceptible to some pyrethroids, so switching to Pounce (or other permethrin formulations), Asana, Warrior, Danitol, Mustang Maxx, or Baythroid may not provide adequate control.

Altacor, Assail, Delegate, Exirel and Rimon are effective against organophosphate-resistant codling moth populations. Consult your state Extension specialists in entomology to plan effective seasonlong programs that make the best use of available products within the label-specified limits and restrictions for each.

Timing Of First Insecticide Spray For Codling Moth Control On Apple And Pear¹

| Degree-days (Base 50°F) After Biofix ² | Insecticide Products |
|---|---|
| 50-75 | Dimilin Rimon |
| 100-200 | Intrepid Confirm |
| 150-250 | Altacor Assail Belay Delegate Exirel |
| 250 | Imidan Avaunt Pyrethroids (Asana, Baythroid, Danitol, Mustang Max, Proaxis, Warrior) Virus (Cyd-X, Carpovirusine, Madex) |

¹ A second spray should be made 10-14 days later.

² Biofix is defined as the date on which pheromone traps detect sustained flight of moths

Apple borers

The dogwood borer and American plum borer are caterpillars that attack burr knot tissue on apple trunks.

Flat-headed and round-headed apple borers are beetle larvae that attack tree trunks, often trees that have received mechanical, cold, or other injury or are generally weakened.

Insecticides currently registered for control of borers on apple are Assail 30SG for dogwood borer and Warrior II for control of tree borer species.

For dogwood borer, the best insecticide timing is at peak egg hatch, which is in late June in the central Midwest. Pheromone mating disruption by Isomate DWB can be used starting at bloom.

For American plum borer, the best timing is at petal fall.

For flat-headed and round-headed apple borers, apply insecticide in the spring.

Apply borer sprays to the lower 4 feet of the trunk and lower branches and soak the bark.

Periodical cicadas

Periodical cicadas are orange to black and about 1 1/2 inches long, have black transparent wings, and appear from May to July. Annual or dog-day cicadas are larger, green to black, and appear each year from July to September. Annual cicadas ordinarily do not cause much damage. Cicada males announce their presence to the voiceless females by making a continuous, high-pitched, shrill sound.

Adult females lay eggs in rows in pockets they cut in small branches and twigs of trees with their long, knife-like egg layer. The eggs hatch in six or seven weeks. The newly hatched nymphs fall to the ground and burrow until they find suitable roots, usually 1 1/2 to 2 feet beneath the soil. With their sucking mouthparts, they immediately begin to suck juices from the roots.

Females prefer oak, hickory, apple, peach, and pear trees, and grapevines for laying eggs. Females damage plants when they make slits in branches and twigs to deposit their eggs. These small twigs and branches turn brown and die and sometimes break off. The damage may be severe in newly planted orchards or on new shade trees or shrubs. Heavy populations of nymphs in the soil also may affect the growth and vigor of certain trees.

You can prevent egg-laying damage by cicadas on young fruit and ornamental trees by covering them with a protective netting, such as cheesecloth. Cover a tree and tie the netting to the trunk below the lower branches. Remove the covering when egg-laying is over. If netting is not an option, you may apply insecticides when egg laying begins and repeat 7 to 10 days later. Pyrethroids are recommended to control periodical cicada, but using these products may lead to mite outbreaks.

Notes on soaps and horticultural oils

SunSpray UFO (UFO = “ultrafine” oil), Saf-T-Side, and M-Pede (a potassium salt of fatty acids, previously called an insecticidal soap) are relatively new insecticides that may be used in certified organic production systems. Summer oils and M-Pede are effective only against insects the sprays contact at the time of application. These sprays provide no residual control. Many questions about their efficacy remain, and their use should be considered experimental.

Nonetheless, they appear to be useful in certain situations.

A summer oil alone, at a concentration of 1-2 percent by volume, provides some control of mites and aphid (rosy apple aphid, apple grain aphid, green apple aphid, and spirea aphid). Limited observations suggest that aphid control is likely to be greatest if you apply oil when clusters are at the 0.25 inch green stage.

M-Pede alone reduces mite, aphid, pear psylla, and white apple leafhopper populations, but control may not be satisfactory or long-lasting unless you apply multiple sprays. Unlike oils, M-Pede is not ovicidal.

If applied alone, a summer oil is likely more effective for aphid and (especially) mite control than M-Pede. Data from Michigan indicate that adding M-Pede at 2 percent by volume to full-rate sprays of Vendex, Kelt-hane, and presumably other miticides, greatly enhances the control they provide.

Phytotoxicity, leaf drop, and fruit blemishes should be major concerns when deciding whether to use summer oil or soap. To prevent damage to foliage or fruits, never use a summer oil with Captan, Sevin, or other sulfur-containing pesticides. Allow at least 14 days between applications of sulfur-containing compounds and the use of a summer oil. Do not apply oils if temperatures exceed 90°F or drying conditions are poor.

Because of concerns about fruit russetting, some authorities suggest that insecticidal soaps should be used only in nonbearing orchards. Applicators must mix oils and soaps at the proper dilution (1-2 percent); concentrated sprays are less effective and more phytotoxic. Deposits of large droplets or the coalescing of droplets on fruit or foliage also increases the likelihood of leaf damage and fruit blemishes.

Effectiveness Of Insecticides For Control Of Minor Apple Insects¹

| Product And Formulation Active Ingredient | IRAC Code ² | Black Stem Borer | Green Aphid | Oblique Banded Leafroller | Potato Leafhopper | Redbanded Leafroller | White apple Leafhopper | REI ³ PHI ⁴ | Max amt ⁵ Max app ⁶ |
|--|---------------------------|---------------------|----------------|---------------------------------|----------------------|-------------------------|---------------------------|--------------------------------------|--|
| Actara (25WDG) | 4A | x | E | x | u | x | E | 12h | 16.5 oz. |
| thiamethoxam | | | | | | | | 14-35d | NA |
| Actara (25WDG) | 4A | x | E | x | u | x | E | 12h | 16.5 fl. oz. |
| thiamethoxam | | | | | | | | 14-35d | NA |
| Admire Pro (4.6F) | 4A | x | E | x | u | x | E | 12h | 10.5 fl. oz. |
| imidacloprid | | | | | | | | 7d | NA |
| Agri-Mek SC (0.7SC) (RUP) | 6 | x | x | x | x | x | G | 12h | 8.5 fl. oz. |
| abamectin | | | | | | | | 28d | 2 |
| Altacor eVo | 28 | x | x | E | x | E | s | 4h | 4.6 fl. oz. |
| chlorantraniliprole | | | | | | | | 5d | 3 |
| Apta (1.34SC) | 21A | x | G | G | u | u | u | 12h | 53.5 fl. oz. |
| tolfenpyrad | | | | | | | | 14d | 2 |
| Asana XL (0.66EC) (RUP) | 3A | x | F | G | x | E | G | 12h | 101.5 fl. oz. |
| esfenvalerate | | | | | | | | 21d | NA |
| Assail 30SG | 4A | x | E | x | G | x | G-E | 12h | 32 oz. |
| acetamiprid | | | | | | | | 7d | 4 |
| Avaunt eVo | 22A | x | x | x | G | G | s | 12h | 24 oz. |
| indoxacarb | | | | | | | | 14d | 4 |

(Continued)

Effectiveness Of Insecticides For Control Of Minor Apple Insects¹ (continued)

| Product And Formulation Active Ingredient | IRAC Code ² | Black Stem Borer | Green Aphid | Oblique Banded Leafroller | Potato Leafhopper | Redbanded Leafroller | White apple Leafhopper | REI ³ PHI ⁴ | Max amt ⁵ Max app ⁶ |
|---|---------------------------|---------------------|----------------|---------------------------------|----------------------|-------------------------|---------------------------|--------------------------------------|--|
| <i>Bacillus thuringiensis</i> (B.t.) (Agree, Dipel, etc.) | 11B | x | x | G | x | G | x | 4-12h | NA |
| <i>Bacillus thuringiensis</i> | | | | | | | | 0d | NA |
| Baythroid XL (1EC) (RUP) | 3A | x | x | G | G | E | G | 12h | 2.8 fl. oz. |
| cyfluthrin | | | | | | | | 7d | NA |
| Belay (2.13SC) | 4A | x | E | u | u | x | E | 12h | 12 fl. oz. |
| clothianidin | | | | | | | | 7d | NA |
| Beleaf 50SG | 29 | x | G | x | x | x | x | 12h | 8.4 oz. |
| fonicamid | | | | | | | | 21d | 3 |
| Centaur WDG (70WDG) | 16 | x | x | x | F | x | F | 12h | 34.5 oz. |
| buprofezin | | | | | | | | 14d | 1 |
| Confirm 2F | 18 | x | x | F | x | E | x | 4h | 120 fl. oz. |
| tebufenozide | | | | | | | | 14d | NA |
| Damoil | UN | x | u | x | x | x | x | 4h | NA |
| mineral oil | | | | | | | | NA | NA |
| Danitol 2.4EC (RUP) | 3A | x | x | G | E | E | E | 24h | 42.6 fl. oz. |
| fenpropathrin | | | | | | | | 14d | NA |
| Delegate WG (25WG) | 5 | x | x | E | x | E | x | 4h | 28 oz. |
| spinetoram | | | | | | | | 7d | 4 |
| Delta Gold (1.5EC) (RUP) | 3A | x | x | u | u | u | u | 12h | 3.6 fl. oz. |
| deltamethrin | | | | | | | | 21d | NA |
| Entrust SC (2SC) | 5 | x | x | G | x | x | x | 4h | 29 fl. oz. |
| spinosad | | | | | | | | 7d | 4 |
| Esteem 35WP, 0.86EC | 7C | x | u | s | x | s | x | 12h | 10 oz. |
| pyriproxifen | | | | | | | | 45d | 2 |
| Exirel (0.83SE) | 28 | x | x | E | x | E | G | 12h | 61.6 fl. oz. |
| cyantraniliprole | | | | | | | | 3d | 3 |
| Imidan 70W | 1B | x | x | x | x | G | x | 168h | 15.5 lb. |
| phosmet | | | | | | | | 7d | NA |
| Intrepid 2F | 18 | x | x | E | x | E | x | 4h | 64 fl. oz. |
| methoxyfenozide | | | | | | | | 14d | NA |
| Lannate SP (90WSB) (RUP) | 1A | x | x | E | x | E | E | 72h | 5 lb. |
| methomyl | | | | | | | | 14d | 5 |
| Movento (2SC) | 23 | x | G | x | x | x | x | 24h | 25 fl. oz. |
| spirotetramat | | | | | | | | 7d | NA |
| Mustang Maxx (0.83EC) (RUP) | 3A | x | x | G | E | E | E | 12h | 24 fl. oz. |
| zeta-cypermethrin | | | | | | | | 14d | NA |
| Neemix 4.5 (0.39L) | UN | x | F | u | u | u | u | 4h | NA |
| azadirachtin | | | | | | | | 0d | NA |

(Continued)

Effectiveness Of Insecticides For Control Of Minor Apple Insects¹ (continued)

| Product And Formulation Active Ingredient | IRAC Code ² | Black Stem Borer | Green Aphid | Oblique Banded Leafroller | Potato Leafhopper | Redbanded Leafroller | White apple Leafhopper | REI ³ PHI ⁴ | Max amt ⁵ Max app ⁶ |
|--|---------------------------|---------------------|----------------|---------------------------------|----------------------|-------------------------|---------------------------|--------------------------------------|--|
| Permethrin 25W (RUP) | 3A | x | G | E-G | x | E | x | 12h | 32 oz. |
| permethrin | | | | | | | | 14d | NA |
| Permethrin 3.2EC (RUP) | 3A | x | x | x | x | E | P | 12h | 20 fl. oz. |
| permethrin | | | | | | | | 14d | NA |
| Portal XLO (0.4EC) | 21A | x | x | x | u | x | F | 12h | 2 pt. |
| fenpyroximate | | | | | | | | 14d | 1 |
| PQZ (1.87SC) | 9B | x | E | x | x | x | x | 12h | 4.8 fl. oz. |
| pyrifluquinazon | | | | | | | | 14d | 2 |
| Proclaim (5SG) (RUP) | 6 | x | x | E | x | E | x | 12h | 14.4 oz. |
| emamectin benzoate | | | | | | | | 14d | NA |
| Rimon 0.83EC | 15 | x | x | E | x | E | u | 12h | 150 fl. oz. |
| novaluron | | | | | | | | 14d | NA |
| Sevin XLR Plus | 1A | x | F | F | x | F | x | 12h | 15 qt. |
| carbaryl | | | | | | | | 3d | 8 |
| Sivanto Prime (1.67SC) | 4D | x | G | x | G | x | G-E | 4h | 28 fl. oz. |
| flupyradifurone | | | | | | | | 14d | NA |
| Surround WP (95WP) | UN | x | x | s | x | s | x | 4h | NA |
| kaolin | | | | | | | | 0d | NA |
| Verdepryn 100SL (0.83SL) | 28 | x | x | x | x | E | u | 4h | 33 fl. oz. |
| cyclaniliprole | | | | | | | | 7d | 3 |
| Versys Inscalis (0.83DC) | 9D | x | G | x | x | x | x | 12h | 7 fl. oz. |
| afidopyropen | | | | | | | | 7d | NA |
| Vydate L (2L) (RUP) | 1A | x | G | x | F | x | E | 48h | 8 pt. |
| oxamyl | | | | | | | | 14d | 4 |
| Warrior II (2.08CS) (RUP) | 3A | x | G | F | x | E | x | 24h | 12.8 fl. oz. |
| lambda-cyhalothrin | | | | | | | | 21d | NA |

Footnotes: See how to read the spray schedule tables section at the beginning of the chapter.

2. PEAR

Pear Spray Schedule

The boxes represent the crop stages where common pests in the Midwest are active. Scouting and/or preventative sprays may be necessary or recommended. Weather and degree day accumulation will impact the exact timing of pest appearance in the orchard.

| Stage | | | | | |
|-------------------------|----------------|-------|---------------------|------------------------|---------------|
| Dormant Until Bud Burst | Pre-Bloom | Bloom | Petal Fall | First And Second Cover | Summer Covers |
| San Jose Scale | | | | San Jose Scale | |
| Pear Psylla | | | | Pear Psylla | |
| | | | | Codling Moth | |
| | Stink Bugs | | | | Stink Bugs |
| European Red Mite Eggs | | | | European Red Mite | |
| Pearleaf Blister Mite | | | | | |
| | Pear Rust Mite | | | Pear Rust Mite | |
| | | | Tarnished Plant Bug | | |
| | | | Plum Curculio | | |
| | | | | Periodical Cicada | |
| | | | | | Mealybug |

How to read the spray schedule tables

Every pear growth stage has important notes on disease or insect management. In some cases, the reader will be directed to the special problems section at the end of the section or chapter. Please make sure to read thoroughly and contact your state Extension specialist with any specific questions.

Key to tables

E = excellent control

G = good control

F = fair control

[r] = fungicide/insecticide resistance possible

s = suppression only

i = ineffective

u = unknown efficacy

x = pest not on the label

¹ Efficacy data in this publication are based on trials conducted across various regions and does not necessarily reflect local efficacy differences or changes over

time. Growers should contact their Extension specialist for the most recent or for state-specific information. The information on this publication is only a guide; the authors and their institutions assume no liability for practices implemented based on this information. Always read and follow pesticide labels. The label is the law. Product registration may vary by state.

² FRAC/IRAC code represents the mode of action of the fungicide/insecticide.

³ PHI refers to the pre-harvest interval, which is the number of days before harvest that the product may not be applied.

⁴ All fungicides/insecticides have a Restricted-Entry Interval (REI). The restricted-entry interval is the time immediately after a pesticide application when entry into the treated area is limited. Check labels for REI. Restrictions in REI may prohibit the use of certain pesticides during harvest.

Applicators must abide by both maximum amount of product per season AND maximum number of applications.

⁵ Max amt refers to the product's maximum amount/acre/year.

⁶ Max app refers to the product's maximum number of applications per year.

RUP refers to restricted use pesticide.

Pear Dormant to Bud Swell - Diseases

Apply before growth starts in the spring and when temperatures are above 45°F.

Disease management notes

- **Fire blight:** If fire blight occurred last year, a fixed copper spray (copper hydroxide, copper oxychloride, basic copper sulfate, Bordeaux mixture) at swollen bud stage is suggested. Label recommendations may vary; be sure to review product labels.

Table 2-1. Pear Diseases - Dormant Through Bud Swell¹

| Product And Formulation Active Ingredient | FRAC Code ² | Fire Blight | REI ³ PHI ⁴ | Max Amt ⁵ Max App ⁶ |
|--|---------------------------|----------------|--------------------------------------|--|
| Badge | M | 5,25-14 pt. | 48 h | 56.3 pt. |
| copper oxychloride | | F-G | 0d | NA |
| Bordeaux | M | 8 lb. | 24 h | NA |
| limed copper sulfate | | F-G | NA | NA |
| C-O-C-S | M | 12-15.6 lb. | 48 h | 31 lb. |
| copper oxychloride, sulfate | | F-G | NA | 1 every 5d |
| Cuprofix Ultra 40 Disperss | M | 7.5-10 lb. | 12h | NA |
| copper sulfate | | F-G | NA | NA |
| Kocide 3000 | M | 7 lb. | 48h | 53.3 lb. |
| copper hydroxide | | F-G | 0d | NA |

Footnotes: See how to read the spray schedule tables section at the beginning of the chapter.

Pear Dormant to Bud Break - Insects

Apply before buds break into green tip.

Insect management notes

Pear psylla

- Adult pear psylla become active when temperature is above 40°F. First eggs of pear psylla are laid on buds and twigs when temperature is above 50°F. Peak psylla egg laying is at the time of bud burst.
- In addition to applying oil, apply an adulticide once psylla adults are seen. Pyrethroids (Asana, Danitol, Mustang Maxx, Permethrin, Pounce) are most effective but should be avoided if resistance is a

concern. Pear psylla populations at some locations are resistant to pyrethroids. Pyrethroids work best when temperatures are cool. Another option is Surround. Surround should be applied every 7-14 days beginning no later than green tip.

- Dimilin is an insect growth regulator that should be applied during egg deposition.

Oil

- Apply when temperatures are above 40°F, never during freezing weather. Do not apply within two weeks of a sulfur spray or later than delayed dormant. Insecticide may be combined with oil during dormant and delayed-dormant periods only. Oil on wood inhibits psylla egg laying/hatching. Apply oil as soon as psylla first eggs are laid and again seven days later if adults are still present.

Sulfur and lime-sulfur

- Should not be applied once green tissue is present on sensitive cultivars (Anjou, Comice or Seckle)
- Lime-sulfur can be applied at the dormant stage, using a high rate (11 gal./acre). A lower rate (3 gal./acre) can be used if not applied until the delayed dormant stage. In addition to controlling pear psylla and San Jose scale, lime sulfur controls pear rust mite and pearleaf blister mite. Elemental sulfur used during the delayed dormant period to control pearleaf blister mite also controls pear psylla. Do not use sulfur or lime-sulfur when temperatures are predicted to exceed 90°F during or within three days of application. Sulfur sprays are most effective when the temperature is above 60°F after application. Not all elemental sulfur products are registered in all Midwest states.

Table 2-2. Pear Insects - Dormant Through Bud Break¹

| Product And Formulation Active Ingredient | IRAC Code ² | Pear Psylla EGG/ADULT | San Jose Scale | European Red Mite | REI ³ PHI ⁴ | Max Amt ⁵ Max App ⁶ |
|--|---------------------------|---------------------------|-----------------------|-----------------------|--------------------------------------|--|
| Acramite 50WS | 20D | x | x | 0.7-1 lb. | 12h | NA |
| bifenazate | | x | x | G | 7d | 1 |
| Agri-Mek SC (0.7SC) (RUP) | 6 | x | x | 2.2-4.2 fl. oz. | 12h | 8.5 fl. oz. |
| abamectin | | x | x | E | 28d | 2 |
| Apollo SC (ISC) | 10A | x | x | 4-8 fl. oz. | 12h | NA |
| clofentezine | | x | x | E | 21d | 1 |
| Asana XL (0.66EC) (RUP) | 3A | 4.8-19.2 fl. oz. | x | x | 12h | 72 fl. oz. |
| esfenvalerate | | G[r] | x | x | 28d | NA |
| Brigade eVo (RUP) | 3A | x | x | 5.1-12.8 fl. oz. | 12h | 32 oz. |
| bifenthrin | | x | x | F | 14d | 3 |
| Centaur WDG (70WDG) | 16 | x | 34.5-46 oz. | x | 12h | 69 oz. |
| buprofezin | | x | E | x | 14d | 2 |
| Damoil | UN | 1-4% | 0.5-4% | see label | 4h | NA |
| mineral oil | | G | G | G | 0d | NA |
| Danitol 2.4EC (RUP) | 3A | 16-21.3 fl. oz. | x | x | 24h | 42.7 fl. oz. |
| fenpropathrin | | G[r] | x | x | 14d | NA |
| Delta Gold (1.5EC) (RUP) | 3A | 1.9 fl. oz. | x | x | 12h | 3.6 fl. oz. |
| deltamethrin | | s | x | x | 21d | NA |
| Diazinon AG 600WBC (RUP) | 1B | 6.5-12.7 fl. oz./100 gal. | 12.7 fl. oz./100 gal. | 12.7 fl. oz./100 gal. | 4d | 102 fl. oz. |
| diazinon | | F | G | u | 21d | 2 |
| Dimethoate (4EC) | 1B | 0.5-1 pt./100 gal. | x | 0.5-1 pt./100 gal. | 10d | 2 pt. |
| dimethoate | | u | x | u | 28d | NA |
| Dimilin 2L (2AF) (RUP) | 15 | 12-48 fl. oz. | x | x | 12h | 64 fl. oz. |
| diflubenzuron | | E | x | x | 14d | 4 |
| Envidor 2SC | 23 | x | x | 16-18 fl. oz. | 12h | 18 fl. oz. |
| spirodiclofen | | x | x | E | 7d | 1 |
| Esteem 35WP | 7C | x | 4-5 oz. | x | 12h | 10 oz. |
| pyriproxifen | | x | E | x | 45d | 2 |
| Grandevo | UN | x | x | 2-3 lb. | 4h | NA |
| <i>Chromobacterium subtsugae</i> | | x | x | u | 0d | NA |
| Kanemite 15SC | 20B | x | x | 21-31 fl. oz. | 12h | 62 fl. oz. |
| acequinocyl | | x | x | E | 14d | 2 |
| Lime-sulfur solution | UN | 3-5 gal. | 3-11 gal. | 3-11 gal. | 48h | NA |
| calcium polysulfide | | E | u | u | PB/PH* | NA |
| Mustang Maxx (0.8EC) (RUP) | 3A | 1.2-4 fl. oz. | x | x | 12h | 24 fl. oz. |
| zeta-cypermethrin | | G[r] | x | x | 14d | NA |
| Nealta (1.67SC) | 25 | x | x | 13.7 fl. oz. | 12h | 274 fl. oz. |
| cyflumetofen | | x | x | E | 7d | 2 |
| Nexter SC (3.75SC) | 21A | x | x | 11-17 fl. oz. | 12h | NA |
| pyridaben | | x | x | E | 7d | 1 |

(Continued)

Table 2-2. Pear Insects - Dormant Through Bud Break¹ (continued)

| Product And Formulation Active Ingredient | IRAC Code ² | Pear Psylla EGG/ADULT | San Jose Scale | European Red Mite | REI ³ PHI ⁴ | Max Amt ⁵ Max App ⁶ |
|--|---------------------------|--------------------------|-------------------|----------------------|--------------------------------------|--|
| Onager Optek (1EC) | 10 | x | x | 12-24 fl. oz. | 12h | 24 fl. oz. |
| hexythiazox | | x | x | E | 28d | 1 |
| Permethrin 3.2EC (RUP) | 3A | 8-16 fl. oz. | x | x | 12h | 26 fl. oz. |
| permethrin | | G[r] | x | x | PB | NA |
| Portal XLO (0.4EC) | 21A | x | x | 2 pt. | 12h | 2 pt. |
| fenpyroximate | | x | x | E | 14d | 1 |
| Pounce 25WP (RUP) | 3A | 12.8-25.6 oz. | x | x | 12h | 41.6 oz. |
| permethrin | | G[r] | x | x | PB* | NA |
| Sivanto Prime (1.67SC) | 4D | x | 10.5-14 fl. oz. | x | 12h | 28 fl. oz. |
| flupyradifurone | | x | s | x | 14d | NA |
| Soap (M-Pede, Des-X, etc.) | UN | 2% | 2% | 2% | 12h | NA |
| potassium salts of fatty acids | | F | F | u | 0d | NA |
| Sulfur (Microfine; 90%) | UN | x | x | 10-60 lb. | 24h | NA |
| sulfur | | x | x | u | 0d | NA |
| Surround WP (95WP) | UN | 50 lb. | x | x | 4h | NA |
| kaolin | | G | x | x | 0d | NA |
| Vendex 50WP (RUP) | 12B | x | x | 1-2 lb. | 48h | 4 lb. |
| fenbutatin-oxide | | x | x | G | 14d | 2 |
| Vydate L (2L) (RUP) | 1A | x | x | 6-8 pt. | 48h | 8 pt. |
| oxamyl | | x | x | G | 14d | 1 |
| Warrior II (2.08CS) (RUP) | 3A | 1.2-2.5 fl. oz. | x | x | 24h | 12.8 fl. oz. |
| lambda-cyhalothrin | | s | x | x | 21d | NA |
| Zeal (72WP) | 10B | x | x | 2-3 oz. | 12h | 3 oz. |
| etoxazole | | x | x | E | 14d | 1 |

*PB = prebloom. PH = postharvest.

Footnotes: See how to read the spray schedule tables section at the beginning of the chapter.

Pear Bud Break To Bloom - Diseases

Apply every 7-10 days after bud break to protect new growth from green tip to bloom.

Disease management notes

- Syllit FL should be mixed with Mancozeb 75DF. Begin applications at 1/4- to 1/2-inch green tip and continue on a 7- to 10-day schedule to bloom.

Table 2-3. Pear Diseases - Bud Break Through Bloom¹

| Product And Formulation Active Ingredient | FRAC Code ² | Fabraea Leaf Spot | Powdery Mildew | Scab | REI ³ PHI ⁴ | Max Amt ⁵ Max App ⁶ |
|--|---------------------------|----------------------|-------------------|---------------|--------------------------------------|--|
| Aprovia (EC) | 7 | x | 5.5-7 fl. oz. | 5.5-7 fl. oz. | 12h | 27.6 fl. oz. |
| benzovindiflupyr | | x | F | G-E | 30d | NA |
| Captan 80 WDG | M | x | 2.5-5 lb. | 5 lb. | 24h | 40 lb. |
| captan | | x | i | G | 0d | NA |
| Cevya | 3 | x | 3-5 fl. oz. | 3-5 fl. oz. | 12h | NA |
| mefentrifluconazole | | x | G-E | E | 0d | NA |

(Continued)

Table 2-3. Pear Diseases - Bud Break Through Bloom¹ (continued)

| Product And Formulation Active Ingredient | FRAC Code ² | Fabraea Leaf Spot | Powdery Mildew | Scab | REI ³ PHI ⁴ | Max Amt ⁵ Max App ⁶ |
|--|---------------------------|----------------------|-------------------|-----------------|--------------------------------------|--|
| C-O-C-S | M | x | x | 0.5-1 lb. | 48 h | 31 lb. |
| copper oxychloride, sulfate | | x | x | u | NA | NA |
| Cuprofix Ultra 40 disperss | M | x | x | 1-2.5 lb. | 12h | NA |
| copper sulfate | | x | x | F-G | NA | NA |
| Ferbam Granulfo (76WDG) | M | 3.5 lb. | x | 3.5 lb. | 24h | NA |
| ferbam | | E | x | F | NA | 3 |
| Flint Extra | 11 | x | 2.5-2.9 fl. oz. | 2.5-2.9 fl. oz. | 12h | 10.5 fl. oz. |
| trifloxystrobin | | x | G[r] | E[r] | 14d | NA |
| Fontelis (SC) | 7 | x | 16-20 fl. oz. | 16-20 fl. oz. | 12h | 61 fl. oz. |
| penthiopyrad | | x | G | E | 28d | NA |
| Inspire Super (EW) | 3+9 | x | 12 fl. oz. | 12 fl. oz. | 12h | 60 fl. oz. |
| difenoconazole + cyprodinil | | x | F | E | 28d | NA |
| Kenja 400SC | 7 | x | 12.5 fl. oz. | 12.5 fl. oz. | 12h | NA |
| isofetamid | | x | s | F | 20d | NA |
| Kocide 3000 | M | x | x | 0.75 lb. | 48h | 53.3 lb. |
| copper hydroxide | | x | x | F-G | 0d | NA |
| Luna Privilege | 7 | x | 2.4-6.8 fl. oz. | 4-6.8 fl. oz. | NA | NA |
| fluopyram | | x | G | G-E | NA | NA |
| Luna Sensation (SC) | 7+11 | x | 5-5.8 fl. oz. | 4-5.8 fl. oz. | 12h | 21 fl. oz. |
| fluopyram + trifloxystrobin | | x | G[r] | E[r] | 14d | 4 |
| Luna Tranquility (SC) | 7+9 | x | 11.2-16 fl. oz. | 11.2-16 fl. oz. | 12h | 54.7 fl. oz. |
| fluopyram + pyrimethanil | | x | G | E | 72d | NA |
| Merivon (2.09SC) | 7+11 | x | 4-5.5 fl. oz. | 4-5.5 fl. oz. | 12h | 22 fl. oz. |
| fluxapyroxad + pyraclostrobin | | x | G | E | 0d | 4 |
| Microthiol Disperss | M | x | 10-20 lb. | 10-20 lb. | 24h | NA |
| sulfur | | x | G | i-F | 0d | NA |
| Miravis (1.67 SC) | 7 | x | 3.4 fl. oz. | 3.4 fl. oz. | 4h | 13.6 fl. oz. |
| pydiflumetofen | | x | G | E | 30d | 4 |
| OSO 5% SC | 19 | x | 6.5-13 fl. oz. | 6.5-13 fl. oz. | 4h | 78 fl. oz. |
| polyoxin D | | x | F | F | 0d | 6 |
| Pristine | 11+7 | x | 14.5-18.5 oz. | 14.5-18.5 oz. | 12h | 74 oz. |
| pyraclostrobin + boscalid | | x | E[r] | E[r] | 0d | 4 |
| Procure 480SC | 3 | x | 8-16 fl. oz. | 8-16 fl. oz. | 12h | 64 fl. oz. |
| triflumizole | | x | E[r] | G[r] | 14d | NA |
| Roper DF Rainshield | M | 3 or 6 lb. | 6 lb. | 6 lb. | 24h | 21 lb. |
| mancozeb | | E | i | G | 77d | 6 |
| Scala SC | 9 | x | x | 7-10 fl. oz. | 12h | 40 fl. oz. |
| pyrimethanil | | x | x | G-E | 72d | NA |
| Sovran (50WG) | 11 | x | 4-6.4 oz. | 3.2-6.4 oz. | 12h | 25.6 oz. |
| kresoxim-methyl | | x | G[r] | E[r] | 30d | 4 |

(Continued)

Table 2-3. Pear Diseases - Bud Break Through Bloom¹ (continued)

| Product And Formulation Active Ingredient | FRAC Code ² | Fabraea Leaf Spot | Powdery Mildew | Scab | REI ³ PHI ⁴ | Max Amt ⁵ Max App ⁶ |
|--|---------------------------|----------------------|-------------------|------------|--------------------------------------|--|
| Syllit FL | U12 | 3 pt. | x | 3 pt. | 48h | 9 pt. |
| dodine | | G | x | E[r] | 7d | 3 |
| Topguard Specialty Crops | 3 | x | 8-12 fl. oz. | 13 fl. oz. | 12h | 52 fl. oz. |
| flutriafol | | x | E | E-G | 14d | 4 |
| Topsin-M WSB | 1 | 1 lb. | 0.75-1 lb. | 0.75-1 lb. | 24h | 4 lb. |
| thiophanate methyl | | E[r] | G[r] | i | 1d | NA |
| Torino (SC) | U6 | x | 6.8 oz. | x | 4h | 6.8 oz. |
| cyflufenamid | | x | E | x | 14d | 1 |
| Vanguard WG | 9 | x | x | 5 oz. | 12h | 30 oz. |
| cyprodinil | | x | x | G | 0d | 2 |
| Ziram 76DF | M3 | 6 | x | 6 lb. | 48h | 42.4 lb. |
| ziram | | E | x | G | 14d | 7 |

Footnotes: See how to read the spray schedule tables section at the beginning of the chapter.

Green Cluster To White Bud - Insects

Insect management notes

Pear psylla

- At green cluster (tight cluster), eggs of pear psylla begin to hatch and young nymphs are found. Apply insecticide when eggs begin to hatch. If pyrethroids (Asana, Danitol, Mustang Maxx, Permethrin, Pounce) were used earlier for adult control, switch to different chemistry (e.g., Dimilin or Centaur, or Esteem, Nexter or Surround).
- Do not use sulfur or lime-sulfur within two weeks of an oil spray. Sulfur or lime-sulfur should not be applied once green tissue is present on sensitive cultivars (Anjou, Comice or Seckle). Sulfur sprays are most effective when the temperature is above 60°F after application. Do not use sulfur or lime-sulfur when temperatures are predicted to exceed 90°F during or within 3 days of application. Not all elemental sulfur or lime sulfur products are registered in all Midwest states.
- Do not apply Actara after green cluster until petal fall.

Table 2-4. Pear Insects - Green Cluster Through White Bud¹

| Product And Formulation Active Ingredient | IRAC Code ² | Pear Psylla NYMPH | San Jose Scale | European Red Mite | REI ³ PHI ⁴ | Max Amt ⁵ Max App ⁶ |
|--|---------------------------|----------------------|-------------------|----------------------|--------------------------------------|--|
| Acramite 50WS | 20D | x | x | 0.7-1 lb. | 12h | NA |
| bifenazate | | x | x | G | 7d | 1 |
| Actara (25WDG) | 4A | 5.5 oz. | x | x | 12h | 16.5 oz. |
| thiamethoxam | | E | x | x | 35d | NA |
| Admire Pro (4.6F) | 4A | 7 fl. oz. | x | x | 12h | 14 fl. oz. |
| imidacloprid | | G | x | x | 7d | NA |
| Agri-Mek SC (0.7SC) (RUP) | 6 | 2.2-4.2 fl. oz. | x | 2.2-4.2 fl. oz. | 12h | 8.5 fl. oz. |
| abamectin | | G | x | E | 28d | 2 |
| Apollo SC (1SC) | 10A | x | x | 4-8 fl. oz. | 12h | NA |
| clofentezine | | x | x | E | 21d | 1 |
| Apta (1.34SC) | 21A | 21-27 fl. oz. | x | x | 12h | 53.5 fl. oz. |
| tolfenpyrad | | G | x | x | 14d | 2 |

(Continued)

Table 2-4. Pear Insects - Green Cluster Through White Bud¹ (continued)

| Product And Formulation Active Ingredient | IRAC Code ² | Pear Psylla NYMPH | San Jose Scale | European Red Mite | REI ³ PHI ⁴ | Max Amt ⁵ Max App ⁶ |
|--|---------------------------|----------------------|------------------------|------------------------|--------------------------------------|--|
| Assail 30SG | 4A | 4-8 oz. | x | x | 12h | 32 oz. |
| acetamiprid | | G | x | x | 7d | 4 |
| Belay (2.13SC) | 4A | 6 fl. oz. | x | x | 12h | 12 fl. oz. |
| clothianidin | | G | x | x | 7d | NA |
| Brigade eVo (RUP) | 3A | x | x | 5.1-12.8 fl. oz. | 12h | 32 fl. oz. |
| bifenthrin | | x | x | F | 14d | NA |
| Centaur WDG (70WDG) | 16 | 34.5-46 oz. | 34.5-46 oz. | x | 12h | 69 oz. |
| buprofezin | | u | E | x | 14d | 2 |
| Damoil | UN | 1-4% | 0.5-4% | See Label | 4h | NA |
| Mineral Oil | | | G | G | 0d | NA |
| Delegate WG (25WG) | 5 | 6-7 oz. | x | x | 4h | 28 oz. |
| spinetoram | | E | x | x | 7d | 4 |
| Diazinon AG 600WBC (RUP) | 1B | x | 12.7 fl. oz./ 100 gal. | 12.7 fl. oz./ 100 gal. | 4d | 102 fl. oz. |
| diazinon | | x | G | u | 21d | 2 |
| Dimethoate (4EC) | 1B | x | x | 0.5-1 pt./100 gal. | 10d | 2 pt. |
| dimethoate | | x | x | u | 28d | NA |
| Dimilin 2L (2AF) (RUP) | 15 | 12-48 fl. oz. | x | x | 12h | 64 fl. oz. |
| diflubenzuron | | E | x | x | 14d | 4 |
| Envidor 2SC | 23 | x | x | 16-18 fl. oz. | 12h | 18 fl. oz. |
| spiroticlofen | | x | x | E | 7d | 1 |
| Esteem 35WP | 7C | 5 oz. | 4-5 oz. | x | 12h | 10 oz. |
| pyriproxifen | | G | E | x | 45d | 2 |
| Grandevo | UN | 2-3 lb. | x | 2-3 lb. | 4h | NA |
| <i>Chromobacterium subtsugae</i> | | u | x | u | 0d | NA |
| Kanemite 15SC | 20B | x | x | 21-31 fl. oz. | 12h | 62 fl. oz. |
| acequinocyl | | x | x | E | 14d | 2 |
| Lime-sulfur solution | UN | x | 3-11 gal. | 3-11 gal. | 48h | NA |
| calcium polysulfide | | x | u | u | PB/PH* | NA |
| Nealta (1.67SC) | 25 | x | x | 13.7 fl. oz. | 12h | 274 fl. oz. |
| cyflumetofen | | x | x | E | 7d | 2 |
| Neemix 4.5 (0.39L) | UN | 7-16 fl. oz. | x | x | 4h | NA |
| azadirachtin | | F | x | x | 0d | NA |
| Nexter SC (3.75SC) | 21A | 11-17 fl. oz. | x | 11-17 fl. oz. | 12h | NA |
| pyridaben | | G | x | E | 7d | 1 |
| Onager Optek (1EC) | 10 | x | x | 12-24 fl. oz. | 12h | 24 fl. oz. |
| hexythiazox | | x | x | E | 28d | 1 |
| Portal XLO (0.4EC) | 21A | 2 pt. | x | 2 pt. | 12h | 2 pt. |
| fenpyroximate | | G | x | E | 14d | 1 |
| Proclaim (5SG) (RUP) | 6 | 3.2-4.8 oz. | x | 3.2-4.8 oz. | 12 or 48h | 14.4 oz. |
| emamectin benzoate | | s | x | s | 14d | NA |

(Continued)

Table 2-4. Pear Insects - Green Cluster Through White Bud¹ (continued)

| Product And Formulation Active Ingredient | IRAC Code ² | Pear Psylla NYMPH | San Jose Scale | European Red Mite | REI ³ PHI ⁴ | Max Amt ⁵ Max App ⁶ |
|--|---------------------------|----------------------|-------------------|----------------------|--------------------------------------|--|
| Sevin XLR Plus (4F) | 1A | 1.5-3 qt. | x | x | 12h | 15 qt. |
| carbaryl | | u | x | x | 3d | 8 |
| Sivanto Prime (1.67SC) | 4D | 10.5-14 fl. oz. | 10.5-14 fl. oz. | x | 12h | 28 fl. oz. |
| flupyradifurone | | G | s | x | 14d | NA |
| Soap (M-Pede, Des-X, etc.) | UN | 2% | 2% | 2% | 12h | NA |
| potassium salts of fatty acids | | F | F | u | 0d | NA |
| Sulfur (Microfine; 90%) | UN | x | x | 10-60 lb. | 24h | NA |
| sulfur | | x | x | u | 0d | NA |
| Surround WP (95WP) | UN | 50 lb. | x | x | 4h | NA |
| kaolin | | G | x | x | 0d | NA |
| Vendex 50WP (RUP) | 12B | x | x | 1-2 lb. | 48h | 4 lb. |
| fenbutatin-oxide | | x | x | G | 14d | 2 |
| Verdepryn 100SL (0.83SL) | 28 | 11 fl. oz. | x | x | 4h | 33 fl. oz. |
| cyclaniliprole | | u | x | x | 7d | 3 |
| Vydate L (2L) (RUP) | 1A | x | x | 6-8 pt. | 48h | 8 pt. |
| oxamyl | | x | x | G | 14d | 1 |
| Zeal (72WP) | 10B | x | x | 2-3 oz. | 12h | 3 oz. |
| etoxazole | | x | x | E | 14d | 1 |

*PB = prebloom. PH = postharvest.

Footnotes: See how to read the spray schedule tables section at the beginning of the chapter.

Pear Bloom To Petal Fall - Diseases

Disease management notes

Fire blight

- Start fire blight sprays at first sign of open blossoms. Repeat sprays at 4- to 5-day intervals through bloom and petal fall on highly susceptible varieties. A minimum of two applications are necessary to provide control. Better control has been obtained

by adding 1 pt of Regulaid to Streptomycin 17W (1 lb.).

- In orchards with documented streptomycin resistant fire blight use Mycoshield 17WP (16 oz./200 ppm) and generic oxytetracycline products alternated with Kasumin 2L (64 oz. per 100 gal.). Do not exceed two sequential treatments per year for use to manage streptomycin-resistant fire blight bacteria.

Table 2-5. Pear Diseases - Bloom Through Petal Fall¹

| Product And Formulation Active Ingredient | FRAC Code ² | Fabraea Leaf Spot | Fire Blight | Powdery Mildew | Scab | Summer Rot | REI ³ PHI ⁴ | Max Amt ⁵ Max App ⁶ |
|--|---------------------------|----------------------|-------------|-------------------|---------------|---------------|--------------------------------------|--|
| Agri-Mycin 17 | 25 | x | 24-48 oz. | x | x | x | 12h | NA |
| streptomycin sulfate | | x | G-E[r] | x | x | x | 50d | NA |
| Aprovia (EC) | 7 | x | x | 5.5-7 fl. oz. | 5.5-7 fl. oz. | 5.5-7 fl. oz. | 12h | 27.6 fl. oz. |
| benzovindiflupyr | | x | x | F | G-E | G-E | 30d | NA |
| Badge | M | x | 0.5-1 pt. | x | x | x | 48h | 56.3 pt. |
| copper oxychloride | | x | F-G | x | x | x | 0d | NA |

(Continued)

Table 2-5. Pear Diseases - Bloom Through Petal Fall¹ (continued)

| Product And Formulation Active Ingredient | FRAC Code ² | Fabraea Leaf Spot | Fire Blight | Powdery Mildew | Scab | Summer Rot | REI ³ PHI ⁴ | Max Amt ⁵ Max App ⁶ |
|--|---------------------------|----------------------|-------------|-------------------|-----------------|---------------|--------------------------------------|--|
| Captan 80 WDG | M | x | x | 2.5-5 lb. | 5 lb. | 2.5-5 lb | 24h | 40 lb. |
| captan | | x | x | i | G | G | 0d | NA |
| Cevya | 3 | x | x | 3-5 fl. oz. | 3-5 fl. oz. | 3-5 fl. oz. | 12h | NA |
| mefentrifluconazole | | x | x | G-E | E | x | 0d | NA |
| C-O-C-S | M | x | 0.5-1 lb. | x | 0.5-1 lb. | x | 48h | 31 lb. |
| copper oxychloride, sulfate | | x | F-G | x | u | x | NA | NA |
| Cuprofix Ultra 40 disperss | M | x | .75 | x | 1-2.5 lb. | x | 12h | NA |
| copper sulfate | | x | F-G | x | F-G | x | NA | NA |
| Ferbam Granulfo (76WDG) | M | 3.5 lb. | x | x | 3.5 lb. | 3.5 lb. | 24h | NA |
| ferbam | | E | x | x | F | G-E | NA | 3 |
| Flint Extra | 11 | x | x | 2.5-2.9 fl. oz. | 2.5-2.9 fl. oz. | 2.9 fl. oz. | 12h | 10.5 fl. oz. |
| trifloxystrobin | | x | x | G[r] | E[r] | x | 14d | NA |
| Fontelis (SC) | 7 | x | x | 16-20 fl. oz. | 16-20 fl. oz. | 16-20 fl. oz. | 12h | 61 fl. oz. |
| penthiopyrad | | x | x | G | E | x | 28d | NA |
| Inspire Super (EW) | 3+9 | x | x | 12 fl. oz. | 12 fl. oz. | x | 12h | 60 fl. oz. |
| difenoconazole + cyprodinil | | x | x | F | E | x | 28d | NA |
| Kasumin 2L | 24 | x | 64 fl. oz. | x | x | x | 12h | 256 fl. oz. |
| kasugamycin | | x | G | x | x | x | 90d | 4 |
| Kenja 400SC | 7 | x | x | 12.5 fl. oz. | 12.5 fl. oz. | x | 12h | NA |
| isofetamid | | x | x | s | F | x | 20d | NA |
| Kocide 3000 | M | x | 0.75 lb. | x | 0.75 lb. | x | 48h | 53.3 lb. |
| copper hydroxide | | x | F-G | x | F-G | x | 0d | NA |
| Luna Privilege | 7 | x | x | 2.4-6.8 fl. oz. | 4-6.8 fl. oz. | x | NA | NA |
| fluopyram | | x | x | G | G-E | x | NA | NA |
| Luna Sensation (SC) | 7+11 | x | x | 5-5.8 fl. oz. | 4-5.8 fl. oz. | 4-5.8 fl. oz. | 12h | 21 fl. oz. |
| fluopyram + trifloxystrobin | | x | x | G[r] | E[r] | x | 14d | 4 |
| Luna Tranquility (SC) | 7+9 | x | x | 11.2-16 fl. oz. | 11.2-16 fl. oz. | x | 12h | 54.7 fl. oz. |
| fluopyram + pyrimethanil | | x | x | G | E | x | 72d | NA |
| Merivon (2.09SC) | 7+11 | x | x | 4-5.5 fl. oz. | 4-5.5 fl. oz. | 4-5.5 fl. oz. | 12h | 22 fl. oz. |
| fluxapyroxad + pyraclostrobin | | x | x | G | E | E | 0d | 4 |
| Microthiol Disperss | M | x | x | 10-20 lb. | 10-20 lb. | x | 24h | NA |
| sulfur | | x | x | G | i-F | x | 0d | NA |
| Miravis (1.67 SC) | 7 | x | x | 3.4 fl. oz. | 3.4 fl. oz. | x | 4h | 13.6 fl. oz. |
| pydiflumetofen | | x | x | G | E | x | 30d | 4 |
| Mycoshield | 41 | x | 1 lb. | x | x | x | 12h | 9 lb. |
| oxytetracycline | | x | G | x | x | x | 60d | 6 |
| OSO 5% SC | 19 | x | x | 6.5-13 fl. oz. | 6.5-13 fl. oz. | 6.5 fl. oz. | 4h | 78 fl. oz. |
| polyoxin D | | x | x | F | F | x | 0d | 6 |

(Continued)

Table 2-5. Pear Diseases - Bloom Through Petal Fall¹ (continued)

| Product And Formulation Active Ingredient | FRAC Code ² | Fabraea Leaf Spot | Fire Blight | Powdery Mildew | Scab | Summer Rot | REI ³ PHI ⁴ | Max Amt ⁵ Max App ⁶ |
|--|---------------------------|----------------------|-------------|-------------------|---------------|---------------|--------------------------------------|--|
| Polyram 80 DF | M3 | x | x | x | 3 lb. | x | 24h | 21 lb. |
| metiran | | x | x | x | G | x | 77d | 7 |
| Pristine | 11+7 | x | x | 14.5-18.5 oz. | 14.5-18.5 oz. | 14.5-18.5 oz. | 12h | 74 oz. |
| pyraclostrobin + boscalid | | x | x | E[r] | E[r] | E-G[r] | 0d | 4 |
| Procure 480SC | 3 | x | x | 8-16 fl. oz. | 8-16 fl. oz. | x | 12h | 64 fl. oz. |
| triflumizole | | x | x | E[r] | G[r] | x | 14d | NA |
| Roper DF Rainshield | M | x | x | 6 lb. | 6 lb. | 3 lb. | 24h | 21 lb. |
| mancozeb | | x | x | i | G | G-E | 77d | 6 |
| Scala SC | 9 | x | x | x | 7-10 fl. oz. | x | 12h | 40 fl. oz. |
| pyrimethanil | | x | x | x | E-G | x | 72d | NA |
| Sovran (50WG) | 11 | x | x | 4-6.4 oz. | 3.2-6.4 oz. | 4-6.4 oz. | 12h | 25.6 oz. |
| kresoxim-methyl | | x | x | G[r] | E[r] | x | 30d | 4 |
| Topguard Specialty Crops | 3 | x | x | 8-12 fl. oz. | 13 fl. oz. | 13 fl. oz. | 12h | 52 fl. oz. |
| flutriafol | | x | x | E | G-E | F-G[r] | 14d | 4 |
| Topsin-M WSB | 1 | 1 lb. | x | 0.75-1 lb. | 0.75-1 lb. | 0.75-1 lb. | 24h | 4 lb. |
| thiophanate methyl | | E[r] | x | G[r] | i | E-G[r] | 1d | NA |
| Torino (SC) | U6 | x | x | 6.8 oz. | x | x | 4h | 6.8 oz. |
| cyflufenamid | | x | x | E | x | x | 14d | 1 |
| Vanguard WG | 9 | x | x | x | 5 oz. | x | 12h | 30 oz. |
| cyprodinil | | x | x | x | G | x | 0d | 2 |
| Ziram 76DF | M3 | 6 lb. | x | x | 6 lb. | 6 lb. | 48h | 42.4 lb. |
| ziram | | E | x | x | G | E | 14d | 7 |

Footnotes: See how to read the spray schedule tables section at the beginning of the chapter.

Pear Petal Fall - Insects

7-10 days after bloom.

Insect management notes

- **Pear psylla:** The pear psylla population should be low at this time if insecticides were applied at earlier stages. Petal fall is peak egg hatch. Nymphs are the target of spray at petal fall. Best options are

Agri-Mek, Delegate and Actara. Agri-Mek is effective now through second cover because leaves are tender; it is not as effective once leaves harden off.

- **Codling moth:** If pheromone mating disruption is used, dispensers should be set up at petal fall. Recommended only in orchards that are 5 acres or larger. See list of products in Apple chapter.

Table 2-6. Pear Insects - Petal Fall¹

| Product And Formulation Active Ingredient | IRAC Code ² | Pear Psylla NYMPH | Plum Curculio | Stink Bug | Tarnished Plant Bug | REI ³ PHI ⁴ | Max Amt ⁵ Max App ⁶ |
|--|---------------------------|----------------------|------------------|--------------|------------------------|--------------------------------------|--|
| Actara (25WDG) | 4A | 5.5 oz. | 4.5-5.5 oz. | x | x | 12h | 16.5 oz. |
| thiamethoxam | | E | G | x | x | 35d | NA |
| Admire Pro (4.6F) | 4A | 7 fl. oz. | x | x | x | 12h | 14 fl. oz. |
| imidacloprid | | G | x | x | x | 7d | NA |

(Continued)

Table 2-6. Pear Insects - Petal Fall¹ (continued)

| Product And Formulation Active Ingredient | IRAC Code ² | Pear Psylla NYMPH | Plum Curculio | Stink Bug | Tarnished Plant Bug | REI ³ PHI ⁴ | Max Amt ⁵ Max App ⁶ |
|--|---------------------------|----------------------|----------------------|---------------------|------------------------|--------------------------------------|--|
| Agri-Mek SC (0.7SC) (RUP) | 6 | 2.2-4.2 fl. oz. | x | x | x | 12h | 8.5 fl. oz. |
| abamectin | | G | x | x | x | 28d | 2 |
| Altacor eVo | 28 | x | 1.3-2.2 oz. | x | x | 4h | 9 fl. oz. |
| chlorantraniliprole | | x | s | x | x | 5d | NA |
| Apta (1.34SC) | 21A | 21-27 fl. oz. | 21-27 fl. oz. | x | x | 12h | 53.5 oz. |
| tolfenpyrad | | G | G | x | x | 14d | 2 |
| Asana XL (0.66EC) (RUP) | 3A | x | 4.8-14.5 fl. oz. | x | x | 12h | 72 fl. oz. |
| esfenvalerate | | x | F | x | x | 28d | NA |
| Assail 30SG | 4A | 4-8 oz. | 8 oz. | x | x | 12h | 32 oz. |
| acetamiprid | | G | G | x | x | 7d | 4 |
| Baythroid XL (1EC) (RUP) | 3A | x | 2.4-2.8 fl. oz. | 2-2.4 fl. oz. | 2-2.4 fl. oz. | 12h | 2.8 fl. oz. |
| beta-cyfluthrin | | x | F | G | E | 7d | NA |
| Belay (2.13SC) | 4A | 6 fl. oz. | 6 fl. oz. | 6 fl. oz. | 4-6 fl. oz. | 12h | 12 fl. oz. |
| clothianidin | | G | G | E | G | 7d | NA |
| Beleaf 50SG | 29 | x | x | x | 2-2.8 oz. | 12h | 8.4 oz. |
| flonicamid | | x | x | x | G | 21d | 3 |
| Brigade eVo (RUP) | 3A | x | 2.6-12.8 fl. oz. | 2.6-12.8 fl. oz. | 2.6-12.8 fl. oz. | 12h | 32 fl. oz. |
| bifenthrin | | x | F | E | E | 14d | 3 |
| Centaur WDG (70WDG) | 16 | 34.5-46 oz. | x | x | x | 12h | 69 oz. |
| buprofezin | | u | x | x | x | 14d | 2 |
| Danitol 2.4EC (RUP) | 3A | x | x | 16-21.3 fl. oz. | 16-21.3 fl. oz. | 24h | 42.7 fl. oz. |
| fenpropathrin | | x | x | G | G | 14d | NA |
| Delegate WG (25WG) | 5 | 6-7 oz. | 6-7 oz. | x | x | 4h | 28 oz. |
| spinetoram | | E | s | x | x | 7d | 4 |
| Delta Gold (1.5EC) (RUP) | 3A | x | 0.9-1.9 fl. oz. | 1.9 fl. oz. | 0.9-1.9 fl. oz. | 12h | 3.6 fl. oz. |
| deltamethrin | | x | F | u | E | 21d | NA |
| Dimilin 2L (2AF) (RUP) | 15 | 12-48 fl. oz. | x | x | x | 12h | 64 fl. oz. |
| diflubenzuron | | E | x | x | x | 14d | 4 |
| Esteem 35WP | 7C | 5 oz. | x | x | x | 12h | 10 oz. |
| pyriproxyfen | | G | x | x | x | 45d | 2 |
| Exirel (0.83SE) | 28 | 13.5-20.5 fl. oz. | 13.5-20.5 fl. oz. | x | x | 12h | 61 fl. oz. |
| cyantraniliprole | | s | G | x | x | 3d | 3 |
| Grandevo | UN | 2-3 lb. | x | x | x | 4h | NA |
| <i>Chromobacterium subtsugae</i> | | u | x | x | x | 0d | NA |
| Imidan 70W | 1B | x | 2.1-5.7 lb. | x | x | 7d | 16 lb. |
| phosmet | | x | G | x | x | 7d | NA |
| Magister SC (1.7SC) | 21A | 32-36 fl. oz. | x | x | x | 12h | 36 fl. oz. |
| fenazaquin | | G | x | x | x | 7d | 1 |

(Continued)

Table 2-6. Pear Insects - Petal Fall¹ (continued)

| Product And Formulation Active Ingredient | IRAC Code ² | Pear Psylla NYMPH | Plum Curculio | Stink Bug | Tarnished Plant Bug | REI ³ PHI ⁴ | Max Amt ⁵ Max App ⁶ |
|--|---------------------------|----------------------|------------------|-----------------|------------------------|--------------------------------------|--|
| Movento (2SC) | 23 | 6-9 fl. oz. | x | x | x | 24h | 25 fl. oz. |
| spirotetramat | | G | x | x | x | 7d | NA |
| Mustang Maxx (0.8EC) (RUP) | 3A | x | 1.2-4 fl. oz. | 1.2-4 fl. oz. | 1.2-4 fl. oz. | 12h | 24 fl. oz. |
| zeta-cypermethrin | | x | F | G | E | 14d | NA |
| Neemix 4.5 (0.39L) | UN | 7-16 fl. oz. | x | 7-16 fl. oz. | 7-16 fl. oz. | 4h | NA |
| azadirachtin | | F | x | u | G | 0d | NA |
| Nexter SC (3.75SC) | 21A | 11-17 fl. oz. | x | x | x | 12h | NA |
| pyridaben | | G | x | x | x | 7d | 1 |
| Portal XLO (0.4EC) | 21A | 2 pt. | x | x | x | 12h | 2 pt. |
| fenpyroximate | | G | x | x | x | 14d | 1 |
| Proclaim (5SG) (RUP) | 6 | 3.2-4.8 oz. | x | x | x | 12 or 48h | 14.4 oz. |
| emamectin benzoate | | s | x | x | x | 14d | NA |
| Sevin XLR Plus (4F) | 1A | 1.5-3 qt. | 1.5-3 qt. | x | 1.5-3 qt. | 12h | 15 qt. |
| carbaryl | | u | F | x | i | 3d | 8 |
| Sivanto Prime (1.67SC) | 4D | 10.5-14 fl. oz. | x | x | x | 12h | 28 fl. oz. |
| flupyradifurone | | G | x | x | x | 14d | NA |
| Soap (M-Pede, Des-X, etc.) | UN | 2% | x | x | x | 12h | NA |
| potassium salts of fatty acids | | F | x | x | x | 0d | NA |
| Surround WP (95WP) | UN | 50 lb. | x | 25-50 lb. | 25-50 lb. | 4h | NA |
| kaolin | | G | x | s | s | 0d | NA |
| Transform WG (50WG) | 4C | 2.75 oz. | x | x | 1.5-2.7 oz. | 24h | 8.5 oz. |
| sulfoxaflor | | s | x | x | u | 7d | 4 |
| Verdepryn 100SL (0.83SL) | 28 | 11 fl. oz. | 5.5-11 fl. oz. | 5.5-11 fl. oz. | x | 4h | 33 fl. oz. |
| cyclaniliprole | | u | G | s | x | 7d | 3 |
| Vydate L (2L) (RUP) | 1A | x | x | 1.5-4 pt. | x | 48h | 8 pt. |
| oxamyl | | x | x | G | x | 14d | 1 |
| Warrior II (2.08CS) (RUP) | 3A | x | 1.2-2.5 fl. oz. | 1.2-2.5 fl. oz. | 1.2-2.5 fl. oz. | 24h | 12.8 fl. oz. |
| lambda-cyhalothrin | | x | G | G | G | 21d | NA |

Footnotes: See how to read the spray schedule tables section at the beginning of the chapter.

Pear Summer Cover - Diseases

10-14 days after petal fall, with applications every 10-14 days as needed, continuing until harvest.

Disease management notes

- Where **Febraea leaf spot** is a problem, use Flint Extra, Sovran or Ziram. Observe preharvest interval requirements.
- Use of Ferbam products is not recommended for later cover sprays due to the presence of unsightly residue at harvest.
- The following products have 28- or 30-day PHI. Be careful of their use close to harvest to avoid MRL issues: Aprovia, Fontelis, Inspire Super, Miravis, Omega and Sovran.
- Luna Tranquility has a 72-day PHI.

Table 2-7. Pear Disease - Summer Cover¹

| Product And Formulation Active Ingredient | FRAC code ² | Bitter Rot | Powdery Mildew | Scab | Sooty Blotch/ Flyspeck | Summer Rots (Black And White) | Fabraea Leaf Spot | REI ³ PHI ⁴ | Max Amt ⁵ Max App ⁶ |
|--|---------------------------|------------------|--------------------|--------------------|------------------------------|---|----------------------|--------------------------------------|--|
| Aprovia (EC) | 7 | 5.5-7 fl. oz. | 5.5-7 fl. oz. | 5.5-7 fl. oz. | 5.5-7 fl. oz. | 5.5-7 fl. oz. | x | 12h | 276 fl. oz. |
| benzovindiflupyr | | G-E | F | G-E | G-E | F | x | 30d | NA |
| Captan 80 WDG | M | 2.5-5 lb. | 2.5-5 lb. | 5 lb. | 2.5-5 lb. | 2.5-5 lb. | x | 24h | 40 lb. |
| captan | | E | i | G | G | F-G | x | 0d | NA |
| Cevya | 3 | 3-5 fl. oz. | 3-5 fl. oz. | 3-5 fl. oz. | 3-5 fl. oz. | 3-5 fl. oz. | x | 12h | NA |
| mefentrifluconazole | | G | G-E | E | E | E | x | 0d | NA |
| C-O-C-S | M | x | x | 0.5-1 lb. | x | x | x | 48 h | 31 lb. |
| copper oxychloride, sulfate | | x | x | u | x | x | x | NA | NA |
| Cuprofix Ultra 40 Disperss | M | x | x | 1-2.5 lb. | x | x | x | 12h | NA |
| copper sulfate | | x | x | F-G | x | x | x | NA | NA |
| Ferbam Granulfo (76WDG) | M | 3.5 lb. | x | 3.5 lb. | 3.5 lb. | 3.5 lb. | 3.5 lb. | 24h | NA |
| ferbam | | F | x | F | F | G | E | NA | 3 |
| Flint Extra | 11 | 2.9 fl. oz. | 2.5-2.9 fl. oz. | 2.5-2.9 fl. oz. | 2.5-2.9 fl. oz. | 2.9 fl. oz. | x | 12h | 10.5 fl. oz. |
| trifloxystrobin | | s[G] | G[r] | E[r] | G | s[G] | x | 14d | NA |
| Fontelis (SC) | 7 | x | 16-20 fl. oz. | 16-20 fl. oz. | x | 16-20 fl. oz. | x | 12h | 61 fl. oz. |
| penthiopyrad | | x | G | E | x | u | x | 28d | NA |
| Inspire Super (EW) | 3+9 | x | 12 fl. oz. | 12 fl. oz. | 12 fl. oz. | x | x | 12h | 60 fl. oz. |
| difenoconazole + cyprodinil | | s | F | E | E | x | x | 28d | NA |
| Kenja 400SC | 7 | x | 12.5 fl. oz. | 12.5 fl. oz. | x | x | x | 12h | NA |
| isofetamid | | x | s | F | x | x | x | 20d | NA |
| Kocide 3000 | M | x | x | 0.75 lb. | x | x | x | 48h | 53.3 lb. |
| copper hydroxide | | x | x | F-G | x | x | x | 0d | NA |
| Luna Privilege | 7 | x | 2.4-6.8 fl. oz. | 4-6.8 fl. oz. | 6.8 fl. oz. | x | x | NA | NA |
| fluopyram | | x | G | G-E | F-G | x | x | NA | NA |
| Luna Sensation (SC) | 7+11 | 4-5.8 fl. oz. | 5-5.8 fl. oz. | 4-5.8 fl. oz. | 4-5.8 fl. oz. | 4-5.8 fl. oz. | x | 12h | 21 fl. oz. |
| fluopyram + trifloxystrobin | | E | G[r] | E[r] | G-E | E | x | 14d | 4 |
| Merivon (2.09SC) | 7+11 | 4-5.5 fl. oz. | 4-5.5 fl. oz. | 4-5.5 fl. oz. | 4-5.5 fl. oz. | 4-5.5 fl. oz. | x | 12h | 22 fl. oz. |
| fluxapyroxad + pyraclostrobin | | E | G | E | E | E | x | 0d | 4 |
| Microthiol Disperss | M | 10-20 lb. | 10-20 lb. | 10-20 lb. | x | x | x | 24h | NA |
| sulfur | | u | G | i-F | x | x | x | 0d | NA |
| Miravis (1.67 SC) | 7 | x | 3.4 fl. oz. | 3.4 fl. oz. | 3.4 fl. oz. | x | x | 4h | 13.6 fl. oz. |
| pydiflumetofen | | x | G | E | G | x | x | 30d | 4 |

(Continued)

Table 2-7. Pear Disease - Summer Cover¹ (continued)

| Product And Formulation Active Ingredient | FRAC code ² | Bitter Rot | Powdery Mildew | Scab | Sooty Blotch/ Flyspeck | Summer Rots (Black And White) | Fabraea Leaf Spot | REI ³ PHI ⁴ | Max Amt ⁵ Max App ⁶ |
|--|---------------------------|------------------|-------------------|-------------------|------------------------------|---|----------------------|--------------------------------------|--|
| OSO 5% SC | 19 | 6.5 fl. oz. | 6.5-13 fl. oz. | 6.5-13 fl. oz. | 6.5 fl. oz. | 6.5 fl. oz. | x | 4h | 78 fl. oz. |
| polyoxin D | | F | F | F | F | F | x | 0d | 6 |
| Pristine | 11+7 | 14.5-18.5 oz. | 14.5-18.5 oz. | 14.5-18.5 oz. | 14.5-18.5 oz. | 14.5-18.5 oz. | x | 12h | 74 oz. |
| pyraclostrobin + boscalid | | F-G | E[r] | E[r] | E | E | x | 0d | 4 |
| Procure 480SC | 3 | x | 8-16 fl. oz. | 8-16 fl. oz. | x | x | x | 12h | 64 fl. oz. |
| triflumizole | | x | E[r] | G[r] | x | x | x | 14d | NA |
| Sovran (50WG) | 11 | x | 4-6.4 oz. | 3.2-6.4 oz. | 4-6.4 oz. | 4-6.4 oz. | x | 12h | 25.6 oz. |
| kresoxim-methyl | | x | G[r] | E[r] | G | G | x | 30d | 4 |
| Topguard Specialty Crops | 3 | 13 fl. oz. | 8-12 fl. oz. | 13 fl. oz. | x | 13 fl. oz. | x | 12h | 52 fl. oz. |
| flutriafol | | u | E | G-E | x | u | x | 14d | 4 |
| Topsin-M WSB | 1 | 1 lb. | 0.75-1 lb. | 0.75-1 lb. | 0.75-1 lb. | 0.75-1 lb. | 1 lb | 24h | 4 lb. |
| thiophanate methyl | | E[r] | G[r] | i | G-E [r] | G | G-E[R] | 1d | NA |
| Torino (SC) | U6 | x | 6.8 oz | x | x | x | x | 4h | 6.8 oz. |
| cyflufenamid | | x | E | x | x | x | x | 14d | 1 |
| Ziram 76DF | M3 | 6 lb. | x | 6 lb. | 6 lb. | 6 lb. | 6 | 48h | 42.4 lb. |
| ziram | | G | x | G | G | i | E | 14d | 7 |

Footnotes: See how to read the spray schedule tables section at the beginning of the chapter.

Pear First To Second Cover - Insects

10-14 days after petal fall and 10-14 days later.

Insect management notes

- Codling moth:** First and second cover are the key times to control codling moth, to kill young larvae as they emerge from eggs. Read the insecticide label to see if an adjuvant is recommended. Optimal timing for the first spray can be determined by using pheromone traps to monitor codling moth, and a degree-day model to track temperature after moth flight begins. See table in Apple chapter for degree-day targets for various insecticides.
- Pear psylla:** Pear psylla control is required in first or second covers only if control at petal-fall was not adequate. Best results occur when pear psylla is treated in its young nymphal stage.
- Pear rust mite:** Pear rust mite is not a problem in most orchards but can be a problem in young orchards or where pyrethroids have been used. First cover is the best time to control pear rust mite if not treated pre-bloom. If pear rust mite is present, note that Agri-Mek, Nexter, Magister and Portal control both pear psylla and pear rust mite.
- San Jose scale "crawlers":** Sprays should target the crawler stage, which usually begins in early summer, around the time of second or third cover. A pheromone trap and a degree-day model can be used to predict the timing of crawler emergence.

Table 2-8. Pear Insects - First Through Second Cover¹

| Product And Formulation Active Ingredient | IRAC Code ² | Pear Psylla NYMPH | Plum Curculio | Pear Rust Mite | Codling Moth | REI ³ PHI ⁴ | Max Amt ⁵ Max App ⁶ |
|---|---------------------------|----------------------|------------------|-------------------|-----------------------|--------------------------------------|--|
| Actara (25WDG) | 4A | 5.5 oz. | 4.5-5.5 oz. | x | x | 12h | 16.5 oz. |
| thiamethoxam | | E | G | x | x | 35d | NA |
| Admire Pro (4.6F) | 4A | 7 fl. oz. | x | x | x | 12h | 14 fl. oz. |
| imidacloprid | | G | x | x | x | 7d | NA |
| Agri-Mek SC (0.7SC) (RUP) | 6 | 1.3-2.2 oz. | x | 1.3-2.2 oz. | x | 12h | 4.6 fl. oz. |
| abamectin | | G | x | E | x | 28d | 6 |
| Altacor eVo | 28 | x | 2.5-4.5 oz. | x | 2.5-4.5 oz. | 4h | 9 fl. oz. |
| chlorantraniliprole | | x | s | x | E | 5d | NA |
| Apta (1.34SC) | 21A | 21-27 fl. oz. | 21-27 fl. oz. | 21-27 fl. oz. | 21-27 fl. oz. | 12h | 53.5 fl. oz. |
| tolfenpyrad | | G | G | u | s | 14d | 2 |
| Asana XL (0.66EC) (RUP) | 3A | x | 4.8-14.5 fl. oz. | x | 4.8-14.5 fl. oz. | 12h | 72 fl. oz. |
| esfenvalerate | | x | F | x | G | 28d | NA |
| Assail 30SG | 4A | 4-8 oz. | 8 oz. | x | 4-8 oz. | 12h | 32 oz. |
| acetamiprid | | G | G | x | E | 7d | 4 |
| <i>Bacillus thuringiensis</i> (B.t.) (Dipel DF, etc.) | 11A | x | x | x | 0.5-2 lb. | 4h | NA |
| <i>Bacillus thuringiensis</i> | | x | x | x | F | 0d | NA |
| Baythroid XL (1EC) (RUP) | 3A | x | 2.4-2.8 fl. oz. | x | 2-2.4 fl. oz. | 12h | 2.8 fl. oz. |
| beta-cyfluthrin | | x | F | x | G | 7d | NA |
| Belay (2.13SC) | 4A | 6 fl. oz. | 6 fl. oz. | x | 6 fl. oz. | 12h | 12 fl. oz. |
| clothianidin | | G | G | x | s | 7d | NA |
| Brigade eVo (RUP) | 3A | x | 2.6-12.8 oz. | x | 2.6-12.8 oz. | 12h | 32 oz. |
| bifenthrin | | x | F | x | G | 14d | NA |
| Centaur WDG (70WDG) | 16 | 34.5-46 oz. | x | x | x | 12h | 69 oz. |
| buprofezin | | u | x | x | x | 14d | 2 |
| Confirm 2F | 18 | x | x | x | 20 fl. oz. | 4h | 20 fl. oz. |
| tebufenozide | | x | x | x | F | 14d | 120 fl. oz. |
| Danitol 2.4EC (RUP) | 3A | x | x | x | 16-21.3 fl. oz. | 24h | 42.7 fl. oz. |
| fenpropathrin | | x | x | x | G | 14d | NA |
| Delegate WG (25WG) | 5 | 6-7 oz. | 6-7 oz. | x | 4.5-7 oz. | 4h | 28 oz. |
| spinetoram | | E | s | x | E | 7d | 4 |
| Delta Gold (1.5EC) (RUP) | 3A | x | 0.9-1.9 fl. oz. | x | 0.9-1.9 fl. oz. | 12h | 3.6 fl. oz. |
| deltamethrin | | x | F | x | G | 21d | NA |
| Diazinon AG 600WBC (RUP) | 1B | x | x | x | 12.7 fl. oz./100 gal. | 4d | 102 fl. oz. |
| diazinon | | x | x | x | F | 21d | 2 |
| Dimilin 2L (2AF) (RUP) | 15 | 12-48 fl. oz. | x | 40-48 fl. oz. | 12-16 fl. oz. | 12h | 64 fl. oz. |
| diflubenzuron | | E | x | u | u | 14d | 4 |
| Entrust SC (2SC) | 5 | x | x | x | 6-10 fl. oz. | 4h | 29 fl. oz. |
| spinosad | | x | x | x | F | 7d | 4 |

(Continued)

Table 2-8. Pear Insects - First Through Second Cover¹ (continued)

| Product And Formulation Active Ingredient | IRAC Code ² | Pear Psylla NYMPH | Plum Curculio | Pear Rust Mite | Codling Moth | REI ³ PHI ⁴ | Max Amt ⁵ Max App ⁶ |
|--|---------------------------|----------------------|-------------------|-------------------|-----------------|--------------------------------------|--|
| Envidor 2SC | 23 | x | x | 16-18 fl. oz. | x | 12h | 18 fl. oz. |
| spirodiclofen | | x | x | E | x | 7d | 1 |
| Esteem 35WP | 7C | 5 oz. | x | x | 4-5 oz. | 12h | 10 oz. |
| pyriproxyfen | | G | x | x | G | 45d | 2 |
| Exirel (0.83SE) | 28 | 13.5-20.5 fl. oz. | 13.5-20.5 fl. oz. | x | 8.5-17 fl. oz. | 12h | 61 fl. oz. |
| cyantraniliprole | | s | G | x | E | 3d | 3 |
| Grandevo | UN | 2-3 lb | x | 2-3 lb. | 1-3 lb. | 4h | NA |
| <i>Chromobacterium subtsugae</i> | | u | x | u | F | 0d | NA |
| Imidan 70W | 1B | x | 2.1-5.7 lb. | x | 2.1-5.7 lb. | 7d | 16 lb. |
| phosmet | | x | G | x | G | 7d | NA |
| Intrepid 2F | 18 | x | x | x | 16 fl. oz. | 4h | 64 fl. oz. |
| methoxyfenozide | | x | x | x | s | 14d | NA |
| Magister SC (1.7SC) | 21A | 32-36 fl. oz. | x | 32-36 fl. oz. | x | 12h | 36 fl. oz. |
| fenazaquin | | G | x | G | x | 7d | 1 |
| Mating disruption | UN | x | x | x | see apple | 0h | NA |
| Isomate, Checkmate | | x | x | x | G | 0d | NA |
| Movento (2SC) | 23 | 6-9 fl. oz. | x | 6-9 fl. oz. | 6-9 fl. oz. | 24h | 25 fl. oz. |
| spirotetramat | | G | x | u | s | 7d | NA |
| Mustang Maxx (0.8EC) (RUP) | 3A | x | 1.2-4 fl. oz. | x | 1.2-4 fl. oz. | 12h | 24 fl. oz. |
| zeta-cypermethrin | | x | F | x | G | 14d | NA |
| Neemix 4.5 (0.39L) | UN | 7-16 fl. oz. | x | x | 4-16 fl. oz. | 4h | NA |
| azadirachtin | | F | x | x | F | 0d | NA |
| Nexter SC (3.75SC) | 21A | 11-17 fl. oz. | x | 11-17 fl. oz. | x | 12h | NA |
| pyridaben | | G | x | E | x | 7d | 1 |
| Portal XLO (0.4EC) | 21A | 2 pt. | x | 2 pt. | x | 12h | 2 pt. |
| fenpyroximate | | G | x | G | x | 14d | 1 |
| Proclaim (5SG) (RUP) | 6 | 3.2-4.8 oz. | x | x | 4.8 oz. | 12 or 48h | 14.4 oz. |
| emamectin benzoate | | s | x | x | F | 14d | NA |
| Sevin XLR Plus (4F) | 1A | 1.5-3 qt. | 1.5-3 qt. | 1.5-3 qt. | 3 qt. | 12h | 15 qt. |
| carbaryl | | u | F | G | F | 3d | 8 |
| Sivanto Prime (1.67SC) | 4D | 10.5-14 fl. oz. | x | x | x | 12h | 28 fl. oz. |
| flupyradifurone | | G | x | x | x | 14d | NA |
| Soap (M-Pede, Des-X, etc.) | UN | 2% | x | 2% | x | 12h | NA |
| potassium salts of fatty acids | | F | x | u | x | 0d | NA |
| Surround WP (95WP) | UN | 50 lb. | x | x | 25-50 lb. | 4h | NA |
| kaolin | | G | x | x | s | 0d | NA |
| Transform WG (50WG) | 4C | 2.75 oz. | x | x | x | 24h | 8.5 oz. |
| sulfoxaflor | | s | x | x | x | 7d | 4 |
| Vendex 50WP (RUP) | 12B | x | x | 1-2 lb. | x | 48h | 4 lb. |
| fenbutatin-oxide | | x | x | G | x | 14d | 2 |

(Continued)

Table 2-8. Pear Insects - First Through Second Cover¹ (continued)

| Product And Formulation Active Ingredient | IRAC Code ² | Pear Psylla NYMPH | Plum Curculio | Pear Rust Mite | Codling Moth | REI ³ PHI ⁴ | Max Amt ⁵ Max App ⁶ |
|--|---------------------------|----------------------|------------------|-------------------|-----------------|--------------------------------------|--|
| Verdepryn 100SL (0.83SL) | 28 | 11 fl. oz. | 5.5-11 fl. oz. | x | 5.5-11 fl. oz. | 4h | 33 fl. oz. |
| cyclaniliprole | | u | G | x | E | 7d | 3 |
| Virus (Cyd-X HP) | 31 | x | x | x | 0.5-3 fl. oz. | 4h | NA |
| <i>Cydia pomonella</i> granulovirus | | x | x | x | G | 0d | NA |
| Virus (Madex HP) | 31 | x | x | x | 0.5-3 fl. oz. | 4h | NA |
| <i>Cydia pomonella</i> granulovirus | | x | x | x | G | 0d | NA |
| Warrior II (2.08CS) (RUP) | 3A | x | 1.2-2.5 fl. oz. | x | 1.2-2.5 fl. oz. | 24h | 12.8 fl. oz. |
| lambda-cyhalothrin | | x | G | x | G | 21d | NA |

Footnotes: See how to read the spray schedule tables section at the beginning of the chapter.

Pear Summer Covers - Insects

Apply at 10- to 14-day intervals observing harvest restrictions and limitations.

Insect management notes

- Pear psylla:** For cultural control, it is very important to prune water sprouts, which appear in early summer. This removal destroys the favorite habitat of pear psylla. Make two applications 10-12 days apart to target second-generation young nymphs, in early summer, usually in about mid-June. The first new summer adults appear about three weeks after full bloom. Adults are found on terminals and water sprouts. The adults are more difficult to control than young nymphs. The first new summer adults appear about three weeks after full bloom. Adults are found on terminals and water sprouts. Third generation nymphs can be controlled in mid-summer, around mid-July.
- San Jose scale:** Sprays should target the crawler stage, which usually begins in early summer, around the time of second or third cover. A pheromone trap and a degree-day model can be used to predict the timing of crawler emergence.
- Codling moth:** Fifth and sixth cover sprays target the new generation of young larvae as they emerge from eggs.
- Stink bug:** The time that stink bugs arrive is variable, sometimes in the pre-bloom period, sometimes in early summer, sometimes not until late summer. Native stink bugs can be present pre-bloom or post-bloom. At locations where brown marmorated stink bug is present, it is most commonly controlled in third, fourth and fifth cover sprays.

Table 2-9. Pear Insects - Summer Cover¹

| Product And Formulation Active Ingredient | IRAC Code ² | European Red Mite | Mealybug | Pear Psylla NYMPH | San Jose Scale CRAWLERS | Codling Moth | Stink Bug | REI ³ PHI ⁴ | Max Amt ⁵ Max App ⁶ |
|--|---------------------------|----------------------|-------------|-------------------------|-------------------------------|-----------------|--------------|--------------------------------------|--|
| Acramite 50WS | 20D | 1 lb. | x | x | x | x | x | 12h | 3 |
| bifenazate | | G | x | x | x | x | x | 7d | 1 |
| Actara (25WDG) | 4A | x | 4.5-5.5 oz. | 5.5 oz. | x | x | x | 12h | 16.5 oz. |
| thiamethoxam | | x | G | E | x | x | x | 35d | NA |
| Admire Pro (4.6F) | 4A | x | 7 fl. oz. | 7 fl. oz. | 2.8 fl. oz. | x | x | 12h | 14 fl. oz. |
| imidacloprid | | x | G | G | F | x | x | 7d | NA |
| Agri-Mek SC (0.7SC) (RUP) | 6 | 2.2-4.2 fl. oz. | x | 2.2-4.2 fl. oz. | x | x | x | 12h | 8.5 fl. oz. |
| abamectin | | E | x | G | x | x | x | 28d | 2 |

(Continued)

Table 2-9. Pear Insects - Summer Cover¹ (continued)

| Product And Formulation Active Ingredient | IRAC Code ² | European Red Mite | Mealybug | Pear Psylla NYMPH | San Jose Scale CRAWLERS | Codling Moth | Stink Bug | REI ³ PHI ⁴ | Max Amt ⁵ Max App ⁶ |
|--|---------------------------|----------------------|---------------|-------------------------|-------------------------------|---------------------|-------------------------|--------------------------------------|--|
| Altacor eVo | 28 | x | x | x | x | 1.3-2.2 oz. | x | 4h | 4.6 oz. |
| chlorantraniliprole | | x | x | x | x | E | x | 5d | 3 |
| Apollo SC (ISC) | 10A | 4-8 fl. oz. | x | x | x | x | x | 12h | NA |
| clofentezine | | E | x | x | x | x | x | 21d | 1 |
| Apta (1.34SC) | 21A | x | 21-27 fl. oz. | 21-27 fl. oz. | x | 21-27 fl. oz. | x | 12h | 53.5 fl. oz. |
| tolfenpyrad | | x | u | G | x | s | x | 14d | 2 |
| Asana XL (0.66EC) (RUP) | 3A | x | x | x | x | 4.8-14.5 fl. oz. | x | 12h | 72 fl. oz. |
| esfenvalerate | | x | x | x | x | G | x | 28d | NA |
| Assail 30SG | 4A | x | 4-8 oz. | 4-8 oz. | 8 oz. | 4-8 oz | x | 12h | 32 oz. |
| acetamiprid | | x | G | G | s | E | x | 7d | 4 |
| Avaunt eVo (30WDG) | 22 | x | x | x | x | 5-6 oz. | x | 12h | 24 oz. |
| indoxacarb | | x | x | x | x | G | x | 28d | 4 |
| <i>Bacillus thuringiensis</i> (B.t.) (Dipel DF, etc.) | 11A | x | x | x | x | 0.5-2 lb. | x | 4h | NA |
| <i>Bacillus thuringiensis</i> | | x | x | x | x | F | x | 0d | NA |
| Baythroid XL (IEC) (RUP) | 3A | x | x | x | 2.4-2.8 fl. oz. | 2-2.4 fl. oz. | 2-2.4 fl. oz. | 12h | 2.8 fl. oz. |
| beta-cyfluthrin | | x | x | x | F | G | G | 7d | NA |
| Belay (2.13SC) | 4A | x | 6 fl. oz. | 6 fl. oz. | 6 fl. oz. | 6 fl. oz. | 6 fl. oz. | 12h | 12 fl. oz. |
| clothianidin | | x | G | G | G | s | E | 7d | NA |
| Brigade eVo (RUP) | 3A | 5.1-12.8 fl. oz. | x | x | 2.6-12.8 fl. oz. | 2.6-12.8 fl. oz. | 2.6- 12.8 fl. oz. | 12h | 32 oz. |
| bifenthrin | | F | x | x | F | G | E | 14d | 3 |
| Centaur WDG (70WDG) | 16 | x | 34.5-46 oz. | 34.5-46 oz. | 34.5-46 oz. | x | x | 12h | 69 oz. |
| buprofezin | | x | G | u | E | x | x | 14d | 2 |
| Confirm 2F | 18 | x | x | x | x | 20 fl. oz. | x | 4h | 120 fl. oz. |
| tebufenozide | | x | x | x | x | F | x | 14d | NA |
| Damoil | UN | see label | x | x | x | 0.5-4% | x | 4h | NA |
| mineral oil | | G | x | x | x | u | x | 0d | NA |
| Danitol 2.4EC (RUP) | 3A | x | x | x | x | 16-21.3 fl. oz. | 16-21.3 fl. oz. | 24h | 42.7 fl. oz. |
| fenpropathrin | | x | x | x | x | G | G | 14d | NA |
| Delegate WG (25WG) | 5 | x | x | 6-7 oz. | x | 4.5-7 oz. | x | 4h | 28 oz. |
| spinetoram | | x | x | E | x | E | x | 7d | 4 |
| Delta Gold (1.5EC) (RUP) | 3A | x | x | x | 0.9-1.9 fl. oz. | 0.9-1.9 fl. oz. | 1.9 fl. oz. | 12h | 3.6 fl. oz. |
| deltamethrin | | x | x | x | u | G | u | 21d | NA |

(Continued)

Table 2-9. Pear Insects - Summer Cover¹ (continued)

| Product And Formulation Active Ingredient | IRAC Code ² | European Red Mite | Mealybug | Pear Psylla NYMPH | San Jose Scale CRAWLERS | Codling Moth | Stink Bug | REI ³ PHI ⁴ | Max Amt ⁵ Max App ⁶ |
|--|---------------------------|---------------------------|---------------------------|-------------------------|-------------------------------|---------------------------|------------------|--------------------------------------|--|
| Diazinon AG 600WBC (RUP) | 1B | 12.7 fl. oz./ 100 gal. | 12.7 fl. oz./ 100 gal. | x | 12.7 fl. oz./ 100 gal. | 12.7 fl. oz./ 100 gal. | x | 4d | 102 fl. oz. |
| diazinon | | u | G | x | G | F | x | 21d | 2 |
| Dimethoate (4EC) | 1B | 0.5-1 pt./100 gal. | x | x | x | x | x | 10d | 2 pt. |
| dimethoate | | u | x | x | x | x | x | 28d | NA |
| Dimilin 2L (2AF) (RUP) | 15 | x | x | 12-48 fl. oz. | x | 12-16 fl. oz. | x | 12h | 64 fl. oz. |
| diflubenzuron | | x | x | E | x | u | x | 14d | 4 |
| Entrust SC (2SC) | 5 | x | x | x | x | 6-10 fl. oz. | x | 4h | 29 fl. oz. |
| spinosad | | x | x | x | x | F | x | 7d | 4 |
| Envidor 2SC | 23 | 16-18 fl. oz. | x | x | x | x | x | 12h | 18 fl. oz. |
| spiroadiclofen | | E | x | x | x | x | x | 7d | 1 |
| Esteem 35WP | 7C | x | x | 5 oz. | 4-5 oz. | 5 oz. | x | 12h | 10 oz. |
| pyriproxyfen | | x | x | G | E | G | x | 45d | 2 |
| Exirel (0.83SE) | 28 | x | x | 13.5-20.5 fl. oz. | x | 8.5-17 fl. oz. | x | 12h | 61 fl. oz. |
| cyantraniliprole | | x | x | s | x | E | x | 3d | 3 |
| Grandevo | UN | 2-3 lb. | 2-3 lb. | 2-3 lb. | 2-3 lb. | 1-3 lb. | x | 4h | NA |
| <i>Chromobacterium subtsugae</i> | | u | u | u | u | F | x | 0d | NA |
| Imidan 70W | 1B | x | 2.1-5.7 lb. | x | x | 2.1-5.7 lb. | x | 7d | 16 lb. |
| phosmet | | x | F | x | x | G | x | 7d | NA |
| Intrepid 2F | 18 | x | x | x | x | 16 fl. oz. | x | 4h | 64 fl. oz. |
| methoxyfenozide | | x | x | x | x | s | x | 14d | NA |
| Kanemite 15SC | 20B | 21-31 fl. oz. | x | x | x | x | x | 12h | 62 fl. oz. |
| acequinocyl | | E | x | x | x | x | x | 14d | 2 |
| Magister SC (1.7SC) | 21A | 32-36 fl. oz. | x | 32-36 fl. oz. | x | x | x | 12h | 36 fl. oz. |
| fenazaquin | | G | x | G | x | x | x | 7d | 1 |
| Mating disruption | UN | x | x | x | x | see apple | x | 0h | NA |
| Isomate, Checkmate | | x | x | x | x | G | x | 0d | NA |
| Movento (2SC) | 23 | 6-9 fl. oz. | 6-9 fl. oz. | 6-9 fl. oz. | 6-9 fl. oz. | 6-9 fl. oz. | x | 24h | 25 fl. oz. |
| spirotetramat | | s | G | G | E | s | x | 7d | NA |
| Mustang Maxx (0.8EC) (RUP) | 3A | x | x | x | x | 1.2-4 fl. oz. | 1.2-4 fl. oz. | 12h | 24 fl. oz. |
| zeta-cypermethrin | | x | x | x | x | G | G | 14d | NA |
| Nealta (1.67SC) | 25 | 13.7 fl. oz. | x | x | x | x | x | 12h | 27.4 fl. oz. |
| cyflumetofen | | E | x | x | x | x | x | 7d | 2 |
| Neemix 4.5 (0.39L) | UN | x | 7-16 fl. oz. | 7-16 fl. oz. | 6-16 fl. oz. | 4-16 fl. oz. | 7-16 fl. oz. | 4h | NA |
| azadirachtin | | x | i | F | u | F | u | 0d | NA |

(Continued)

Table 2-9. Pear Insects - Summer Cover¹ (continued)

| Product And Formulation Active Ingredient | IRAC Code ² | European Red Mite | Mealybug | Pear Psylla NYMPH | San Jose Scale CRAWLERS | Codling Moth | Stink Bug | REI ³ PHI ⁴ | Max Amt ⁵ Max App ⁶ |
|--|---------------------------|----------------------|----------|-------------------------|-------------------------------|--------------------|--------------------|--------------------------------------|--|
| Nexter SC (3.75SC) | 21A | 11-17 fl. oz. | x | 11-17 fl. oz. | x | x | x | 12h | NA |
| pyridaben | | E | x | G | x | x | x | 7d | 1 |
| Onager Optek (1EC) | 10 | 12-24 fl. oz. | x | x | x | x | x | 12h | 24 fl. oz. |
| hexythiazox | | E | x | x | x | x | x | 28d | 1 |
| Portal XLO (0.4EC) | 21A | 2 pt. | 2 pt. | 2 pt. | x | x | x | 12h | 2 pt. |
| fenpyroximate | | E | u | G | x | x | x | 14d | 1 |
| Proclaim (5SG) (RUP) | 6 | 3.2-4.8 oz. | x | 3.2-4.8 oz. | x | 4.8 oz. | x | 12 or 48h | 14.4 oz. |
| emamectin benzoate | | s | x | s | x | F | x | 14d | NA |
| Sevin XLR Plus (4F) | 1A | x | x | 1.5-3 qt. | 1.5-3 qt. | 3 qt. | x | 12h | 15 qt. |
| carbaryl | | x | x | u | F | F | x | 3d | 8 |
| Sivanto Prime (1.67SC) | 4D | x | x | 10.5-14 fl. oz. | 10.5-14 fl. oz. | x | x | 12h | 28 fl. oz. |
| flupyradifurone | | x | x | G | G | x | x | 14d | NA |
| Soap (M-Pede, Des-X, etc.) | UN | 2% | 2% | 2% | 2% | x | x | 12h | NA |
| potassium salts of fatty acids | | u | G | F | F | x | x | 0d | NA |
| Sulfur (Microfine; 90%) | UN | 10-60 lb. | x | x | x | x | x | 24h | NA |
| sulfur | | u | x | x | x | x | x | 0d | NA |
| Surround WP (95WP) | UN | x | x | 50 lb. | x | 25-50 lb. | 25-50 lb. | 4h | NA |
| kaolin | | x | x | G | x | s | s | 0d | NA |
| Transform WG (50WG) | 4C | x | x | 2.75 oz. | 2.75 oz. | x | x | 24h | 8.5 oz. |
| sulfoxaflor | | x | x | s | s | x | x | 7d | 4 |
| Vendex 50WP (RUP) | 12B | 1-2 lb. | x | x | x | x | x | 48h | 4 lb. |
| fenbutatin-oxide | | G | x | x | x | x | x | 14d | 2 |
| Verdepryn 100SL (0.83SL) | 28 | x | x | 11 fl. oz. | x | 5.5-11 fl. oz. | 5.5-11 fl. oz. | 4h | 33 fl. oz. |
| cyclaniliprole | | x | x | u | x | E | s | 7d | 3 |
| Virus (Cyd-X HP) | 31 | x | x | x | x | 0.5-3 fl. oz. | x | 4h | NA |
| <i>Cydia pomonella</i> granulovirus | | x | x | x | x | G | x | 0d | NA |
| Virus (Madex HP) | 31 | x | x | x | x | 0.5-3 fl. oz. | x | 4h | NA |
| <i>Cydia pomonella</i> granulovirus | | x | x | x | x | G | x | 0d | NA |
| Vydate L (2L) (RUP) | 1A | 6-8 pt. | x | x | x | x | 1.5-4 pt. | 48h | 8 pt. |
| oxamyl | | G | x | x | x | x | G | 14d | 1 |
| Warrior II (2.08CS) (RUP) | 3A | x | x | x | 1.2-2.5 fl. oz. | 1.2-2.5 fl. oz. | 1.2-2.5 fl. oz. | 24h | 12.8 fl. oz. |
| lambda-cyhalothrin | | x | x | x | F | G | G | 21d | NA |
| Zeal (72WP) | 10B | 2-3 oz. | x | x | x | x | x | 12h | 3 oz. |
| etoxazole | | E | x | x | x | x | x | 14d | 1 |

Footnotes: See how to read the spray schedule tables section at the beginning of the chapter.

3. CHERRY

Cherry Spray Schedule

The shaded/colored boxes represent the crop stages where common pests in the Midwest are active. Scouting and/or preventative sprays may be necessary or recommended. Weather and degree day accumulation will impact the exact timing of pest appearance in the orchard.

| Stage | | | | | | | |
|-------------------|---------------------|------------------|-------------------|-------------------|------------------|-------------------------|-------------------|
| Dormant | Early To Full Bloom | Petal Shuck Fall | First Cover | Second Cover | Additional Cover | Pre-Harvest | Post-Harvest |
| | | | | | Borers | | |
| | | | Cherry Fruit Fly | | | | |
| | | Plum Curculio | | | | | |
| | | | | | | | Japanese Beetle |
| | | | | | | | Periodical Cicada |
| | | | | | | Spotted-wing Drosophila | |
| European Red Mite | | | | European Red Mite | | | |
| | | | Periodical Cicada | | | | |

How to read the spray schedule tables

Every cherry growth stage has important notes on disease or insect management. In some cases, the reader will be directed to the special problems section at the end of the section or chapter. Please make sure to read thoroughly and contact your state Extension specialist with any specific questions.

Key to tables

E = excellent control

G = good control

F = fair control

[r] = fungicide/insecticide resistance possible

s = suppression only

i = ineffective

u = unknown efficacy

x = pest not on the label

¹ Efficacy data in this publication are based on trials conducted across various regions and does not necessarily reflect local efficacy differences or changes over time. Growers should contact their Extension specialist for the most recent or for state-specific information. The information on this publication is only a guide; the authors and their institutions assume no liability for

practices implemented based on this information. Always read and follow pesticide labels. The label is the law. Product registration may vary by state.

² FRAC/IRAC code represents the mode of action of the fungicide/insecticide.

³ PHI refers to the pre-harvest interval, which is the number of days before harvest that the product may not be applied.

⁴ All fungicides/insecticides have a Restricted-Entry Interval (REI). The restricted-entry interval is the time immediately after a pesticide application when entry into the treated area is limited. Check labels for REI. Restrictions in REI may prohibit the use of certain pesticides during harvest.

Applicators must abide by both maximum amount of product per season AND maximum number of applications.

⁵ Max amt refers to the product's maximum amount/acre/year.

⁶ Max app refers to the product's maximum number of applications per year.

RUP refers to restricted use pesticide.

Cherry Dormant - Diseases

Before buds break in the spring.

Notes on disease management

- Bacterial canker:** Bacterial canker is generally more serious on sweet than tart cherry. Dormant copper sprays are an effective method for control of bacterial canker. All stone fruit, including cherry, are extremely sensitive to copper. Sprays must be timed to reduce *Pseudomonas syringae* inoculum without causing phytotoxicity. Tart cherries may require continued protection through bloom.

- Application of copper for bacterial canker at this time may reduce the overwintering inoculum of the cherry leaf spot pathogen.
- Make first application before fall rains and a second at late dormant.
- Phytophthora:** See Phytophthora Disease Management of Stone Fruit on page 153.

Table 3-1. Cherry Diseases - Dormant¹

| Product And Formulation | Active Ingredient | FRAC Code ² | Bacterial Canker | Cherry Leaf Spot Dormant | REI ³ PHI ⁴ | Max Amt ⁵ Max App ⁶ |
|----------------------------|------------------------------|------------------------|------------------|--------------------------|-----------------------------------|---|
| Badge SC | | M | 3.5-14 pt. | 3.5-14 pt. | 24h | 63.4 pt. |
| | copper sulfate + oxychloride | | P-G | F | 0d | NA |
| C-O-C-S WDG | | M | x | 8-15 lb. | 48h | 35 lb. |
| | copper oxychloride | | x | F | 21d | 3 |
| Cuprofix Ultra 40 Disperss | | M | 5-8 lb. | 5-8 lb. | 48h | 45 lb. |
| | copper sulfate | | P-G | P-G | 120d | NA |
| Kocide 3000 | | M | x | 2.2-3.5 lb. | 48h | 60 lb. |
| | copper hydroxide | | x | G | 0d | 4 |

Footnotes: See how to read the spray schedule tables section at the beginning of the chapter.

Cherry Late Dormant to Prebloom - Insects

Notes on insect management

- European red mite:** If European red mites have been a problem in the past, apply superior oil or Envior 2SC during the dormant stage to control mite eggs. Other miticides will be more effective if delayed until eggs are hatching (petal-fall). Oil

applied by late dormant will also control aphid eggs. When spraying oil, check labels for temperature restrictions (e.g., only when temperatures are above 40F° or never during freezing weather).

- San Jose scale:** Generally controlled dormant to delayed dormant where they have been a problem. Do not use Imidan 70W on sweet cherries.

Table 3-2. Cherry Insects - Dormant Through Prebloom¹

| Product And Formulation | Active Ingredient | IRAC Code ² | Aphid | European Red Mite | San Jose Scale | REI ³ PHI ⁴ | Max Amt ⁵ Max App ⁶ |
|---------------------------|-------------------|------------------------|---------------|-------------------|----------------|-----------------------------------|---|
| Acramite 50WS | | 20D | x | 1 lb. | x | 12h | 1 lb. |
| | bifenazate | | x | G | x | 3d | 1 |
| Actara (25WDG) | | 4A | 3-4 oz. | x | x | 12h | 11 oz. |
| | thiamethoxam | | E | x | x | 14d | NA |
| Agri-Mek SC (0.7SC) (RUP) | | 6 | x | 2.25-4.25 fl. oz. | x | 12h | 8.5 fl. oz. |
| | abamectin | | x | G | x | 21d | 2 |
| Apta (1.34SC) | | 21A | 17-21 fl. oz. | x | x | 12h | 53.5 fl. oz. |
| | tolfenpyrad | | u | x | x | 14d | 2 |

(Continued)

Table 3-2. Cherry Insects - Dormant Through Prebloom¹ (continued)

| Product And Formulation Active Ingredient | IRAC Code ² | Aphid | European Red Mite | San Jose Scale | REI ³ PHI ⁴ | Max Amt ⁵ Max App ⁶ |
|--|------------------------|------------------------|----------------------------|-------------------|--------------------------------------|--|
| Apollo SC (1SC) | 10A | x | 2-8 fl. oz. | x | 12h | NA |
| clofentezine | | x | E | x | 21d | 1 |
| Asana XL (0.66EC) (RUP) | 3A | 4.8-14.5 fl. oz. | x | x | 12h | 72.5 fl. oz. |
| esfenvalerate | | u | x | x | 14d | NA |
| Assail 30SC | 4A | 2.1-4.5 fl. oz. | x | 4.5-6.7 fl. oz. | 12h | 32 oz. |
| acetamiprid | | E | x | F | 7d | 4 |
| Baythroid XL (1EC) (RUP) | 3A | 2.4-2.8 fl. oz. | x | x | 12h | 5.6 fl. oz. |
| beta-cyfluthrin | | E | x | x | 7d | NA |
| Beleaf 50SG | 29 | 2-2.8 oz. | x | x | 12h | 8.4 fl. oz. |
| flonicamid | | E | x | x | 14d | 3 |
| Centaur WDG (70WDG) | 16 | x | x | 34.5 oz. | 12h | 69 oz. |
| buprofezin | | x | x | E | 14d | 2 |
| Danitol 2.4EC (RUP) | 3A | 10.3-21.3 fl. oz. | 10.3-21.3 fl. oz. | x | 24h | 42.7 fl. oz. |
| fenpropathrin | | u | F | x | 3d | NA |
| Diazinon AG600 WBC (RUP) | 1B | 12.75 fl. oz./100 gal. | 6.5-12.25 fl. oz./100 gal. | x | 4d | 102 fl. oz. |
| diazinon | | u | u | x | 21d | 2 |
| Envidor 2SC | 23 | x | 16-18 fl. oz. | x | 12h | 18 fl. oz. |
| spirodiclofen | | x | E | x | 7d | 1 |
| Esteem 35WP | 7C | x | x | 4-5 oz. | 12h | 15 oz. |
| pyriproxifen | | x | x | E | 14d | 3 |
| Exirel (0.83SE) | 28 | 13.5-20.5 fl. oz. | x | x | 12h | 61.5 fl. oz. |
| cyantraniliprole | | E | x | x | 3d | 3 |
| Imidan 70W | 1B | x | x | 2.13-4.25 lb. | 7/14d | 13 lb. |
| phosmet | | x | x | E | 7d | NA |
| Malathion 5EC | 1B | 2.8 pt. | x | x | 12h | NA |
| malathion | | G | x | x | 3d | 4 |
| Mustang Maxx (0.83EC) (RUP) | 3A | 1.28-4 fl. oz. | x | x | 12h | 24 fl. oz. |
| zeta-cypermethrin | | u | x | x | 3d | 6 |
| Nealta | 25 | x | 13.7 | x | 12h | 27.4 |
| cyflumetofen | | x | u | x | 7d | 2 |
| Neemix 4.5 | UN | 5-7 fl. oz. | x | x | 4h | NA |
| azadirachtin | | F | x | x | 0d | NA |
| Nexter (75WP) | 21 | x | 4.4-10.7 oz. | x | 12h | 21.3 oz. |
| pyridaben | | x | G | x | 7d | 2 |
| Oil (superior) | UN | x | see label | see label | 4h | NA |
| mineral oil | | x | E | G | 0d | NA |
| Onager Optek (1EC) | 10A | x | 12-24 oz. | x | 12h | 24 fl. oz. |
| hexythiazox | | x | E | x | 7d | 1 |

(Continued)

Table 3-2. Cherry Insects - Dormant Through Prebloom¹ (continued)

| Product And Formulation Active Ingredient | IRAC Code ² | Aphid | European Red Mite | San Jose Scale | REI ³ PHI ⁴ | Max Amt ⁵ Max App ⁶ |
|--|------------------------|------------------|----------------------|-------------------|--------------------------------------|--|
| Portal XLO (0.4EC) | 21A | x | 2 pt. | x | 12h | 4 pt. |
| fenpyroximate | | x | E | x | 7d | 2 |
| PQZ (1.87SC) | 9B | 2.4-3.2 fl. oz. | x | x | 12h | 4.8 fl. oz. |
| pyrifluquinazon | | E | x | x | 7h | 2 |
| Pyganic 5EC | 3A | 4.5-15.6 fl. oz. | 4.5-15.6 fl. oz. | 4.5-15.6 fl. oz. | 12h | NA |
| pyrethrins | | u | u | u | 0d | 10 |
| Savey 50DF | 10A | x | 3-6 oz. | x | 12h | 6 oz. |
| hexythiazox | | x | E | x | 28d | 1 |
| Sevin XLR Plus (4F) | 1A | 2-3 qt. | x | 4-5 qt. | 12h | 14 qt. |
| carbaryl | | u | x | u | 3d | 3 |
| Sivanto Prime (1.67SC) | 4D | 7-14 fl. oz. | x | 10.5-14 fl. oz. | 4h | 28 fl. oz. |
| flupyradifurone | | E | x | G | 14d | NA |
| Transform 75 WG | 4C | 0.75-1.5 oz. | x | 2.75 oz. | | 8.5 oz. |
| sulfoxaflor | | E | x | x | 7d | 4 |
| Vendex 50WP (RUP) | 12B | x | 1-2 lb. | x | 2d | 3 lb. |
| fenbutatin-oxide | | x | G | x | 14d | 2 |
| Versys Inscalis (0.83DC) | 9D | 1.5 fl. oz. | x | x | 12h | 3 fl. oz. |
| afidopyropen | | G | x | x | 7d | NA |
| Warrior II (2.08CS) (RUP) | 3A | 1.2-2.5 fl. oz. | x | x | 24h | 12.8 fl. oz. |
| lambda-cyhalothrin | | u | x | x | 14d | NA |
| Zeal (72WP) | 10B | x | 2-3 oz. | x | 12h | 3 oz. |
| etoxazole | | x | E | x | 7d | 1 |

Footnotes: See how to read the spray schedule tables section at the beginning of the chapter.

Cherry Early Bloom To Petal Fall And Shuck Split - Diseases

Early bloom is also referred to as popcorn or white bud, followed by bloom. Petal fall describes when flower petals are dropping. Shuck split describes when the shucks fall from expanding fruit.

Disease management notes

The two major diseases of concern are brown rot and leaf spot. Many fungicides are labeled for disease management at this time.

- Rovral is recommended early, as use after petal fall is prohibited.
- Vanguard is only labeled for tart cherries. Make the second and final application at full bloom.
- Bravo may not be used after shuck split.

Brown rot: Infection is rare at early bloom but may occur if brown rot cankers and mummies are present coupled with warm (above 60F°), wet weather. It continues to be a risk throughout bloom, and whenever weather is warm and wet.

Cherry leaf spot

- Cherry leaf spot management begins at early bloom for tart cherries; for sweet cherries, it begins at petal fall.
- Syllit F is labeled for use west of the Mississippi River.

Bacterial canker: For sour cherries only: Badge SC one to two times after petal fall, using lower to moderate rates. Do not apply to sweet cherry or the English Morello variety as severe injury may result.

Table 3-3. Cherry Diseases - Early Bloom Through Petal Fall And Shuck Split¹

| Product And Formulation Active Ingredient | FRAC Code ² | Brown Rot | Cherry Leaf Spot | REI ³ PHI ⁴ | Max Amt ⁵ Max App ⁶ |
|--|---------------------------|-----------------|---------------------|--------------------------------------|--|
| Abound (SC) | 11 | 12-15.5 fl. oz. | 12-15 fl. oz. | 4h | 90 fl. oz. |
| azoxystrobin | | G[r] | G[r] | 0d | 5 |
| Badge SC | M | 3.5-5 pt. | 1.5-5 pt. | 24h | 63.4 pt. |
| copper sulfate + oxychloride | | F | F-G | 0d | NA |
| Bravo Weather Stik | M5 | 3-4 pt. | 3-4 pt. | 12h | 20.5 pt. |
| chlorothalonil | | F-G | E | thru SS | NA |
| C-O-C-S WDG | M | 1-2.9 lb. | 1-2.9 lb. | 48h | 35 lb. |
| copper oxychloride | | F | F | 21d | 3 |
| Cabrio EG (20EG) | 11 | 9.5 oz. | 9.5 oz. | 12h | 47.5 oz. |
| pyraclostrobin | | F-E | G [r] | 0d | 5 |
| Captan 80WDG | M | 2.5 lb. | 2.5 lb. | 24h | 17.5 lb. |
| captan | | G | F-G | 0d | NA |
| Cevya | 3 | 3-5 fl. oz. | 3-5 fl. oz. | 12h | NA |
| mefentrifluconazole | | E | E | 0d | see label |
| Elevate 50WDG | 17 | 1-1.5 lb. | x | 12h | 6 lb. |
| fenhexamid | | G-E | x | 0d | NA |
| Elite 45DF | 3 | 4-8 fl. oz. | 4-8 oz. | 12h | 3 lb. |
| tebuconazole | | E[r] | G-E[r] | 0d | NA |
| Flint Extra | 11 | 2.5-3.8 fl. oz. | 2.5-3.8 fl. oz. | 12h | 15.2 fl. oz. |
| trifloxystrobin | | E | E | 1d | 4 |
| Fontelis (SC) | 7 | 14-20 fl. oz. | 14-20 fl. oz. | 12h | 61 fl. oz. |
| penthiopyrad | | E | F-G | 0d | NA |
| Indar 2F | 3 | 6 fl. oz. | 6 fl. oz. | 12h | 48 fl. oz. |
| fenbuconazole | | E[r] | E[r] | 0d | 8 |
| Inspire Super (EW) | 3+9 | 16-20 fl. oz. | x | 12h | 80 fl. oz. |
| difenoconazole + cyprodinil | | E | x | 2d | 4 |
| Kenja 400SC | 7 | 12.5 fl. oz. | x | 12 | 37.5 fl. oz. |
| isofetamid | | E | x | 1d | 3 |
| Luna Experience (SC) | 7+3 | 6-10 fl. oz. | 6-10 fl. oz. | 12h | 34 fl. oz. |
| fluopyram + tebuconazol | | G-E | x | 0d | NA |
| Luna Privilege | 7 | 4-6.8 fl. oz. | 6.8 fl. oz. | 12h | 13.7 fl. oz. |
| fluopyram | | G-E | s | 0d | NA |
| Luna Sensation (SC) | 7+11 | 5-5.6 fl. oz. | 5-6 fl. oz. | 12h | 27.1 fl. oz. |
| fluopyram + trifloxystrobin | | E | E-G | 1d | 4 |
| Merivon | 7+11 | 4-6.7 fl. oz. | 4-6.7 fl. oz. | 12h | 20.1 fl. oz. |
| fluxapyroxad + pyraclostrobin | | E | G-E[r] | 0d | 3 |
| Pristine | 7+11 | 10.5-14.5 oz. | 10.5-14.5 oz. | 12h | 72.5 oz. |
| pyaclostrobin + boscalid | | G | E | 0d | 5 |
| Procure 480SC | 3 | 10-16 fl. oz. | 10-16 fl. oz. | 12h | 56 fl. oz. |
| triflumizole | | G[r] | G[r] | 1d | 4 |

(Continued)

Table 3-3. Table 3-3. Cherry Diseases - Early Bloom Through Petal Fall And Shuck Split¹ (continued)

| Product And Formulation Active Ingredient | FRAC Code ² | Brown Rot | Cherry Leaf Spot | REI ³ PHI ⁴ | Max Amt ⁵ Max App ⁶ |
|--|---------------------------|---------------|---------------------|--------------------------------------|--|
| Quadris Top | 11+3 | 12-14 fl. oz. | x | 12h | 56 fl. oz. |
| azoxystrobin + difenoconazole | | E | x | 0d | NA |
| Quash | 3 | 2.5-4 oz. | 4 oz. | 12h | 10.5-12 oz. |
| metconazole | | G[r] | G[r] | 14d | 3 |
| Quilt Xcel | 11+3 | 14 fl. oz. | 14 fl. oz. | 12h | 70 fl. oz. |
| azoxystrobin + propiconazole | | E | G | 0d | 5 |
| Rally 40WSP | 3 | 2.5-6 oz. | 2.5-6 oz. | 24h | 3.2 lb. |
| myclobutanil | | E | E[r] | 0d | NA |
| Rovral 4F | 2 | 1-2 pt. | x | 24h | 4 pt. |
| iprodione | | E | x | 60d | 2 |
| Sulfur, Microthiol Disperss | M2 | 18 lb. | x | 24h | NA |
| sulfur | | F | x | NA | NA |
| Syllit F | U12 | 3 pt. | 1.5-3 pt. | 48h | 12 pt. |
| dodine | | s | G | 7d | 6 |
| Tilt (EC) | 3 | 4 fl. oz. | 4 fl. oz. | 12h | 20 fl. oz. |
| propiconazole | | G | G[r] | 0d | 5 |
| Topguard EQ | 3+11 | 6-8 oz. | x | 6-8 oz. | 12h |
| flutriafol + azoxystrobin | | G | x | E | 7d |
| Topguard Specialty Crop | 3 | 14 fl. oz. | 14 fl. oz. | 12h | 56 fl. oz. |
| flutriafol | | E | G | 7d | 4 |
| Topsin M WSB | 1 | 1-1.5 lb. | 1.5 lb. | 48h | 4 lb. |
| thiophanate-methyl | | G | F-G | 1d | NA |
| Vanguard WG (75WG) | 9 | 5 oz. | x | 12h | 30 oz. |
| cyprodinil | | G | x | 2d | 4 |
| Ziram 76DF | M3 | 5-6 lb. | 5-6 lb. | 48hr | 24.2 lb. |
| ziram | | F-i | F-i | 14d | 4 |

Footnotes: See how to read the spray schedule tables section at the beginning of the chapter.

Cherry Full Bloom - Insects

Insect management notes

- **Save the bees! Do not apply insecticides during bloom.**

Cherry Petal Fall To Shuck Split - Insects

When petals are shed from bloom through shucks split, with shucks falling from expanding fruit.

Insect management notes

- **Plum curculio:** Do not use Imidan 70W on sweet cherries.
- **European red mites:** Agri-Mek SC, Apollo SC, or Zeal miticides may be applied to suppress developing populations of European red mites.

Table 3-4. Cherry Insects - Petal Fall Through Shuck Split¹

| Product And Formulation Active Ingredient | IRAC Code ² | Black Cherry Aphid | European Red Mite | Leafroller | Plum Curculio | San Jose Scale | REI ³ PHI ⁴ | Max Amt ⁵ Max App ⁶ |
|--|------------------------|--------------------|-------------------|-------------------|-------------------|-----------------|--------------------------------------|--|
| Acramite 50WS | 20 | x | 0.75-1 lb. | x | x | x | 12h | NA |
| bifenazate | | x | E | x | x | x | 3d | 1 |
| Actara (25WDG) | 4A | 3-4 fl. oz. | x | x | 4.5-5.5 oz. | x | 12h | 11 oz. |
| thiamethoxam | | G-E | x | x | G | x | 14d | NA |
| Admire Pro (4.6F) | 4A | 1.4-2.8 fl. oz. | x | x | 2.8 fl. oz. | 1.4-2.8 fl. oz. | 12h | 10.5/14 fl. oz. |
| imidacloprid | | G-E | x | x | s | G | 7-21d | NA |
| Agri-Mek SC (0.7SC) (RUP) | 6 | x | 2.25-4.25 fl. oz. | x | x | x | 12h | 8.5 fl. oz. |
| abamectin | | x | G | x | x | x | 21d | 2 |
| Apollo SC (ISC) | 10A | x | 2-8 fl. oz. | x | x | x | 12h | NA |
| clofentezine | | x | E | x | x | x | 21d | 1 |
| Apta (1.34SC) | 21A | 17-27 fl. oz. | x | 21-27 fl. oz. | 21-27 fl. oz. | x | 12h | 53.5 fl. oz. |
| tolfenpyrad | | u | x | G | G | x | 14d | 2 |
| Asana XL (0.66EC) (RUP) | 3A | x | x | x | 4.8-14.5 fl. oz. | x | 12h | 72.5 fl. oz. |
| esfenvalerate | | x | x | x | G | x | 14d | NA |
| Assail 30SC | 4A | 2.1-4.5 fl. oz. | x | x | 4.5-6.7 fl. oz. | 4.5-6.7 fl. oz. | 12h | 26.9 fl. oz. |
| acetamiprid | | G-E | x | x | G | G | 7d | 4 |
| Avaunt eVo (30WDG) | 22A | x | x | x | 5-6 oz. | x | 12h | 24 oz. |
| indoxacarb | | x | x | x | E | x | 14d | 4 |
| Baythroid XL (1EC) (RUP) | 3A | x | x | 4.8-14.5 fl. oz. | 2.4-2.8 fl. oz. | x | 12h | 5.6 fl. oz. |
| beta-cyfluthrin | | x | x | E | G | x | 7d | NA |
| Beleaf 50SG | 29 | 2.0-2.8 fl. oz. | x | x | x | x | 12h | 8.4 fl. oz. |
| flonicamid | | G-E | x | x | x | x | 14d | 3 |
| Centaur WDG | 16 | x | x | x | x | 34.5 oz. | 12h | 69 oz. |
| buprofezin | | x | x | x | x | E | 14d | 2 |
| Danitol 2.4EC (RUP) | 3A | x | x | 10.6-21.3 fl. oz. | 10.6-21.3 fl. oz. | x | 24h | 42.7 fl. oz. |
| fenpropathrin | | x | x | G | G | x | 3d | NA |
| Delegate WG (25WG) | 5 | x | x | 4.5-7 oz. | 6-7 oz. | x | 4h | 28 oz. |
| spinetoram | | x | x | E | s | x | 7d | 4 |
| Envidor 2SC | 23 | x | 16-18 fl. oz. | x | x | x | 12h | 18 fl. oz. |
| spirodiclofen | | x | E | x | x | x | 7d | 1 |
| Entrust | 5 | x | x | 4-6 fl. oz. | x | x | 4h | 29 fl. oz. |
| spinosad | | x | x | G | x | x | 7d | 3 |
| Esteem | 7C | x | x | x | x | 4-5 fl. oz. | 12h | 15 oz. |
| pyriproxyfen | | x | x | x | x | E | 14d | 3 |
| Exirel (0.83SE) | 28 | 13.5-20.5 fl. oz. | x | 10-20.5 fl. oz. | 13.5-20.5 fl. oz. | x | 12h | 61.5 fl. oz. |
| cyantraniliprole | | G | x | E | G | x | 3d | 3 |
| Imidan 70W | 1B | x | x | 2.12-4.25 lb. | 2.12 lb. | x | 72h | 7.5 lb. |
| phosmet | | x | x | E | G | x | 7/14d | NA |

(Continued)

Table 3-4. Cherry Insects - Petal Fall Through Shuck Split¹ (continued)

| Product And Formulation Active Ingredient | IRAC Code ² | Black Cherry Aphid | European Red Mite | Leafroller | Plum Curculio | San Jose Scale | REI ³ PHI ⁴ | Max Amt ⁵ Max App ⁶ |
|--|------------------------|--------------------|-------------------|------------------|-----------------|------------------|--------------------------------------|--|
| Intrepid | 18 | x | x | 8-16 fl. oz. | x | x | 24h | 25.6 fl. oz. |
| | | x | x | G-E | x | x | 14d | NA |
| Magister SC (1.7SC) | 21A | x | 32-36 fl. oz. | x | x | x | 12h | 36 fl. oz. |
| fenazaquin | | x | u | x | x | x | 3d | 1 |
| Movento MPC | 23 | 10-14 fl. oz. | 10-14 fl. oz. | x | x | 10-14 fl. oz. | 24h | 24 fl. oz. |
| spirotetramat | | G-E | s | x | x | G-E | 7d | NA |
| Mustang Maxx (0.83EC) (RUP) | 3A | x | x | 1.2-4 fl. oz. | 1.2-4 fl. oz. | x | 12h | 24 fl. oz. |
| zeta-cypermethrin | | x | x | E | G | x | 3d | 6 |
| Nexter (75WP) | 21 | x | 4.4-10.6 oz. | x | x | x | 12h | 21.3 oz. |
| pyridaben | | x | u | x | x | x | 300d | 2 |
| Nexter SC (3.75SC) | 21A | x | 7.5-17 fl. oz. | x | x | x | 12h | NA |
| pyridaben | | x | u | x | x | x | 300d | 2 |
| Oil (superior) | UN | x | see label | see label | x | x | 4h | UN |
| mineral oil | | x | u | u | x | x | dor- mant | NA |
| Onager Optek (1EC) | 10 | x | 12-24 fl. oz. | x | x | x | 12h | 24 fl. oz. |
| hexythiazox | | x | u | x | x | x | 7d | 1 |
| Portal XLO (0.4EC) | 21A | x | 2 pts. | x | x | x | 12h | 4 pt. |
| fenpyroximate | | x | u | x | x | x | 7d | 2 |
| Pounce 25WP | 3A | x | x | x | 6.4-12.8 oz. | x | 12h | 38.4 oz. |
| permethrin | | x | x | x | G | x | 3d | 3 |
| Pyganic 5EC | 3A | 4.5-15.6 fl. oz. | x | 4.5-15.6 fl. oz. | x | 4.5-15.6 fl. oz. | 12h | NA |
| pyrethrins | | F | x | F | G | F | 0d | 10 |
| Rimon 0.83 EC | 15 | x | x | 20-50 fl. oz. | x | x | 12d | 150 fl. oz. |
| novaluron | | x | x | G-E | x | x | 8d | NA |
| Savey 50DF | 10A | x | 3-6 oz. | x | x | x | 12h | 6 oz. |
| hexythiazox | | x | E | x | x | x | 28d | 1 |
| Sevin XLR Plus (4F) | 1A | x | x | 2-3 qt. | 2-3 qt. | x | 12h | 14 qt. |
| carbaryl | | x | x | F | F | x | 3d | 3 |
| Surround WP (95WP) | UN | 25-50 lb. | x | 25-50 lb. | 25-50 lb. | x | 4h | NA |
| kaolin | | s | x | u | s | x | 0d | NA |
| Vendex 50WP (RUP) | 12B | x | 1.5-3 lb. | x | x | x | 48h | 4.5 lb. |
| fenbutatin-oxide | | x | G | x | x | x | 14d | 2 |
| Verdepryn 100SL | 28 | x | x | 5.5-11 fl. oz. | 5.5-11 fl. oz. | x | 4h | 33 fl. oz. |
| cyclaniliprole | | x | x | E | u | x | 7d | 3 |
| Warrior II (2.08CS) (RUP) | 3A | x | x | 1.2-2.5 fl. oz. | 1.2-2.5 fl. oz. | x | 24h | 12.8 fl. oz. |
| lambda-cyhalothrin | | x | x | u | G | x | 14d | NA |
| Zeal (72WP) | 10B | x | 2-3 oz. | x | x | x | 12h | 3 oz. |
| etoxazole | | x | E | x | x | x | 7d | 1 |

Footnotes: See how to read the spray schedule tables section at the beginning of the chapter.

Cherry Shuck Split To First Cover - Insects

Insect management notes

- Do not use Imidan 70W on sweet cherries.

Table 3-5. Cherry Insects - Shuck Split Through First Cover¹

| Product And Formulation Active Ingredient | IRAC Code ² | Cherry Fruit Fly | Japanese Beetle | Plum Curculio | REI ³ PHI ⁴ | Max Amt ⁵ Max App ⁶ |
|--|---------------------------|---------------------|--------------------|-------------------|--------------------------------------|--|
| Admire Pro (4.6F) | 4A | 2-2.8 fl. oz. | 1.4-2.8 fl. oz. | 2.8 fl. oz. | 12h | 10.5/14 fl. oz. |
| imidacloprid | | G | F | s | 7/21d | NA |
| Apta (1.34SC) | 21A | 14-27 fl. oz. | x | 21-27 fl. oz. | 12h | 53.5 fl. oz. |
| tolfenpyrad | | u | x | G | 14d | 2 |
| Asana XL 30SC | 3A | 4.5-6.7 fl. oz. | 4.5-6.7 fl. oz. | 4.5-6.7 fl. oz. | 12h | 26.9 fl. oz. |
| esfenvalerate | | G | x | G | 14d | NA |
| Assail 30SG | 4A | 5.3-8 oz. | 5.3-8 oz. | 5.3-8 oz. | 12h | 32 oz. |
| acetamiprid | | G | G | G | 7d | 4 |
| Avaunt eVo (30WDG) | 22 | x | x | 5-6 oz. | 12h | 24 oz. |
| indoxacarb | | x | x | E | 14d | 4 |
| Baythroid XL (1EC) (RUP) | 3A | 2.4-2.8 fl. oz. | x | 2.4-2.8 fl. oz. | 12h | 5.6 fl. oz. |
| beta-cyfluthrin | | G | x | G | 7d | NA |
| Danitol 2.4EC (RUP) | 3A | 16-21.3 fl. oz. | 10.7-21.3 fl. oz. | 10.7-21.3 fl. oz. | 24h | 42.7 fl. oz. |
| fenpropathrin | | u | E | G | 3d | NA |
| Delegate WG (25WG) | 5 | 6-7 oz. | x | 6-7 oz. | 4h | 28 oz. |
| spinetoram | | s | x | s | 7d | 4 |
| Exirel (0.83SE) | 28 | 10-17 fl. oz. | 13.5-20.5 fl. oz. | 13.5-20.5 fl. oz. | 12h | 61.5 fl. oz. |
| cyantraniliprole | | E | G | G | 3d | 3 |
| Imidan 70W | 1B | 2.1 lb. | 2.1 lb. | 2.1 lb. | 72h | 7.5 lb. |
| phosmet | | E | G | G | 7/14d | NA |
| Malathion 5EC | 1B | 2.8 pt. | 2.8 pt. | x | 12h | NA |
| malathion | | G | G | x | 3d | 4 |
| Mustang Maxx (0.83EC) (RUP) | 3A | 1.28-4 fl. oz. | x | 1.28-4 fl. oz. | 12h | 24 fl. oz. |
| zeta-cypermethrin | | F | x | G | 3d | 6 |
| Pounce 25WP | 3A | x | x | 6.4-12.8 oz. | 12h | 38.4 oz. |
| permethrin | | x | x | G | 3d | 3 |
| Pyganic 5EC | 3A | 4.5-15.6 fl. oz. | 4.5-15.6 fl. oz. | x | 12h | NA |
| pyrethrins | | i | F | x | 0d | 10 |
| Sevin XLR Plus (4F) | 1A | 2-3 qt. | 2-3 qt. | 2-3 qt. | 12h | 14 qt. |
| carbaryl | | G | E | F | 3d | 3 |
| Surround WP (95WP) | UN | 25-50 lb. | 25-50 lb. | 25-50 lb. | 4h | NA |
| kaolin | | s | F | s | 0d | NA |
| Verdepryn 100SL | 28 | 5.5-11 fl. oz. | 5.5-11 fl. oz. | 5.5-11 fl. oz. | 4h | 33 fl. oz. |
| cyclaniliprole | | G | u | u | 7d | 3 |
| Warrior II (2.08CS) (RUP) | 3A | 1.2-2.5 fl. oz. | 1.2-2.5 fl. oz. | 1.2-2.5 fl. oz. | 24h | 12.8 fl. oz. |
| lambda-cyhalothrin | | G | E | G | 14d | NA |

Footnotes: See how to read the spray schedule tables section at the beginning of the chapter.

Cherry Summer Cover To Harvest Sprays - Diseases

Applications begin 10-14 days after shuck split as needed through harvest.

Notes on disease management

- **Leaf spot:** Syllit is labeled for use west of the Mississippi River.

Table 3-6. Cherry Diseases - Summer Cover Through Harvest¹

| Product And Formulation Active Ingredient | FRAC Code ² | Brown Rot | Cherry Leaf Spot | REI ³ PHI ⁴ | Max Amt ⁵ Max App ⁶ |
|--|------------------------|-----------------|------------------|--------------------------------------|--|
| Abound (SC) | 11 | 12-15.5 fl. oz. | 12-15 fl. oz. | 4h | 90 fl. oz. |
| azoxystrobin | | G[r] | G[r] | 0d | 5 |
| Badge SC | M | 3.5-5 pt. | 1.5-5 pt. | 24h | 63.4 pt. |
| copper sulfate + oxychloride | | F | F-G | 0d | NA |
| C-O-C-S WDG | M | 1-2.9 lb. | 1-2.9 lb. | 48h | 35 lb. |
| copper oxychloride | | F | F | 21d | 3 |
| Cabrio EG (20EG) | 11 | 9.5 oz. | 9.5 oz. | 12h | 47.5 oz. |
| pyraclostrobin | | F-E | G[r] | 0d | 5 |
| Captan 80WDG | M | 2.5 lb. | 2.5 lb. | 24h | 17.5 lb. |
| captan | | G | F-G | 0d | NA |
| CaptEvote 68WDG | M+17 | 3.7 lb. | 3.75 lb. | 24h | 18.7 lb. |
| captan + fenhexamid | | E | G | 0d | NA |
| Cevya | 3 | 3-5 fl. oz. | 3-5 fl. oz. | 12h | NA |
| mefentrifluconazole | | E | E | 0d | see label |
| Elevate 50WDG | 17 | 1-1.5 lb. | x | 12h | 6 lb. |
| fenhexamid | | G-E | x | 0d | NA |
| Elite 45DF | 3 | 4-8 fl. oz. | 4-8 oz. | 12h | 3 lb. |
| tebuconazole | | E [r] | E-G[r] | 0d | NA |
| Flint Extra | 11 | 2.5-3.8 fl. oz. | 2.5-3.8 fl. oz. | 12h | 15.2 oz. |
| trifloxystrobin | | E | E | 1d | 4 |
| Fontelis (SC) | 7 | 14-20 fl. oz. | 14-20 fl. oz. | 12h | 61 fl. oz. |
| penthiopyrad | | E | F-G | 0d | NA |
| Indar 2F | 3 | 6 fl. oz. | 6 fl. oz. | 12h | 48 fl. oz. |
| fenbuconazole | | E[r] | E[r] | 0d | 8 |
| Inspire Super (EW) | 3+9 | 16-20 fl. oz. | x | 12h | 80 fl. oz. |
| difenoconazole + cyprodinil | | E | x | 2d | 4 |
| Kenja 400SC | 7 | 12.5 fl. oz. | x | 12 | 37.5 fl. oz. |
| isofetamid | | E | x | 1d | 3 |
| Luna Experience (SC) | 7+3 | 6-10 fl. oz. | 6-10 fl. oz. | 12h | 34 fl. oz. |
| fluopyram + tebuconazol | | G-E | x | 0d | NA |
| Luna Privilege | 7 | 4-6.8 fl. oz. | 6.8 fl. oz. | 12h | 13.7 fl. oz. |
| fluopyram | | G-E | s | 0d | NA |
| Luna Sensation (SC) | 7+11 | 5-5.6 fl. oz. | 5-6 fl. oz. | 12h | 27.1 fl. oz. |
| fluopyram + trifloxystrobin | | E | E-G | 1d | 4 |

(Continued)

Table 3-6. Cherry Diseases - Summer Cover Through Harvest¹ (continued)

| Product And Formulation Active Ingredient | FRAC Code ² | Brown Rot | Cherry Leaf Spot | REI ³ PHI ⁴ | Max Amt ⁵ Max App ⁶ |
|--|---------------------------|---------------|---------------------|--------------------------------------|--|
| Merivon XBF | 7+11 | 4-6.7 fl. oz. | 4-6.7 fl. oz. | 12h | 20.1 fl. oz. |
| fluxapyroxad + pyraclostrobin | | E | E-G | 0d | 3 |
| Pristine | 7+11 | 10.5-14.5 oz. | 10.5-14.5 oz. | 12h | 72.5 oz. |
| pyaclostrobin + boscalid | | G | E | 0d | 5 |
| Procure 480SC | 3 | 10-16 fl. oz. | 10-16 fl. oz. | 12h | 56 fl. oz. |
| triflumizole | | G[r] | G[r] | 1d | 4 |
| Quadris Top | 11+3 | 12-14 fl. oz. | x | 12h | 56 fl. oz. |
| azoxystrobin + difenoconazole | | E | x | 0d | NA |
| Quash | 3 | 2.5-4 oz. | 4 oz. | 12h | 10.5-12 oz. |
| metconazole | | G[r] | G[r] | 14d | 3 |
| Quilt Xcel | 11+3 | 14 fl. oz. | x | 12h | 70 fl. oz. |
| azoxystrobin + propiconazole | | E | G | 0d | 5 |
| Rally 40WSP | 3 | 2.5-6 oz. | 2.5-6 oz. | 24h | 3.2 lb. |
| myclobutanil | | E | E[r] | 0d | NA |
| Sulfur, Microthiol Disperss | M2 | 18 lb. | x | 24h | NA |
| sulfur | | F | x | NA | NA |
| Syllit F | U12 | 3 pt. | 1.5-3 pt. | 48h | 12 pt. |
| dodine | | s | G | 7d | 6 |
| Tilt (EC) | 3 | 4 fl. oz. | 4 fl. oz. | 12h | 20 fl. oz. |
| propiconazole | | G | G[r] | 0d | 5 |
| Topguard EQ | 3+11 | 6-8 oz. | x | 6-8 oz. | 12h |
| flutriafol + azoxystrobin | | G | x | E | 7d |
| Topguard Specialty Crop | 3 | 14 fl. oz. | 14 fl. oz. | 12h | 56 fl. oz. |
| flutriafol | | E | G | 7d | 4 |
| Topsin M WSB | 1 | 1-1.5 lb. | 1.5 lb. | 48h | 4 lb. |
| thiophanate-methyl | | G | F-G | 1d | NA |
| Vanguard WG (75WG) | 9 | 5 oz. | x | 12h | 30 oz. |
| cyprodinil | | G | x | 2d | 4 |
| Ziram 76DF | M3 | 5-6 lb. | x | 48hr | 24.2 lb. |
| ziram | | F | x | 14d | 4 |

Footnotes: See how to read the spray schedule tables section at the beginning of the chapter.

Additional Summer Cover Sprays To Harvest - Insects

Insect management notes

- Imidan 70W: Do not use on sweet cherries.
- **Lesser peachtree borer:** Control of the first generation of lesser peachtree borer with trunk sprays

is during the time of peak moth flight, generally from early May to early June, depending on latitude and spring temperatures. See Borers of Peach, Cherry, and Plum Trees, page 152.

- **Spotted-wing Drosophila:** Can begin to attack fruit when they change color and soften before harvest.

Table 3-7. Cherry Insects - Summer Cover Through Harvest¹

| Product And Formulation Active Ingredient | IRAC Code ² | Cherry Fruit Fly | Japanese Beetle | Borers | Spotted-Wing Drosophila | REI ³ PHI ⁴ | Max Amt ⁵ Max App ⁶ |
|--|---------------------------|-------------------------------|--------------------|------------------|----------------------------|--------------------------------------|--|
| Actara (25WDG) | 4A | 4.5-5.5 oz. | x | x | x | 12h | 11 oz. |
| thiamethoxam | | G | x | x | x | 14d | NA |
| Admire Pro (4.6F) | 4A | 2-2.8 fl. oz. | 1.4-2.8 fl. oz. | x | x | 12h | 10.5/14 fl. oz. |
| imidacloprid | | G | F | x | x | 7/21d | NA |
| Altacor eVo (35WDG) | 28 | 1.5-2.2 fl. oz. | x | x | x | 4h | 4.6 fl. oz. |
| chlorantraniliprole | | s | x | x | x | 10d | 3 |
| Apta (1.34SC) | 21A | 14-27 fl. oz. | x | x | 21-27 fl. oz. | 12h | 53.5 fl. oz. |
| tolfenpyrad | | u | x | x | s | 14d | 2 |
| Asana XL (0.66EC) (RUP) | 3A | 4.8-14.5 fl. oz. | x | 4.8-14.5 fl. oz. | see label | 12h | 72.5 fl. oz. |
| esfenvalerate | | G | x | G | x | 14d | NA |
| Assail 30SG | 4A | 4.5-6.7 fl. oz. | 4.5-6.7 fl. oz. | x | x | 12h | 26.9 fl. oz. |
| acetamiprid | | G | G | x | x | 7d | 4 |
| Baythroid XL (1EC) (RUP) | 3A | 2.4-2.8 fl. oz. | x | 1.4-2 fl. oz. | see label | 12h | 5.6 fl. oz. |
| beta-cyfluthrin | | G | x | G | G | 7d | NA |
| Danitol 2.4EC (RUP) | 3A | 16-21.3 fl. oz. | 10.6-21.3 fl. oz. | x | 10.6-21.3 fl. oz. | 24h | 42.7 fl. oz. |
| fenpropathrin | | u | E | x | E | 3d | NA |
| Delegate WG (25WG) | 5 | 6-7 fl. oz. | x | x | 4.5-7 oz. | 4h | 28 oz. |
| spinetoram | | s | x | x | G | 7d | 4 |
| Diazinon AG 600WBC (RUP) | 1B | 6.5-12.7 fl. oz./ 100 gal. | x | x | x | 96h | 102 fl. oz. |
| diazinon | | G | x | x | x | 21d | 2 |
| Entrust SC (2SC) | 5 | 4-8 fl. oz. | x | x | 4-8 fl. oz. | 4h | 29 fl. oz. |
| spinosad | | F | x | x | G-E | 7d | 3 |
| Exirel (0.83SE) | 28 | 10-17 fl. oz. | 13.5-20.5 fl. oz. | x | 13.5-20.5 fl. oz. | 12h | 61.5 fl. oz. |
| cyantraniliprole | | E | G | x | E | 3d | 3 |
| Imidan 70W | 1B | 2.12 lb. | 2.12 lb. | x | 2.1 lb. | 72h | 7.5 lb. |
| phosmet | | E | G | x | E | 7/14d | NA |
| Malathion 5EC | 1B | 2.8 pt. | 2.8 pt. | x | see label | 12h | NA |
| malathion | | G | G | x | G | 3d | 4 |
| Movento MPC | 23 | 10-14 fl. oz. | x | x | 10-14 fl. oz. | 24h | 24 fl. oz. |
| spirotetramat | | u | x | x | G | 7d | NA |
| Mustang Maxx (0.83EC) (RUP) | 3A | 1.28-4 fl. oz. | x | 1.28-4 fl. oz. | 4 fl. oz. | 12h | 24 fl. oz. |
| zeta-cypermethrin | | F | x | G | E | 3d | 6 |
| Pounce 25WP | 3A | x | x | 6.4-12.8 fl. oz. | see label | 12h | 38.4 oz. |
| permethrin | | x | x | F | u | 3d | 3 |
| Pyganic 5EC | 3A | 4.5-15.6 fl. oz. | 4.5-15.6 fl. oz. | x | 4.5-15.6 fl. oz. | 12h | NA |
| pyrethrins | | i | F | x | F | 0d | 10 |
| Rimon 0.83EC | 15 | 20-40 fl. oz. | x | 20 fl. oz. | 20-40 fl. oz. | 12h | 150 fl. oz. |
| novaluron | | u | x | u | G | 8d | NA |

(Continued)

Table 3-7. Cherry Insects - Summer Cover Through Harvest¹ (continued)

| Product And Formulation Active Ingredient | IRAC Code ² | Cherry Fruit Fly | Japanese Beetle | Borers | Spotted-Wing Drosophila | REI ³ PHI ⁴ | Max Amt ⁵ Max App ⁶ |
|--|---------------------------|---------------------|--------------------|----------------------|----------------------------|--------------------------------------|--|
| Sevin XLR Plus (4F) | 1A | 2-3 qt. | 2-3 qt. | 2-3 qt. | x | 12h | 14 qt. |
| carbaryl | | G | E | u | x | 3d | 3 |
| Surround WP (95WP) | UN | 25-50 lb. | 25-50 lb. | x | x | 4h | NA |
| kaolin | | s | F | x | x | 0d | NA |
| Verdepryn 100SL | 28 | 5.5-11 fl. oz. | 5.5-11 fl. oz. | 5.5-11 fl. oz. | 5.5-11 fl. oz. | 4h | 33 fl. oz. |
| cyclaniliprole | | G | u | u | u | 7d | 3 |
| Warrior II (2.08CS) (RUP) | 3A | 1.2-2.5 fl. oz. | 1.2-2.5 fl. oz. | 1.28-2.56 fl. oz. | see label | 24h | 12.8 fl. oz. |
| lambda-cyhalothrin | | G | E | G | E | 14d | NA |

Footnotes: See how to read the spray schedule tables section at the beginning of the chapter.

Cherry Postharvest - Disease

Disease management notes

- The goal is to keep trees from defoliating without driving fungicide resistance. Rely on Bravo, coppers and sulfur for disease management at this time.

Table 3-8. Cherry Diseases - Postharvest¹

| Product And Formulation Active Ingredient | FRAC Code ² | Cherry Leaf Spot | Powdery Mildew | REI ³ PHI ⁴ | Max Amt ⁵ Max app ⁶ |
|--|---------------------------|---------------------|-------------------|--------------------------------------|--|
| Badge SC | M | 1.5-5 pt. | x | 24h | 63.4 pt. |
| copper sulfate + oxychloride | | F-G | x | 0d | NA |
| Bravo Weather Stik | M5 | 3-4 pt. | x | 12h | 20.5 pt. |
| chlorothalonil | | E | x | through shuck split | NA |
| C-O-C-S WDG | M | 1-2.9 lb. | x | 48h | 35 lb. |
| copper oxychloride | | F | x | 21d | 3 |
| Captan 80WDG | M | 2.5 lb. | x | 24h | 17.5 lb. |
| captan | | F-G | x | 0d | NA |
| Cuprofix Ultra 40 Disperss | M | 3.75 lb. | x | 48h | 45 lb. |
| copper sulfate | | P-G | x | 120d | NA |
| Sulfur, Microthiol Disperss | M2 | x | 10-30 lb. | 24h | NA |
| sulfur | | x | G | NA | NA |
| Syllit F | U12 | 1.5-3 pt. | x | 48h | 12 pt. |
| dodine | | G | x | 7d | 6 |
| Ziram 76DF | M3 | x | x | 48hr | 24.2 lb. |
| ziram | | F-i | x | 14d | 4 |

Footnotes: See how to read the spray schedule tables section at the beginning of the chapter.

Cherry Postharvest - Insects (see Summer Covers)

Insect management notes

Japanese beetle

- Damage by adult beetles feeding on leaves is sporadic and transient in July to August in most of the region.
- Leaves are skeletonized and may turn brown and fall. Defoliation is not fatal but may be stressful.

Special Comments on Cherry Schedule

Pro-Gibb on cherries

The active ingredient in Pro-Gibb is a natural plant hormone, gibberellin A3. You can use it to maintain and extend high fruiting capacity of bearing tart cherry trees and to reduce the occurrence of "blind" nodes by stimulating lateral vegetative buds and a more productive balance of lateral shoots and spurs.

Apply 4 to 18 fl. oz. of Pro-Gibb 4% per acre from 14 to 28 days after bloom when 1-3 inches of terminal shoot extension has occurred in sufficient water to provide uniform coverage. Application rate depends on tree age and vigor. See label.

Since Pro-Gibb acts on buds that will flower the following growing year, responses will not begin to be visible until the year after application. Shoot, spur and flower changes will be evident two or three years after the program is started.

Applications must be applied annually to promote spur development and yield improvement.

Prohexidione-calcium (Pro-Ca) for vegetative control

Pro-Ca products are labeled for vegetative control on sweet cherry. These include Apogee, Cryova, Kudos, and generics. Refer to labels for registration and use information.

Fungicides for Phytophthora root rot management

See page 133 for Phytophthora Management on Bearing and Non-bearing Stone Fruit.

4. PEACH

Peach Insect Pests

The shaded boxes represent the crop stages where common pests in the Midwest are active; scouting and preventative sprays may be necessary/recommended. Weather and degree day accumulation will impact the exact timing of pest appearance in the orchard. **MD= mating disruption and pheromone traps.**

| Dormant | Pink | Full Bloom | Petal Fall | Shuck Split | First Cover | Second Cover | Additional Covers | Pre-Harvest |
|---------------------|------------|------------|---------------------|-------------------------|-----------------|--------------|-------------------|-------------------------|
| | | | | | | | Green June Beetle | |
| | | | | | | | Japanese Beetle | |
| | PTB MD | | | | Peachtree Borer | | | Peachtree Borer |
| | FM MD | | Oriental Fruit Moth | | | | | |
| | | | Plum Curculio | | | | | |
| | Stink Bugs | | Stink Bugs | | | | | |
| San Jose Scale | | | | | San Jose Scale | | | |
| | | | | European Red Mite | | | | |
| | | | | Green Peach Aphid | | | | |
| | | | | | | | | Spotted-Wing Drosophila |
| Tarnished Plant Bug | | | Tarnished Plant Bug | | | | | |
| | | | | Two-Spotted Spider Mite | | | | |
| | | | | | | | | |

Peach Spray Schedule

How to read the spray schedule tables

Every peach growth stage has important notes on disease or insect management. In some cases, the reader will be directed to the special problems section at the end of the section or chapter. Please make sure to read thoroughly and contact your state Extension specialist with any specific questions.

Key to tables

E = excellent control

G = good control

F = fair control

[r] = fungicide/insecticide resistance possible

s = suppression only

i = ineffective

u = unknown efficacy

x = pest not on the label

¹Efficacy data in this publication are based on trials conducted across various regions and does not necessarily reflect local efficacy differences or changes over time. Growers should contact their Extension specialist for the most recent or for state-specific information. The information on this publication is only a guide; the authors and their institutions assume no liability for practices implemented based on this information. Always read and follow pesticide labels. The label is the law. Product registration may vary by state.

² FRAC/IRAC code represents the mode of action of the fungicide/insecticide.

³ PHI refers to the pre-harvest interval, which is the number of days before harvest that the product may not be applied.

⁴ All fungicides/insecticides have a Restricted-Entry Interval (REI). The restricted-entry interval is the time immediately after a pesticide application when entry

into the treated area is limited. Check labels for REI. Restrictions in REI may prohibit the use of certain pesticides during harvest.

Applicators must abide by both maximum amount of product per season AND maximum number of applications.

⁵ Max amt refers to the product's maximum amount/acre/year.

⁶ Max app refers to the product's maximum number of applications per year.

RUP refers to restricted use pesticide.

Peach Dormant To Delayed Dormant - Diseases

After leaves drop in the fall or as buds swell in spring.

Notes on disease management

Peach leaf curl

- To effectively control peach leaf curl, fungicides must be applied before bud swell. Best control is

achieved by applying in late autumn at 50% leaf fall and again at delayed dormant in late winter before floral buds begin to open; second best time is in early spring prior to bud break.

Bacterial diseases (bacterial canker and shot hole)

- Copper pesticides: Using copper at the dormant stage may reduce the overwintering inoculum of the bacteria that cause bacterial infection. As season progresses, reduce the rate of copper applied to reduce the risk of phytotoxicity. Copper applied using tank mixes with a pH of less than 6.5 may result in phytotoxicity issues.
- Be sure to incorporate oxytetracycline compounds to reduce the risk of bacterial pathogens evolving resistance to copper products.
- The addition of 1 to 3 pounds of hydrated lime per copper application may reduce crop injury.

Table 4-1. Peach Diseases - Dormant Through Delayed Dormant¹

| Product And Formulation Active Ingredient | FRAC Code ² | Bacterial Canker | Bacterial Spot Dormant | Peach Leaf Curl | Phytophthora Crown, Collar And Root Rot | REI ³ PHI ⁴ | Max Amt ⁵ Max App ⁶ |
|--|---------------------------|---------------------|------------------------------|--------------------|---|--------------------------------------|--|
| Badge SC | M | 3.5-14 pt. | 3.5-14 pt. | 3.5-14 pt. | x | 24h | 63.4 pt. |
| copper sulfate + oxychloride | | x | F-G | F-G | x | 0d | NA |
| Bravo Weather Stik | M3 | x | x | 3.1-4.1 pt. | x | 12h | 20.5 pt. |
| chlorothalonil | | x | x | G | x | shuck-split | NA |
| C-O-C-S WDG | M | 12-15.6 lb. | 12-15.6 lb. | 1-2.9 lb. | x | 48h | 35 lb. |
| copper oxychloride | | G | F-G | F-G | x | 21d | 3 |
| Cuprofix Ultra 40 disperss | M | 5-7.5 lb. | 5-7.5 lb. | 5-7.5 lb. | x | 12h | 45 lb. |
| copper hydroxide | | G | F-G | F-G | x | 120d | NA |
| Ferbam Granuflo | M | x | x | 4.5 lb. | x | 24h | 3.4 lb. |
| ferbam | | x | x | G-E | x | 21d | 3 |
| Kocide 3000 | M | 3.5-7 lb. | 3.5-7 lb. | 3.5-7 lb. | x | 48h | 60 lb. |
| copper hydroxide | | G | F-G | F-G | x | 0d | NA |
| Ridomil Gold SL | 4 | x | x | x | 2 qt./A or 1.5 oz. per 1000 sq. ft. | 48 | 1.5 gal. |
| mefenoxam | | x | x | x | E | NA | 3 |
| Thiram Granuflo | M3 | x | x | 3.5 lb. | x | 24h | 21.2 lb. |
| thiram | | x | x | G | x | 7d | NA |
| Ziram 76DF | M3 | x | x | 3.75-8 lb. | x | 48h | 48.2 lb. |
| ziram | | x | x | E | x | 30d | 6 |

Footnotes: See how to read the spray schedule tables section at the beginning of the chapter.

Peach Dormant - Insects

After leaves drop in the fall or before buds swell in spring.

Notes on insect management

- **Mites and San Jose scale:** When spraying superior oil, check labels for temperature restrictions (e.g., only when temperatures are above 40F° or never during freezing weather).

Table 4-2. Peach Insects - Dormant¹

| Product And Formulation | Active Ingredient | IRAC Code ² | San Jose Scale | REI ³ PHI ⁴ | Max Amt ⁵ Max App ⁶ |
|--------------------------|-------------------|------------------------|------------------------|--------------------------------------|--|
| Assail 30SG | | 4A | 5.3-8 oz. | 12h | 32 oz. |
| | acetamiprid | | F | 7d | 4 |
| Belay (2.13SC) | | 4A | 6 fl. oz. | 12h | 12 fl. oz. |
| | clothianidin | | G | 21d | NA |
| Centaur WDG (70WDG) | | 16 | 34.5 oz. | 12h | 69 oz. |
| | buprofezin | | E | 14d | 2 |
| Diazinon AG 600WBC (RUP) | | 1B | 12.75 fl. oz./100 gal. | 4d | 51 fl. oz. |
| | diazinon | | F | 21d | 2 |
| Esteem 35WP | | 7C | 4-5 oz. | 12h | 15 oz. |
| | pyriproxifen | | E | 14d | 3 |
| Imidan 70W | | 1B | 2.1-4.25 lb. | 4d | 17 lb. |
| | phosmet | | E | 14d | NA |
| Neemix 4.5 (0.39L) | | UN | 7-16 fl. oz. | 4h | NA |
| | azadirachtin | | G | 0d | NA |
| Damoil | | UN | 0.25-0.67% | 4h | NA |
| | mineral oil | | G | NA | NA |
| Pyganic 5EC | | 3A | 4.5-15.6 fl. oz. | 12h | 15.6 fl. oz. |
| | pyrethrins | | u | 0d | 1 |
| Sevin XLR Plus (4F) | | 1A | 4-5 qt. | 12h | 14 qt. |
| | carbaryl | | u | 3d | 3 |
| Sivanto Prime | | 4D | 10.5-14 fl. oz. | 4h | 28 fl. oz. |
| | flupyradifurone | | G | 14d | NA |

Footnotes: See how to read the spray schedule tables section at the beginning of the chapter.

Peach Pink - Diseases

Notes on disease management

Copper pesticides: Copper rates are tied to crop development, with rates reduced as the season progresses to minimize the risk of phytotoxicity. When using coppers post-bloom, do not apply during extended dews or foggy conditions. Spotting of leaves and defoliation may occur from copper use after bud swell. Generally, 2-2.5 lb. cu/A at early bud break; 1-2 lb. cu/A at pink bud; 1 lb. cu/A at blossoms opening; 0.5 lb. cu/A at petal fall; and 0.10-0.25 lb. cu/A + Mycoshield at shuck-split. The recommended amount is copper (not the commercial compound), so depends on the formulation amount of commercial compound

would be different. The addition of 1 to 3 pounds of hydrated lime at blossom opening may reduce crop injury.

- Under severe disease pressure, use the Ziram 76DF higher rate.
- Fungicides with the FRAC code 3, 7, or 11 are not recommended at this time, unless disease pressures are particularly high. They are better deployed later in the season.
- Using Bravo Weather Stik (BWS) plus copper plus Fontelis alternated at 7-day intervals with BWS plus copper plus Inspire Super from pink to full bloom stages has been reported very effective against diseases during this period.

Table 4-3. Peach Diseases - Pink¹

| Product And Formulation Active Ingredient | FRAC Code ² | Bacterial Spot/Shot Hole | Brown Rot | Peach Leaf Curl | REI ³ PHI ⁴ | Max Amt ⁵ Max App ⁶ |
|--|---------------------------|--------------------------------|-----------------|--------------------|--------------------------------------|--|
| Abound (SC) | 11 | x | 12-15.5 fl. oz. | x | 4h | 90 fl. oz. |
| azoxystrobin | | x | F-E[r] | x | 0d | See label |
| Badge SC | M | 0.5-2 pt. | 3.5-5.25 pt. | 3.5-5.25 pt. | 24h | 63.4 pt. |
| copper sulfate + oxychloride | | F-G | F | F-G | 0d | NA |
| Bravo Weather Stik | M3 | x | 3.1-4.1 pt. | 3.1-4.1 pt. | 12h | 20.5 pt. |
| chlorothalonil | | x | G | G | shuck-split | NA |
| Captan | M4 | x | 2.5-5 lb. | x | 24h | 40 lb. |
| captan | | x | G | x | 0d | NA |
| Cevya | 3 | x | 3-5 fl. oz. | x | 12h | 15 fl. oz. |
| mefentrifluconazole | | x | E | x | 0d | See label |
| C-O-C-S WDG | M | 1-2.9 lb. | 1-2.9 lb. | 1-2.9 lb. | 48h | 35 lb. |
| copper oxychloride | | F-G | F | F-G | 21d | 3 |
| Cuprofix Ultra 40 disperss | M | 1.5 lb. | 1.5 lb. | 1.5 lb. | 12h | 45 lb. |
| copper hydroxide | | F-G | F | F-G | 120d | NA |
| Elevate 50 WDG | 17 | x | 1-1.5 lb. | x | 12h | 6 lb. |
| fenhexamid | | x | G-E | x | 0d | NA |
| Elite 45DF | 3 | x | 4-8 oz. | x | 12h | 3 lb. |
| tebuconazole | | x | E | x | 0d | NA |
| Ferbam Granufflo | M | x | x | 4.5 lb. | 24h | 3.4 lb. |
| ferbam | | x | x | G-E | 21d | 3 |
| Flint Extra | 11 | x | 2.5-3.8 fl. oz. | x | 12h | 15.2 fl. oz. |
| trifloxystrobin | | x | s (G) | E | 1d | 4 |
| Fontelis (SC) | 7 | x | 14-20 fl. oz. | x | 12h | 61 fl. oz. |
| penthiopyrad | | x | G-E | x | 0d | NA |
| Indar 2F | 3 | x | 6 fl. oz. | x | 12h | 48 fl. oz. |
| fenbuconazole | | x | E[r] | x | 0d | 8 |
| Inspire Super (EW) | 3+9 | x | 16-20 fl. oz. | x | 12h | 80 fl. oz. |
| difenoconazole + cyprodinil | | x | E | x | 2d | 4 |
| Kenja 400 SC | 7 | x | 12.5 fl. oz. | x | 12h | 37.5 fl. oz. |
| isofetamid | | x | E | x | 1d | 3 |
| Kocide 3000 | M | 3.5-5 lb. | 3-5 lb. | 3.5-5 lb. | 48h | 60 lb. |
| copper hydroxide | | F-G | F | F-G | 0d | NA |
| Luna Experience (SC) | 7+3 | x | 6-10 fl. oz. | 6-10 fl. oz. | 12h | 34 fl. oz. |
| fluopyram + tebuconazol | | x | G-E | G-E | 0d | NA |
| Luna Privilege | 7 | x | 4-6.8 fl. oz. | x | 12h | 13.7 fl. oz. |
| fluopyram | | x | E | x | 0d | NA |
| Luna Sensation (SC) | 7+11 | x | 5-7.6 fl. oz. | 5-7.6 fl. oz. | 12h | 27.1 fl. oz. |
| fluopyram + trifloxystrobin | | x | E | G-E | 1d | 4 |
| Merivon XBF | 7+11 | x | 4-6.7 fl. oz. | x | 12h | 20.1 fl. oz. |
| fluxapyroxad + pyraclostrobin | | x | E | x | 0d | 3 |

(Continued)

Table 4-3. Peach Diseases - Pink¹ (continued)

| Product And Formulation Active Ingredient | FRAC Code ² | Bacterial Spot/Shot Hole | Brown Rot | Peach Leaf Curl | REI ³ PHI ⁴ | Max Amt ⁵ Max App ⁶ |
|--|---------------------------|--------------------------------|-----------------|--------------------|--------------------------------------|--|
| Microthiol Disperss | M | x | 10-20 lb. | 10-20 lb. | 24h | NA |
| sulfur | | x | F-P | G-E | 0d | NA |
| Miravis | 7 | x | 3.4-5.1 fl. oz. | x | 4h | 20.4 fl. oz. |
| pydiflumetofen | | x | E | x | 0d | 4 |
| Ph-D | 19 | x | x | x | 4h | NA |
| polyoxin D | | x | x | x | 0h | NA |
| Pristine | 7+11 | x | 10.5-14.5 oz. | 10.5-14.5 oz. | 12h | 72.5 oz. |
| pyaclostrobin + boscalid | | x | G[r] | s(G) | 0d | 5 |
| Quadris Top | 11+3 | x | 12-14 fl. oz. | x | 12h | 56 fl. oz. |
| azoxystrobin + difenoconazole | | x | E | x | 0d | 4 |
| Quash | 3 | x | 2.5-3.5 fl. oz. | x | 12h | 12 fl. oz. |
| metconazole | | x | G-E | x | 14d | 3 |
| Quilt Xcel | 11+3 | x | 14 fl. oz. | x | 12h | 70 fl. oz. |
| azoxystrobin + propiconazole | | x | E | x | 0d | 5 |
| Rally 40WSP | 3 | x | 2.5-6 oz. | x | 24h | 3.25 lb. |
| myclobutanil | | x | G | x | 0h | NA |
| Rovral 4F | 2 | x | 1-2 pt. | x | 24h | 4 pt. |
| iprodione | | x | E | x | PF | 2 |
| Scala (SC) | 9 | x | 9-18 fl. oz. | x | 12h | 54 fl. oz. |
| pyrimethanil | | x | G-E | x | 2d | 3 |
| Syllit F | U12 | x | 3 pt. | 3 pt. | 48h | 9 pt. |
| dodine | | x | s | E | petal fall | 3 |
| Thiram Granuflo | M3 | x | 3.5 lb. | 3.5 lb. | 24h | 21.2 lb. |
| thiram | | x | G | G | 7d | NA |
| Tilt (EC) | 3 | x | 4 fl. oz. | x | 12h | 20 fl. oz. |
| propiconazole | | x | E | x | 0d | 5 |
| Topguard EQ | 3+11 | x | 6-8 fl. oz. | x | 12h | NA |
| flutriafol + azoxystrobin | | x | G | x | 7d | 4 |
| Topguard Specialty Crop | 3 | x | 14 fl. oz. | 14 fl. oz. | 12h | 56 fl. oz. |
| flutriafol | | x | E | G | 7d | 4 |
| Topsin M WSB | 1 | x | 1-1.5 lb. | x | 48h | 4 lb. |
| thiophanate-methyl | | x | E[r] | x | 1d | NA |
| Vanguard WG (75WG) | 9 | x | 5 oz. | x | 12h | 30 oz. |
| cyprodinil | | x | G-E | x | 2d | 4 |
| Ziram 76DF | M3 | x | 4.5-8 lb. | 3.75-8 lb. | 48h | 48.2 lb. |
| ziram | | x | G | E | 30d | 6 |

Footnotes: See how to read the spray schedule tables section at the beginning of the chapter.

Peach Pink - Insects

Notes on insect management

- Oriental fruit moth monitoring:** Put pheromone traps to monitor Oriental fruit moth in place now to determine the need for sprays at petal fall. **For mating disruption, see Mating Disruption for Peach Pests, page 153.**
- Peachtree borers:** Pheromone traps should be deployed pre bloom. For mating disruption, see Mating Disruption for Peach Pests page 153.
- Tarnished plant bug, stink bugs:** Apply insecticides before any blooms open.

Table 4-4. Peach Insects - Pink¹

| Product And Formulation | Active Ingredient | IRAC Code ² | Plant Bug/Stink Bug | REI ³ PHI ⁴ | Max Amt ⁵ Max App ⁶ |
|-----------------------------|--------------------|------------------------|---------------------|--------------------------------------|--|
| Asana XL (0.66EC) (RUP) | | 3A | 4.8-14.5 fl. oz. | 12h | 72.5 fl. oz. |
| | esfenvalerate | | G | 14d | NA |
| Assail 30SG | | 4A | 5.3-8 oz. | 12h | 32 oz. |
| | acetamiprid | | F | 7d | 4 |
| Baythroid XL (1EC) (RUP) | | 3A | 2-2.4 fl. oz. | 12h | 5.6 fl. oz. |
| | beta-cyfluthrin | | E | 7d | NA |
| Belay (2.13SC) | | 4A | 6 fl. oz. | 12h | 12 fl. oz. |
| | clothianidin | | E | 21d | NA |
| Beleaf 50SG | | 29 | 2-2.8 oz. | 12h | 8.4 oz. |
| | flonicamid | | G | 14d | 3 |
| Danitol 2.4EC (RUP) | | 3A | 10.7-21.3 fl. oz. | 24h | 42.7 fl. oz. |
| | fenpropathrin | | E | 3d | NA |
| Lannate LV | | 1A | 3 pt. | 4d | 18 pt. |
| | methomyl | | G | 4d | 6 |
| Mustang Maxx (0.83EC) (RUP) | | 3A | 1.28-4 fl. oz. | 12h | 24 fl. oz. |
| | zeta-cypermethrin | | E | 14d | NA |
| Neemix 4.5 (0.39L) | | UN | 7-16 fl. oz. | 4h | NA |
| | azadirachtin | | E | 0d | NA |
| Pyganic 5EC | | 3A | 4.5-15.6 fl. oz. | 12h | 15.6 fl. oz. |
| | pyrethrins | | u | 0d | 1 |
| Rimon 0.83EC | | 15 | 20-40 fl. oz. | 12h | 150 fl. oz. |
| | novaluron | | u | 8d | NA |
| Scorpion 35SL | | 4A | 5.25-7 fl. oz. | 12h | 14.25 fl. oz. |
| | dinotefuran | | E | 3/21d | NA |
| Sevin XLR Plus (4F) | | 1A | 2-3 qt. | 12h | 14 qt. |
| | carbaryl | | F | 3d | 3 |
| Venom (70SG) | | 4A | 3-4 oz. | 12h | 6 oz. |
| | dinotefuran | | E | 3d | NA |
| Warrior II (2.08CS) (RUP) | | 3A | 1.2-2.5 fl. oz. | 24h | 12.8 fl. oz. |
| | lambda-cyhalothrin | | E | 14d | NA |

Footnotes: See how to read the spray schedule tables section at the beginning of the chapter.

(Continued)

Peach Full Bloom - Diseases

Notes on disease management

- Use 12 oz. of Mycoshield/100 gallons of water (=150 ppm), varying volume depending upon tree size and foliar development.
- Quadris Top and Quilt Xcel contain azoxystrobin, which is known to be phytotoxic to certain apple varieties. Do not use where drift might affect apples.

- Application of Captan or Bravo as a tank mix is not recommended at this time due to the risk of phytotoxicity.
- The use of copper is not recommended during bloom to reduce phytotoxicity and protect pollinators.

Check with your local Extension Specialist for using copper during bloom.

Table 4-5. Peach Diseases - Full Bloom¹

| Product And Formulation Active Ingredient | FRAC Code ² | Brown Rot | REI ³ PHI ⁴ | Max Amt ⁵ Max App ⁶ |
|--|---------------------------|-----------------|--------------------------------------|--|
| Abound (SC) | 11 | 12-15.5 fl. oz. | 4h | 90 fl. oz. |
| azoxystrobin | | F-G[r] | 0d | See label |
| Bravo Weather Stik | M3 | 3.1-4.1 pt. | 12h | 20.5 pt. |
| chlorothalonil | | G | shuck-split | NA |
| Captan | M4 | 2.5-5 lb. | 24h | 40 lb. |
| captan | | G | 0d | NA |
| Cevya | 3 | 3-5 fl. oz. | 12h | 15 fl. oz. |
| mefentrifluconazole | | E | 0d | See label |
| Elevate 50 WDG | 17 | 1-1.5 lb. | 12h | 6 lb. |
| fenhexamid | | G-E | 0d | NA |
| Elite 45DF | 3 | 4-8 oz. | 12h | 3 lb. |
| tebuconazole | | E | 0d | NA |
| Flint Extra | 11 | 2.5-3.8 fl. oz. | 12h | 15.2 fl. oz. |
| trifloxystrobin | | s (G) | 1d | 4 |
| Fontelis (SC) | 7 | 14-20 fl. oz. | 12h | 61 fl. oz. |
| penthiopyrad | | E | 0d | NA |
| Indar 2F | 3 | 6 fl. oz. | 12h | 48 fl. oz. |
| fenbuconazole | | E[r] | 0d | 8 |
| Inspire Super (EW) | 3+9 | 16-20 fl. oz. | 12h | 80 fl. oz. |
| difenoconazole + cyprodinil | | E | 2d | 4 |
| Kenja 400 SC | 7 | 12.5 fl. oz. | 12h | 37.5 fl. oz. |
| isofetamid | | E | 1d | 3 |
| Luna Experience (SC) | 7+3 | 6-10 fl. oz. | 12h | 34 fl. oz. |
| fluopyram + tebuconazol | | G-E | 0d | NA |
| Luna Privilege | 7 | 4-6.8 fl. oz. | 12h | 13.7 fl. oz. |
| fluopyram | | E | 0d | NA |
| Luna Sensation (SC) | 7+11 | 5-7.6 fl. oz. | 12h | 27.1 fl. oz. |
| fluopyram + trifloxystrobin | | E | 1d | 4 |
| Merivon XBF | 7+11 | 4-6.7 fl. oz. | 12h | 20.1 fl. oz. |
| fluxapyroxad + pyraclostrobin | | E | 0d | 3 |

(Continued)

Table 4-5. Peach Diseases - Full Bloom¹ (continued)

| Product And Formulation Active Ingredient | FRAC Code ² | Brown Rot | REI ³ PHI ⁴ | Max Amt ⁵ Max App ⁶ |
|--|---------------------------|-----------------|--------------------------------------|--|
| Microthiol Disperss | M | 10-20 lb. | 24h | NA |
| sulfur | | F-P | 0d | NA |
| Miravis | 7 | 3.4-5.1 fl. oz. | 4h | 20.4 fl. oz. |
| pydiflumetofen | | E | 0d | 4 |
| Mycoshield | 41 | x | 12h | 12 lb. |
| oxytetracycline | | x | 21d | 8 |
| Ph-D | 19 | 6.2 oz. | 4h | NA |
| polyoxin D | | x | 0h | NA |
| Pristine | 7+11 | 10.5-14.5 oz. | 12h | 72.5 oz. |
| pyaclostrobin + boscalid | | E-G[r] | 0d | 5 |
| Quadris Top | 11+3 | 12-14 fl. oz. | 12h | 56 fl. oz. |
| azoxystrobin + difenoconazole | | G-E | 0d | 4 |
| Quash | 3 | 2.5-3.5 fl. oz. | 12h | 12 fl. oz. |
| metconazole | | G-E | 14d | 3 |
| Quilt Xcel | 11+3 | 14 fl. oz. | 12h | 70 fl. oz. |
| azoxystrobin + propiconazole | | E | 0d | 5 |
| Rally 40WSP | 3 | 2.5-6 oz. | 24h | 3.25 lb. |
| myclobutanil | | G | 0h | NA |
| Rovral 4F | 2 | 1-2 pt. | 24h | 4 pt. |
| iprodione | | E | PF | 2 |
| Scala (SC) | 9 | 9-18 fl. oz. | 12h | 54 fl. oz. |
| pyrimethanil | | E-G | 2d | 3 |
| Thiram Granuflo | M3 | 3.5 lb. | 24h | 21.2 lb. |
| thiram | | G | 7d | NA |
| Tilt (EC) | 3 | 4 fl. oz. | 12h | 20 fl. oz. |
| propiconazole | | E | 0d | 5 |
| Topguard EQ | 3+11 | 6-8 fl. oz. | 12h | NA |
| flutriafol + azoxystrobin | | G | 7d | 4 |
| Topguard Specialty Crop | 3 | 14 fl. oz. | 12h | 56 fl. oz. |
| flutriafol | | E | 7d | 4 |
| Topsin M WSB | 1 | 1-1.5 lb. | 48h | 4 lb. |
| thiophanate-methyl | | G[r] | 1d | NA |
| Vanguard WG (75WG) | 9 | 5 oz. | 12h | 30 oz. |
| cyprodinil | | G-E | 2d | 4 |
| Ziram 76DF | M3 | 4.5-8 lb. | 48h | 48.2 lb. |
| ziram | | G | 30d | 6 |

Footnotes: See how to read the spray schedule tables section at the beginning of the chapter.

Peach Full Bloom - Insects

Save the bees! Insecticide use for insects or mites is not recommended at this time.

Peach Petal Fall To Shuck Split - Diseases

Brown rot

- Last application for Rovral (iprodione) at petal fall. Do not apply Rovral after petal fall.
- Last application of Bravo or chlorothalonil product for control of brown rot and scab at shuck split.

Bacterial spot management:

- Some labels specify shuck-split, first and /or second cover sprays for application timings. Carefully read the label whenever using copper products to avoid phytotoxicity.

- If bacterial spot has been a problem, apply at 7-day intervals from petal fall (<5% shuck split) through first cover.
- For control of bacterial spot, reduce the rate of copper as the season progresses to reduce the risk of phytotoxicity. Do not apply under extended wet or humid conditions.
- Application of Bravo Weather Stik (BWS) plus Fontelis plus Mycoshield alternated at 7-day intervals with BWS plus Inspire Super plus Mycoshield from petal fall through shuck-split has been reported effective in controlling fungal and bacterial diseases during this period.

Peach scab applications begin now and are critical from shuck split through second-third cover.

Table 4-6. Peach Diseases - Petal Fall Through Shuck Split¹

| Product And Formulation Active Ingredient | FRAC Code ² | Bacterial Spot/Shot Hole | Brown Rot | Peach Scab | Powdery Mildew/Peach Rusty Spot | REI ³ PHI ⁴ | Max Amt ⁵ Max App ⁶ |
|--|---------------------------|--------------------------------|-----------------|-----------------|---------------------------------------|--------------------------------------|--|
| Abound (SC) | 11 | x | 12-15.5 fl. oz. | 12-15.5 fl. oz. | 12-15.5 fl. oz. | 4h | 90 fl. oz. |
| azoxystrobin | | x | F-G[r] | G | F-G | 0d | See label |
| Badge SC | M | 0.5-2 pt. | x | x | x | 24h | 63.4 pt. |
| copper sulfate + oxychloride | | F-G | x | x | x | 0d | NA |
| Bravo Weather Stik | M3 | x | 3.1-4.1 pt. | 3.1-4.1 pt. | 3.1-4.1 pt. | 12h | 20.5 pt. |
| chlorothalonil | | x | G | G | x | shuck-split | NA |
| Captan | M4 | x | 2.5-5 lb. | 2.5-5 lb. | 2.5-5 lb. | 24h | 40 lb. |
| captan | | x | G | G | F | 0d | NA |
| Cevya | 3 | x | 3-5 fl. oz. | 3-5 fl. oz. | 4-5 fl. oz. | 12h | 15 fl. oz. |
| mefentrifluconazole | | x | E | E | E | 0d | See label |
| C-O-C-S WDG | M | 1-2.9 lb. | 1-2.9 lb. | x | x | 48h | 35 lb. |
| copper oxychloride | | F-G | F | x | x | 21d | 3 |
| Cuprofix Ultra 40 disperss | M | 1 lb. | 1 lb. | x | x | 12h | 45 lb. |
| copper hydroxide | | G-F | F | x | x | 120d | NA |
| Elevate 50 WDG | 17 | x | 1-1.5 lb. | x | x | 12h | 6 lb |
| fenhexamid | | x | G-E | x | x | 0d | NA |
| Elite 45DF | 3 | x | 4-8 oz. | x | 4-8 oz. | 12h | 3 lb. |
| tebuconazole | | x | E | x | E | 0d | NA |
| Flint Extra | 11 | x | 2.5-3.8 fl. oz. | 2.5-3.8 fl. oz. | 2.5-3.8 fl. oz. | 12h | 15.2 fl. oz. |
| trifloxystrobin | | x | s (G) | E | E | 1d | 4 |
| Fontelis (SC) | 7 | x | 14-20 fl. oz. | 14-20 fl. oz. | 14-20 fl. oz. | 12h | 61 fl. oz. |
| penthiopyrad | | x | G-E | F-G | F-G | 0d | NA |
| Indar 2F | 3 | x | 6 fl. oz. | 6 fl. oz. | x | 12h | 48 fl. oz. |
| fenbuconazole | | x | E[r] | F | x | 0d | 8 |

(Continued)

Table 4-6. Peach Diseases - Petal Fall Through Shuck Split¹ (continued)

| Product And Formulation Active Ingredient | FRAC Code ² | Bacterial Spot/Shot Hole | Brown Rot | Peach Scab | Powdery Mildew/Peach Rusty Spot | REI ³ PHI ⁴ | Max Amt ⁵ Max App ⁶ |
|--|---------------------------|--------------------------------|-----------------|-----------------|---------------------------------------|--------------------------------------|--|
| Inspire Super (EW) | 3+9 | x | 16-20 fl. oz. | 16-20 fl. oz. | 16-20 fl. oz. | 12h | 80 fl. oz. |
| difenoconazole + cyprodinil | | x | E | F-G | F-G | 2d | 4 |
| Kenja 400 SC | 7 | x | 12.5 fl. oz. | 12.5 fl. oz. | x | 12h | 37.5 fl. oz. |
| isofetamid | | x | E | G | x | 1d | 3 |
| Kocide 3000 | M | 02.5-0.5 lb. | x | x | x | 48h | 60 lb |
| copper hydroxide | | G-F | x | x | x | 0d | NA |
| Luna Experience (SC) | 7+3 | x | 6-10 fl. oz. | 8-10 fl. oz. | 6-10 fl. oz. | 12h | 34 fl. oz. |
| fluopyram + tebuconazol | | x | G-E | E | G | 0d | NA |
| Luna Privilege | 7 | x | 4-6.8 fl. oz. | 4.8-6.8 fl. oz. | 4-6.84 fl. oz. | 12h | 13.7 fl. oz. |
| fluopyram | | x | E | F | G | 0d | NA |
| Luna Sensation (SC) | 7+11 | x | 5-7.6 fl. oz. | 5-7.6 fl. oz. | 5-7.6 fl. oz. | 12h | 27.1 fl. oz. |
| fluopyram + trifloxystrobin | | x | E | F | G-E | 1d | 4 |
| Merivon XBF | 7+11 | x | 4-6.7 fl. oz. | 4-6.7 fl. oz. | 4-6.7fl. oz. | 12h | 20.1 fl. oz. |
| fluxapyroxad + pyraclostrobin | | x | E | G-E | G-E | 0d | 3 |
| Microthiol Disperss | M | x | 10-20 lb. | x | 10-20 lb. | 24h | NA |
| sulfur | | x | F-P | x | F-P | 0d | NA |
| Miravis | 7 | x | 3.4-5.1 fl. oz. | 3.4-5.1 fl. oz. | 3.4-5.1 fl. oz. | 4h | 20.4 fl. oz. |
| pydiflumetofen | | x | E | E-G | E-G | 0d | 4 |
| Mycoshield | 41 | 12 oz./100 g. | x | x | x | 12h | 12 lb. |
| oxytetracycline | | E[r] | x | x | x | 21d | 8 |
| Ph-D | 19 | x | x | 6.2 oz. | 6.2 oz. | 4h | NA |
| polyoxin D | | x | x | G | u | 0h | NA |
| Pristine | 7+11 | x | 10.5-14.5 oz. | 10.5-14.5 oz. | 10.5-14.5 oz. | 12h | 72.5 oz. |
| pyaclostrobin + boscalid | | x | E-G[r] | E-G | E[r] | 0d | 5 |
| Quadris Top | 11+3 | x | 12-14 fl. oz. | 12-14 fl. oz. | 12-14 fl. oz. | 12h | 56 fl. oz. |
| azoxystrobin + difenoconazole | | x | E | u | G | 0d | 4 |
| Quash | 3 | x | 2.5-3.5 fl. oz. | 2.5-3.5 fl. oz. | 3.5-4 fl. oz. | 12h | 12 fl. oz. |
| metconazole | | x | G-E | G | E | 14d | 3 |
| Quilt Xcel | 11+3 | x | 14 fl. oz. | 14 fl. oz. | 14 fl. oz. | 12h | 70 fl. oz. |
| azoxystrobin + propiconazole | | x | E | G | G | 0d | 5 |
| Quintec | 13 | x | x | x | 7 fl. oz. | 12h | 28 fl. oz. |
| quinoxifen | | x | x | x | E | 7d | 4 |
| Rally 40WSP | 3 | x | 2.5-6 oz. | x | 2.5-6 oz. | 24h | 3.25 lb. |
| myclobutanil | | x | G | x | G-E | 0h | NA |
| Rovral 4F | 2 | x | 1-2 pt. | 1-2 qt. | x | 24h | 4 pt. |
| iprodione | | x | E | u | x | PF | 2 |
| Scala (SC) | 9 | x | 9-18 fl. oz. | x | x | 12h | 54 fl. oz. |
| pyrimethanil | | x | G-E | x | x | 2d | 3 |

(Continued)

Table 4-6. Peach Diseases - Petal Fall Through Shuck Split¹ (continued)

| Product And Formulation Active Ingredient | FRAC Code ² | Bacterial Spot/Shot Hole | Brown Rot | Peach Scab | Powdery Mildew/Peach Rusty Spot | REI ³ PHI ⁴ | Max Amt ⁵ Max App ⁶ |
|--|---------------------------|--------------------------------|-------------|---------------|---------------------------------------|--------------------------------------|--|
| Syllit F | U12 | x | 3 pt. | 3 pt. | x | 48h | 9 pt. |
| dodine | | x | s | F | x | petal fall | 3 |
| Thiram Granuflo | M3 | x | 3.5 lb. | 3.5 lb. | x | 24h | 21.2 lb. |
| thiram | | x | G | G | x | 7d | NA |
| Tilt (EC) | 3 | x | 4 fl. oz. | x | 4 fl. oz. | 12h | 20 fl. oz. |
| propiconazole | | x | E | x | G | 0d | 5 |
| Topguard EQ | 3+11 | x | 6-8 fl. oz. | 6-8 fl. oz. | 6-8 fl. oz. | 12h | NA |
| flutriafol + azoxystrobin | | x | G | u | E | 7d | 4 |
| Topguard Specialty Crop | 3 | x | 14 fl. oz. | 14 fl. oz. | 14 fl. oz. | 12h | 56 fl. oz. |
| flutriafol | | x | E | s(G) | G | 7d | 4 |
| Topsin M WSB | 1 | x | 1-1.5 lb. | 1-1.5 lb. | 1-1.5 lb. | 48h | 4 lb. |
| thiophanate-methyl | | x | G[r] | G | G[r] | 1d | NA |
| Vanguard WG (75WG) | 9 | x | 5 oz. | x | x | 12h | 30 oz. |
| cyprodinil | | x | G-E | x | x | 2d | 4 |
| Ziram 76DF | M3 | x | 4.5-8 lb. | 4.5-8 lb. | x | 48h | 48.2 lb. |
| ziram | | x | G | G | x | 30d | 6 |

Footnotes: See how to read the spray schedule tables section at the beginning of the chapter.

Peach Petal Fall To Shuck Split - Insects

Table 4-7. Peach Insects - Petal Fall Through Shuck Split¹

| Product And Formulation Active Ingredient | IRAC Code ² | Borers | Cherry Fruit Fly | European Red Mite | Green Peach Aphid | Japanese Beetle | Oriental Fruit Moth | Plant Bug/ Stink Bug | Plum Curculio | San Jose Scale | Spotted-Wing Drosophila | REI ³ PHI ⁴ | Max Amt ⁵ Max App ⁶ |
|--|---------------------------|----------------|---------------------|----------------------|----------------------|--------------------|------------------------|-------------------------|------------------|--------------------|----------------------------|--------------------------------------|--|
| Acramite 50WS | 25 | x | x | 1 lb. | x | x | x | x | x | x | x | 12h | NA |
| bifenazate | | x | x | E | x | x | x | x | x | x | x | 3h | 1 |
| Actara (25WDG) | 4A | x | 4.5-5.5 oz. | x | 3-4 oz. | x | x | 4.5-5.5 oz. | 4.5-5.5 oz. | x | x | 12h | 11 oz. |
| thiamethoxam | | x | F | x | E | x | x | G | G | x | x | 14d | NA |
| Admire Pro (4.6F) | 4A | x | 2-2.8 fl. oz. | x | 1.4-2.8 fl. oz. | 1.4-2.8 fl. oz. | x | 2.8 fl. oz. | 2.8 fl. oz. | 1.4-2.8 fl. oz. | x | 12h | 8.4 fl. oz. |
| imidacloprid | | x | F | x | E | G | x | s | s | F | x | 0d | NA |
| Altacor eVo | 28 | 1.5-2.2 oz. | 1.5-2.2 oz. | x | x | x | 1.5-2.2 oz. | x | x | x | x | 4h | 4.6 oz. |
| chlorantraniliprole | | u | s | x | x | x | E | x | x | x | x | 10d | 3 |
| Apta (1.34SC) | 21A | x | 14-27 fl. oz. | x | 17-27 fl. oz. | x | x | 21-27 fl. oz. | 21-27 fl. oz. | x | 21-27 fl. oz. | 12h | 53.5 fl. oz. |
| tolfenpyrad | | x | u | x | G | x | x | S | G | x | s | 14d | 2 |

(Continued)

Table 4-7. Peach Insects - Petal Fall Through Shuck Split¹ (continued)

| Product And Formulation Active Ingredient | IRAC Code ² | Borers | Cherry Fruit Fly | European Red Mite | Green Peach Aphid | Japanese Beetle | Oriental Fruit Moth | Plant Bug/ Stink Bug | Plum Curculio | San Jose Scale | Spotted-Wing Drosophila | REI ³ PHI ⁴ | Max Amt ⁵ Max App ⁶ |
|--|------------------------|------------------|------------------|------------------------|------------------------|-------------------|------------------------|-------------------------|-------------------|------------------------|-------------------------|--------------------------------------|--|
| Asana XL (0.66EC) (RUP) | 3A | 4.8-14.5 fl. oz. | 4.8-14.5 fl. oz. | x | x | x | 4.8-14.5 fl. oz. | 4.8-14.5 fl. oz. | 4.8-14.5 fl. oz. | x | x | 12h | 72.5 fl. oz. |
| esfenvalerate | | G | G | x | x | x | E[r] | G | G | x | x | 14d | NA |
| Assail 30SG | 4A | 5.3-8 oz. | 5.3-8 oz. | x | 2.5-5.3 oz. | 5.3-8 oz. | 5.3-8 oz. | 5.3-8 oz. | 5.3-8 oz. | 5.3-8 oz. | x | 12h | 32 oz. |
| acetamiprid | | G | F | x | E | G | E | F | G | F | x | 7d | 4 |
| Avaunt eVo (30WDG) | 22A | x | x | x | x | x | 6 oz. | x | 5-6 oz. | x | x | 12h | 24 oz. |
| indoxacarb | | x | x | x | x | x | G | x | E | x | x | 14d | 4 |
| Baythroid XL (1EC) (RUP) | 3A | 1.4-2 fl. oz. | 2.4-2.8 fl. oz. | x | x | x | 2-2.4 fl. oz. | 2-2.4 fl. oz. | 2.4-2.8 fl. oz. | x | x | 12h | 5.6 fl. oz. |
| beta-cyfluthrin | | G | G | x | x | x | E[r] | E | G | x | x | 7d | NA |
| Belay (2.13SC) | 4A | x | x | x | 3 - 6 fl. oz. | x | x | 6 fl. oz. | 6 fl. oz. | 6 fl. oz. | x | 12h | 12 fl. oz. |
| clothianidin | | x | x | x | E | x | x | E | G | G | x | 21d | NA |
| Beleaf 50SG | 29 | x | x | x | 2-2.8 oz. | x | x | 2-2.8 oz. | x | x | x | 12h | 8.4 oz. |
| flonicamid | | x | x | x | E | x | x | G | x | x | x | 14d | 3 |
| Danitol 2.4EC (RUP) | 3A | x | 16-21.3 fl. oz. | 10.7-21.3 fl. oz. | 10.7-21.3 fl. oz. | 10.7-21.3 fl. oz. | 10.7-21.3 fl. oz. | 10.7-21.3 fl. oz. | 10.7-21.3 fl. oz. | x | 10.7-21.3 fl. oz. | 24h | 42.7 fl. oz. |
| fenpropathrin | | x | G | i | F | E | E[r] | E | G | x | E | 3d | NA |
| Delegate WG (25WG) | 5 | x | 6-7 oz. | x | x | x | 6-7 oz. | x | 6-7 oz. | x | 4.5-7 oz. | 4h | 28 oz. |
| spinetoram | | x | s | x | x | x | E | x | s | x | G | 1d | 4 |
| Diazinon AG 600WBC (RUP) | 1B | x | x | 12.75 fl. oz./100 gal. | 12.75 fl. oz./100 gal. | x | 12.75 fl. oz./100 gal. | x | x | 12.75 fl. oz./100 gal. | x | 4d | 51 fl. oz. |
| diazinon | | x | x | i | G | x | G | x | x | F | x | 21d | 2 |
| Dimilin 2L (RUP) | 15 | x | x | x | x | x | 8-16 fl. oz. | x | 8-16 fl. oz. | x | x | 12h | 32 fl. oz. |
| | | x | x | x | x | x | x | x | u | x | x | 14d | 2 |
| Entrust SC (2SC) | 5 | x | 4-8 fl. oz. | x | x | x | 4-8 fl. oz. | x | x | x | 4-8 fl. oz. | 4h | 29 fl. oz. |
| spinosad | | x | F | x | x | x | F | x | x | x | G-E | 7d | 3 |
| Esteem 35WP | 7C | x | x | x | x | x | 4-5 oz. | x | x | 4-5 oz. | x | 12h | 15 oz. |
| pyriproxyfen | | x | x | x | x | x | s | x | x | E | x | 14d | 3 |
| Exirel (0.83SE) | 28 | x | 10-17 fl. oz. | x | x | 13.5-20.5 fl. oz. | 10-20.5 fl. oz. | x | 13.5-20.5 fl. oz. | x | 13.5-20.5 fl. oz. | 12h | 61.5 fl. oz. |
| cyantraniliprole | | x | E | x | x | G | E | x | G | x | E | 3d | 3 |

(Continued)

Table 4-7. Peach Insects - Petal Fall Through Shuck Split¹ (continued)

| Product And Formulation Active Ingredient | IRAC Code ² | Borers | Cherry Fruit Fly | European Red Mite | Green Peach Aphid | Japanese Beetle | Oriental Fruit Moth | Plant Bug/ Stink Bug | Plum Curculio | San Jose Scale | Spotted-Wing Drosophila | REI ³ PHI ⁴ | Max Amt ⁵ Max App ⁶ |
|--|------------------------|------------------|------------------|-------------------|-------------------|------------------|---------------------|-------------------------|----------------|------------------|-------------------------|--------------------------------------|--|
| Imidan 70W | 1B | x | 2.1-2.5 lb. | x | x | 2.1-4.25 lb. | 2.13-4.25 lb. | x | 2.1-4.25 lb. | 2.1-4.25 lb. | 2.1-4.25 lb. | 4d | 17 lb. |
| phosmet | | x | G | x | x | G | E[r] | x | G | E | E | 14d | NA |
| Intrepid 2F | 18 | x | x | x | x | x | 10-16 fl. oz. | x | x | x | x | 4h | 64 fl. oz. |
| methoxyfenozide | | x | x | x | x | x | G | x | x | x | x | 7d | NA |
| Lannate LV | 1A | x | x | x | 3 pt. | x | 3 pt. | 3 pt. | x | x | x | 4d | 18 pt. |
| methomyl | | x | x | x | G | x | F | G | x | x | x | 4d | 6 |
| Malathion 5EC | 1B | x | x | 2.5-4.8 pt. | 2.5-4.8 pt. | 2.5-4.8 pt. | 4.8 pt. | x | 4.8 pt. | x | x | 24h | 4.8 pt. |
| malathion | | | | u | u | u | u | | u | | | 7d | 3 |
| Mustang Maxx (0.83EC) (RUP) | 3A | 1.28-4 fl. oz. | 1.28-4 fl. oz. | x | x | x | 1.2-4 fl. oz. | 1.2-4 fl. oz. | 1.2-4 fl. oz. | x | 4 fl. oz. | 12h | 24 fl. oz. |
| zeta-cypermethrin | | G | F | x | x | x | E[r] | E | G | x | E | 14d | NA |
| Neemix 4.5 (0.39L) | UN | 7-16 fl. oz. | 7-16 fl. oz. | x | 5-7 fl. oz. | x | 7-16 fl. oz. | 7-16 fl. oz. | x | 7-16 fl. oz. | x | 4h | NA |
| azadirachtin | | F | u | x | G | x | u | E | i | G | x | 0d | NA |
| Nexter SC (3.755C) | 21 | x | x | 7.5-17 fl. oz. | x | x | x | x | x | x | x | 12h | 34 fl. oz. |
| pyridaben | | x | x | u | x | x | x | x | x | x | x | 7d | 2 |
| Pounce 25WP | 3A | 6.4-16 oz. | x | x | x | x | 6.4-16 oz. | x | 6.4-16 oz. | x | x | 12h | 48 oz. |
| permethrin | | F | x | x | x | x | E[r] | x | G | x | x | 14d | NA |
| Pyganic 5EC | 3A | 4.5-15.6 fl. oz. | 4.5-15.6 fl. oz. | 4.5-15.6 fl. oz. | 4.5-15.6 fl. oz. | 4.5-15.6 fl. oz. | x | 4.5-15.6 fl. oz. | x | 4.5-15.6 fl. oz. | 4.5-15.6 fl. oz. | 12h | 15.6 fl. oz. |
| pyrethrins | | u | i | u | G | u | x | u | x | u | F | 0d | 1 |
| Rimon 0.83EC | 15 | 20 fl. oz. | 20-40 fl. oz. | x | x | x | 20-40 fl. oz. | 20-40 fl. oz. | x | x | 20-40 fl. oz. | 12h | 150 fl. oz. |
| novaluron | | u | u | x | x | x | E | u | x | x | G | 8d | NA |
| Scorpion 35SL | 4A | 5.25-7 fl. oz. | x | x | 3.5-7 fl. oz. | x | x | 5.25-7 fl. oz. | 5.25-7 fl. oz. | x | x | 12h | 14.25 fl. oz. |
| dinotefuran | | s | x | x | s | x | x | E | s | x | x | 3/21d | NA |
| Sevin XLR Plus (4F) | 1A | 2-3 qt. | 2-3 qt. | x | x | 2-3 qt. | 2-3 qt. | 2-3 qt. | 2-3 qt. | 4-5 qt. | x | 12h | 14 qt. |
| carbaryl | | u | G | x | x | E | F | F | F | u | x | 3d | 3 |
| Surround WP (95WP) | UN | x | 25-50 lb. | x | x | 25-50 lb. | 25-50 lb. | x | 25-50 lb. | x | x | 4h | NA |
| kaolin | | x | s | x | x | s | s | x | s | x | x | 0d | NA |

(Continued)

Table 4-7. Peach Insects - Petal Fall Through Shuck Split¹ (continued)

| Product And Formulation Active Ingredient | IRAC Code ² | Borers | Cherry Fruit Fly | European Red Mite | Green Peach Aphid | Japanese Beetle | Oriental Fruit Moth | Plant Bug/ Stink Bug | Plum Curculio | San Jose Scale | Spotted-Wing Drosophila | REI ³ PHI ⁴ | Max Amt ⁵ Max App ⁶ |
|--|------------------------|-----------------|------------------|-------------------|-------------------|-----------------|---------------------|-------------------------|-----------------|----------------|-------------------------|--------------------------------------|--|
| Venom (70SG) | 4A | 3-4 oz. | x | x | 2-4 oz. | x | x | 3-4 oz. | 3-4 oz. | x | x | 12h | . |
| dinotefuran | | u | x | x | s | x | x | E | u | x | x | 3d | NA |
| Verdepryn 100SL (0.83SL) | 28 | x | 5.5-11 fl. oz. | x | 5.5-11 fl. oz. | x | 5.5-11 fl. oz. | 5.5-11 fl. oz. | 5.5-11 fl. oz. | x | 5.5-11 fl. oz. | 4h | 33 fl. oz. |
| cyclaniliprole | | x | G | x | E | x | E | s | u | x | u | 7d | 3 |
| Warrior II (2.08CS) (RUP) | 3A | 1.2-2.5 fl. oz. | 1.2-2.5 fl. oz. | x | x | 1.2-2.5 fl. oz. | 1.2-2.5 fl. oz. | 1.2-2.5 fl. oz. | 1.2-2.5 fl. oz. | x | x | 24h | 12.8 fl. oz. |
| lambda-cyhalothrin | | G | G | x | x | E | G[r] | E | G | x | x | 14d | NA |

Footnotes: See how to read the spray schedule tables section at the beginning of the chapter.

Peach First Cover - Diseases

7-10 days after shuck split.

Table 4-8. Peach Diseases - First Cover¹

| Product And Formulation Active Ingredient | FRAC Code ² | Bacterial Spot/ Shot Hole | Brown Rot | Peach Scab | Powdery Mildew/ Peach Rusty Spot | REI ³ PHI ⁴ | Max Amt ⁵ Max App ⁶ |
|--|------------------------|------------------------------|-----------------|-----------------|-------------------------------------|--------------------------------------|--|
| Abound (SC) | 11 | x | 12-15.5 fl. oz. | 9-15.5 fl. oz. | 12-15.5 fl. oz. | 4h | 90 fl. oz. |
| azoxystrobin | | x | F-G[r] | G | F | 0d | See label |
| Badge SC | M | 0.5-2 pt. | x | x | x | 24h | 63.4 pt. |
| copper sulfate + oxychloride | | F-G | x | x | x | 0d | NA |
| Captan | M4 | x | 2.5-5 lb. | 2.5-5 lb. | 2.5-5 lb. | 24h | 40 lb. |
| captan | | x | G | G | F | 0d | NA |
| Cevya | 3 | x | 3-5 fl. oz. | 3-5 fl. oz. | 4-5 fl. oz. | 12h | 15 fl. oz. |
| mefentrifluconazole | | x | E | E | E | 0d | See label |
| Elevate 50 WDG | 17 | x | 1-1.5 lb. | x | x | 12h | 6 lb. |
| fenhexamid | | x | G-E | x | x | 0d | NA |
| Elite 45DF | 3 | x | 4-8 oz. | x | 4-8 oz. | 12h | 3 lb. |
| tebuconazole | | x | E | x | E | 0d | NA |
| Flint Extra | 11 | x | 2.5-3.8 fl. oz. | 2.5-3.8 fl. oz. | 2.5-3.8 fl. oz. | 12h | 15.2 fl. oz. |
| trifloxystrobin | | x | s (G) | E | E | 1d | 4 |
| Fontelis (SC) | 7 | x | 14-20 fl. oz. | 14-20 fl. oz. | 14-20 fl. oz. | 12h | 61 fl. oz. |
| penthiopyrad | | x | G-E | F-G | F-G | 0d | NA |
| Indar 2F | 3 | x | 6 fl. oz. | 6 fl. oz. | x | 12h | 48 fl. oz. |
| fenbuconazole | | x | E[r] | F | x | 0d | 8 |

(Continued)

Table 4-8. Peach Diseases - First Cover¹ (continued)

| Product And Formulation Active Ingredient | FRAC Code ² | Bacterial Spot/ Shot Hole | Brown Rot | Peach Scab | Powdery Mildew/ Peach Rusty Spot | REI ³ PHI ⁴ | Max Amt ⁵ Max App ⁶ |
|--|---------------------------|---------------------------------|-----------------|-----------------|---|--------------------------------------|--|
| Inspire Super (EW) | 3+9 | x | 16-20 fl. oz. | 16-20 fl. oz. | 16-20 fl. oz. | 12h | 80 fl. oz. |
| difenoconazole + cyprodinil | | x | E | F-G | F-G | 2d | 4 |
| Kenja 400 SC | 7 | x | 12.5 fl. oz. | 12.5 fl. oz. | x | 12h | 37.5 fl. oz. |
| isofetamid | | x | E | G | x | 1d | 3 |
| Kocide 3000 | M | 0.25-0.5 lb. | x | x | x | 48h | 60 lb. |
| copper hydroxide | | G-F | x | x | x | 0d | NA |
| Luna Experience (SC) | 7+3 | x | 6-10 fl. oz. | 8-10 fl. oz. | 6-10 fl. oz. | 12h | 34 fl. oz. |
| fluopyram + tebuconazol | | x | G-E | E | G | 0d | NA |
| Luna Privilege | 7 | x | 4-6.8 fl. oz. | 4.8-6.8 fl. oz. | 4-6.8 fl. oz. | 12h | 13.7 fl. oz. |
| fluopyram | | x | E | F | G | 0d | NA |
| Luna Sensation (SC) | 7+11 | x | 5-7.6 fl. oz. | 5-7.6 fl. oz. | 5-7.6 fl. oz. | 12h | 27.1 fl. oz. |
| fluopyram + trifloxystrobin | | x | E | F | G-E | 1d | 4 |
| Merivon XBF | 7+11 | x | 4-6.7 fl. oz. | 4-6.7 fl. oz. | 4-6.7 fl. oz. | 12h | 20.1 fl. oz. |
| fluxapyroxad + pyraclostrobin | | x | E | G-E | G-E | 0d | 3 |
| Microthiol Disperss | M | x | 10-20 lb. | x | 10-20 lb. | 24h | NA |
| sulfur | | x | F-P | x | F-P | 0d | NA |
| Miravis | 7 | x | 3.4-5.1 fl. oz. | 3.4-5.1 fl. oz. | 3.4-5.1 fl. oz. | 4h | 20.4 fl. oz. |
| pydiflumetofen | | x | E | G-E | G-E | 0d | 4 |
| Mycoshield | 41 | 12 oz./100 g. | x | x | x | 12h | 12 lb. |
| oxytetracycline | | E[r] | x | x | x | 21d | 8 |
| Ph-D | 19 | x | x | 6.2 oz. | 6.2 oz. | 4h | NA |
| polyoxin D | | x | x | G | u | 0h | NA |
| Pristine | 7+11 | x | 10.5-14.5 oz. | 10.5-14.5 oz. | 10.5-14.5 oz. | 12h | 72.5 oz. |
| pyraclostrobin + boscalid | | x | G[r] | G-E | E[r] | 0d | 5 |
| Quadris Top | 11+3 | x | 12-14 fl. oz. | 12-14 fl. oz. | 12-14 fl. oz. | 12h | 56 fl. oz. |
| azoxystrobin + difenoconazole | | x | E | u | G | 0d | 4 |
| Quash | 3 | x | 2.5-3.5 fl. oz. | 2.5-3.5 fl. oz. | 3.5-4 fl. oz. | 12h | 12 fl. oz. |
| metconazole | | x | G-E | G | E | 14d | 3 |
| Quilt Xcel | 11+3 | x | 14 fl. oz. | 14 fl. oz. | 14 fl. oz. | 12h | 70 fl. oz. |
| azoxystrobin + propiconazole | | x | E | G | G | 0d | 5 |
| Quintec | 13 | x | x | x | 7 fl. oz. | 12h | 28 fl. oz. |
| quinoxifen | | x | x | x | E | 7d | 4 |
| Rally 40WSP | 3 | x | 2.5-6 oz. | x | 2.5-6 oz. | 24h | 3.25 lb. |
| myclobutanil | | x | G | x | G-E | 0h | NA |
| Scala (SC) | 9 | x | 9-18 fl. oz. | x | x | 12h | 54 fl. oz. |
| pyrimethanil | | x | G-E | x | x | 2d | 3 |
| Thiram Granuflo | M3 | x | 3.5 lb. | 3.5 lb. | x | 24h | 21.2 lb. |
| thiram | | x | G | G | x | 7d | NA |

(Continued)

Table 4-8. Peach Diseases - First Cover¹ (continued)

| Product And Formulation Active Ingredient | FRAC Code ² | Bacterial Spot/ Shot Hole | Brown Rot | Peach Scab | Powdery Mildew/ Peach Rusty Spot | REI ³ PHI ⁴ | Max Amt ⁵ Max App ⁶ |
|--|---------------------------|---------------------------------|-------------|---------------|---|--------------------------------------|--|
| Tilt (EC) | 3 | x | 4 fl. oz. | x | 4 fl. oz. | 12h | 20 fl. oz. |
| propiconazole | | x | E | x | G | 0d | 5 |
| Topguard EQ | 3+11 | x | 6-8 fl. oz. | 6-8 fl. oz. | 6-8 fl. oz. | 12h | NA |
| flutriafol + azoxystrobin | | x | G | u | E | 7d | 4 |
| Topguard Specialty Crop | 3 | x | 14 fl. oz. | x | 14 fl. oz. | 12h | 56 fl. oz. |
| flutriafol | | x | E | s(G) | G | 7d | 4 |
| Topsin M WSB | 1 | x | 1-1.5 lb. | 1-1.5 lb. | 1-1.5 lb. | 48h | 4 lb. |
| thiophanate-methyl | | x | G[r] | G | G[r] | 1d | NA |
| Vanguard WG (75WG) | 9 | x | 5 oz. | x | x | 12h | 30 oz. |
| cyprodinil | | x | G-E | x | x | 2d | 4 |
| Ziram 76DF | M3 | x | 4.5-8 lb. | 4.5-8 lb. | x | 48h | 48.2 lb. |
| ziram | | x | G | G | x | 30d | 6 |

Footnotes: See how to read the spray schedule tables section at the beginning of the chapter.

Peach First To Second Cover - Insects

7-10 days after shuck split. Second cover occurs 7-14 days after first cover.

Notes on insect management

San Jose scale

- Movento 2SC, must be tank mixed with a spray adjuvant.

Lesser peachtree borer

- Control of the first generation of lesser peachtree borer with trunk sprays is during the time of peak moth flight, generally from early May to early June, depending on latitude and spring temperatures. See Borers of Peach, Cherry, and Plum Trees, page 152.

Table 4-9. Peach Insects - First Through Second Cover¹

| Product And Formulation Active Ingredient | IRAC Code ² | Borers | Oriental Fruit Moth | Plant Bug/ Stink Bug | Plum Curculio | San Jose Scale | REI ³ PHI ⁴ | Max Amt ⁵ Max App ⁶ |
|--|---------------------------|------------------|------------------------|----------------------------|------------------|-------------------|--------------------------------------|--|
| Actara (25WDG) | 4A | x | x | 4.5-5.5 oz. | 4.5-5.5 oz. | x | 12h | 11 oz. |
| thiamethoxam | | x | x | G | G | x | 14d | NA |
| Admire Pro (4.6F) | 4A | x | x | 2.8 fl. oz. | 2.8 fl. oz. | 1.4-2.8 fl. oz. | 12h | 8.4 fl. oz. |
| imidacloprid | | x | x | s | s | F | 0d | NA |
| Altacor eVo | 28 | 1.5-2.2 oz. | 1.5-2.2 oz. | x | x | x | 4h | 4.6 oz. |
| chlorantraniliprole | | u | E | x | x | x | 10d | 3 |
| Apta (1.34SC) | 21A | x | x | 21-27 fl. oz. | 21-27 fl. oz. | x | 12h | 53.5 fl. oz. |
| tolfenpyrad | | x | x | s | G | x | 14d | 2 |
| Asana XL (0.66EC) (RUP) | 3A | 4.8-14.5 fl. oz. | 4.8-14.5 fl. oz. | 4.8-14.5 fl. oz. | 4.8-14.5 fl. oz. | x | 12h | 72.5 fl. oz. |
| esfenvalerate | | G | E[r] | G | G | x | 14d | NA |

(Continued)

Table 4-9. Peach Insects - First Through Second Cover¹ (continued)

| Product And Formulation Active Ingredient | IRAC Code ² | Borers | Oriental Fruit Moth | Plant Bug/ Stink Bug | Plum Curculio | San Jose Scale | REI ³ PHI ⁴ | Max Amt ⁵ Max App ⁶ |
|--|---------------------------|---------------|------------------------|----------------------------|-------------------|------------------------|--------------------------------------|--|
| Assail 30SG | 4A | x | 5.3-8 oz. | 5.3-8 oz. | 5.3-8 oz. | 5.3-8 oz. | 12h | 32 oz. |
| acetamiprid | | x | E | F | G | F | 7d | 4 |
| Avaunt eVo (30WDG) | 22 | x | 5.6 oz. | x | 5-6 oz. | x | 12h | 24 oz. |
| indoxacarb | | x | G | x | E | x | 14d | 4 |
| Baythroid XL (1EC) (RUP) | 3A | 1.4-2 fl. oz. | 2-2.4 fl. oz. | 2-2.4 fl. oz. | 2.4-2.8 fl. oz. | x | 12h | 5.6 fl. oz. |
| beta-cyfluthrin | | G | E[r] | E | G | x | 7d | NA |
| Belay (2.13SC) | 4A | x | x | 6 fl. oz. | 6 fl. oz. | 6 fl. oz. | 12h | 12 fl. oz. |
| clothianidin | | x | x | E | G | G | 21d | NA |
| Beleaf 50SG | 29 | x | x | 2-2.8 oz. | x | x | 12h | 8.4 oz. |
| flonicamid | | x | x | G | x | x | 14d | 3 |
| Centaur WDG (70WDG) | 16 | x | x | x | x | 34.5 oz. | 12h | 69 oz. |
| buprofezin | | x | x | x | x | E | 14d | 2 |
| Danitol 2.4EC (RUP) | 3A | x | 10.7-21.3 fl. oz. | 10.7-21.3 fl. oz. | 10.7-21.3 fl. oz. | x | 24h | 42.7 fl. oz. |
| fenpropathrin | | x | E[r] | E | G | x | 3d | NA |
| Delegate WG (25WG) | 5 | x | 6-7 oz. | x | 6-7 oz. | x | 4h | 28 oz. |
| spinetoram | | x | E | x | s | x | 1d | 4 |
| Diazinon AG 600WBC (RUP) | 1B | x | 12.75 fl. oz./100 gal. | x | x | 12.75 fl. oz./100 gal. | 4d | 51 fl. oz. |
| diazinon | | x | G | x | x | F | 21d | 2 |
| Dimilin 2L (RUP) | 15 | x | 8-16 fl. oz. | x | 8-16 fl. oz. | x | 12h | 32 fl. oz. |
| diflubenzuron | | x | u | x | u | x | 14d | 2 |
| Entrust SC (2SC) | 5 | x | 4-8 fl. oz. | x | x | x | 4h | 29 fl. oz. |
| spinosad | | x | F | x | x | x | 7d | 3 |
| Esteem 35WP | 7C | x | 4-5 oz. | x | x | 4-5 oz. | 12h | 15 oz. |
| pyriproxifen | | x | s | x | x | E | 14d | 3 |
| Exirel (0.83SE) | 28 | x | 10-20.5 fl. oz. | x | 13.5-20.5 fl. oz. | x | 12h | 61.5 fl. oz. |
| cyantraniliprole | | x | E | x | G | x | 3d | 3 |
| Imidan 70W | 1B | x | 2.13-4.25 lb. | x | 2.1-4.25 lb. | 2.1-4.25 lb. | 4d | 17 lb. |
| phosmet | | x | E[r] | x | G | E | 14d | NA |
| Intrepid 2F | 18 | x | 10-16 fl. oz. | x | x | x | 4h | 64 fl. oz. |
| methoxyfenozide | | x | G | x | x | x | 7d | NA |
| Lannate LV | 1A | x | 3 pt. | 3 pt. | x | x | 4d | 18 pt. |
| methomyl | | x | F | G | x | x | 4d | 6 |
| Malathion 5EC (8EC) | 1B | x | 4.8 pt. | x | 4.8 pt. | x | 24h | 4.8 pt. |
| malathion | | x | u | x | u | x | 7d | 3 |

(Continued)

Table 4-9. Peach Insects - First Through Second Cover¹ (continued)

| Product And Formulation Active Ingredient | IRAC Code ² | Borers | Oriental Fruit Moth | Plant Bug/ Stink Bug | Plum Curculio | San Jose Scale | REI ³ PHI ⁴ | Max Amt ⁵ Max App ⁶ |
|--|---------------------------|---------------------|------------------------|----------------------------|--------------------|---------------------|--------------------------------------|--|
| Movento (2SC) | 23 | x | x | x | x | 6-9 fl. oz. | 24h | 15.3 fl. oz. |
| spirotetramat | | x | x | x | x | G | 7d | NA |
| Mustang Maxx (0.83EC) (RUP) | 3A | 1.2-4 fl. oz. | 1.2-4 fl. oz. | 1.2-4 fl. oz. | 1.2-4 fl. oz. | x | 12h | 24 fl. oz. |
| zeta-cypermethrin | | G | E[r] | E | G | x | 14d | NA |
| Neemix 4.5 (0.39L) | UN | 7-16 fl. oz. | 7-16 fl. oz. | 7-16 fl. oz. | x | 7-16 fl. oz. | 4h | NA |
| azadirachtin | | F | u | E | i | G | 0d | NA |
| Pounce 25WP | 3A | 6.4-16 oz. | 6.4-16 oz. | x | 6.4-16 oz. | x | 12h | 48 fl. oz. |
| permethrin | | F | E[r] | x | G | x | 14d | NA |
| Pyganic 5EC | 3A | 4.5-15.6 fl. oz. | x | 4.5-15.6 fl. oz. | x | 4.5-15.6 fl. oz. | 12h | 15.6 fl. oz. |
| pyrethrins | | u | x | u | x | u | 0d | 1 |
| Rimon 0.83EC | 15 | 20 fl. oz. | 20-40 fl. oz. | 20-40 fl. oz. | x | x | 12h | 150 fl. oz. |
| novaluron | | u | E | u | x | x | 8d | NA |
| Scorpion 35SL | 4A | 5.25-7 fl. oz. | x | 5.25-7 fl. oz. | 5.25-7 fl. oz. | x | 12h | 14.25 fl. oz. |
| dinotefuran | | s | x | E | s | x | 3/21d | NA |
| Sevin XLR Plus (4F) | 1A | 2-3 qt. | 2-3 qt. | 2-3 qt. | 2-3 qt. | 2-3 qt. | 12h | 14 qt. |
| carbaryl | | u | F | F | F | u | 3d | 3 |
| Sivanto Prime (1.67 SL) | 4D | x | x | x | x | 10.5-14 fl. oz. | 4h | 28 fl. oz. |
| flupyradifurone | | x | x | x | x | G | 14d | NA |
| Surround WP (95WP) | UN | 25-50 lb. | 25-50 lb. | x | 25-50 lb. | x | 4h | NA |
| kaolin | | x | s | x | s | x | 0d | NA |
| Venom (70SG) | 4A | 3-4 oz. | x | 3-4 oz. | 3-4 oz. | x | 12h | 6 oz. |
| dinotefuran | | u | x | E | u | x | 3d | NA |
| Verdepryn 100SL (0.83SL) | 28 | x | 5.5-11 fl. oz. | 5.5-11 fl. oz. | 5.5-11 fl. oz. | x | 4h | 33 fl. oz. |
| cyclaniliprole | | x | E | s | u | x | 7d | 3 |
| Warrior II (2.08CS) (RUP) | 3A | 1.2-2.5 fl. oz. | 1.2-2.5 fl. oz. | 1.2-2.5 fl. oz. | 1.2-2.5 fl. oz. | x | 24h | 12.8 fl. oz. |
| lambda-cyhalothrin | | G | G[r] | E | G | x | 14d | NA |

Footnotes: See how to read the spray schedule tables section at the beginning of the chapter.

Peach Summer Cover Sprays - Diseases

10 days after first cover. Until ~3 weeks prior to harvest

- Be aware of PHI when applying fungicides to early harvested varieties.
- Remember application limits when applying cop-pers to control bacterial spot.

- PHI of Mycoshield is 21 days.
- When environmental conditions are conducive for brown rot, scab or powdery mildew, maintain fungicide schedule every 7-14 days.

Table 4-10. Peach Diseases - Summer Cover¹

| Product And Formulation Active Ingredient | FRAC Code ² | Bacterial Spot/Shot Hole | Brown Rot | Peach Scab | Powdery Mildew/Peach Rusty Spot | REI ³ PHI ⁴ | Max Amt ⁵ Max App ⁶ |
|--|---------------------------|--------------------------------|-----------------|-----------------|---------------------------------------|--------------------------------------|--|
| Abound (SC) | 11 | x | 12-15.5 fl. oz. | 9-15.5 fl. oz. | 12-15.5 fl. oz. | 4h | 90 fl. oz. |
| azoxystrobin | | x | F-G[r] | G | F | 0d | See label |
| Badge SC | M | 0.5-2 pt. | x | x | x | 24h | 63.4 pt. |
| copper sulfate + oxychloride | | F-G | x | x | x | 0d | NA |
| Captan | M4 | x | 2.5-5 lb. | 2.5-5 lb. | 2.5-5 lb. | 24h | 40 lb. |
| captan | | x | G | G | F | 0d | NA |
| Cevya | 3 | x | 3-5 fl. oz. | 3-5 fl. oz. | 4-5 fl. oz. | 12h | 15 fl. oz. |
| mefentrifluconazole | | x | E | E | E | 0d | See label |
| Elevate 50 WDG | 17 | x | 1-1.5 lb. | x | x | 12h | 6 lb. |
| fenhexamid | | x | G-E | x | x | 0d | NA |
| Elite 45DF | 3 | x | 4-8 oz. | x | 4-8 oz. | 12h | 3 lb. |
| tebuconazole | | x | E | x | E | 0d | NA |
| Flint Extra | 11 | x | 2.5-3.8 fl. oz. | 2.5-3.8 fl. oz. | 2.5-3.8 fl. oz. | 12h | 15.2 fl. oz. |
| trifloxystrobin | | x | s (G) | E | E | 1d | 4 |
| Fontelis (SC) | 7 | x | 14-20 fl. oz. | 14-20 fl. oz. | 14-20 fl. oz. | 12h | 61 fl. oz. |
| penthiopyrad | | x | G-E | F-G | F-G | 0d | NA |
| Indar 2F | 3 | x | 6 fl. oz. | 6 fl. oz. | x | 12h | 48 fl. oz. |
| fenbuconazole | | x | E[r] | F | x | 0d | 8 |
| Inspire Super (EW) | 3+9 | x | 16-20 fl. oz. | 16-20 fl. oz. | 16-20 fl. oz. | 12h | 80 fl. oz. |
| difenoconazole + cyprodinil | | x | E | F-G | F-G | 2d | 4 |
| Kenja 400 SC | 7 | x | 12.5 fl. oz. | 12.5 fl. oz. | x | 12h | 37.5 fl. oz. |
| isofetamid | | x | E | G | x | 1d | 3 |
| Kocide 3000 | M | 0.25-0.5 lb. | x | x | x | 48h | 60 lb. |
| copper hydroxide | | G-F | x | x | x | 0d | NA |
| Luna Experience (SC) | 7+3 | x | 6-10 fl. oz. | 8-10 fl. oz. | 6-10 fl. oz. | 12h | 34 fl. oz. |
| fluopyram + tebuconazol | | x | G-E | E | G | 0d | NA |
| Luna Privilege | 7 | x | 4-6.8 fl. oz. | 4.8-6.8 fl. oz. | 4-6.8 fl. oz. | 12h | 13.7 fl. oz. |
| fluopyram | | x | E | F | G | 0d | NA |
| Luna Sensation (SC) | 7+11 | x | 5-7.6 fl. oz. | 5-7.6 fl. oz. | 5-7.6 fl. oz. | 12h | 27.1 fl. oz. |
| fluopyram + trifloxystrobin | | x | E | F | G-E | 1d | 4 |
| Merivon XBF | 7+11 | x | 4-6.7 fl. oz. | 4-6.7 fl. oz. | 4-6.7 fl. oz. | 12h | 20.1 fl. oz. |
| fluxapyroxad + pyraclostrobin | | x | E | G-E | G-E | 0d | 3 |
| Microthiol Disperss | M | x | 10-20 lb. | x | 10-20 lb. | 24h | NA |
| sulfur | | x | F-P | x | F-P | 0d | NA |
| Miravis | 7 | x | 3.4-5.1 fl. oz. | 3.4-5.1 fl. oz. | 3.4-5.1 fl. oz. | 4h | 20.4 fl. oz. |
| pydiflumetofen | | x | E | E-G | E-G | 0d | 4 |
| Mycoshield | 41 | 12 oz./100 g. | x | x | x | 12h | 12 lb. |
| oxytetracycline | | E[r] | x | x | x | 21d | 8 |
| Ph-D | 19 | x | x | 6.2 oz. | 6.2 oz. | 4h | NA |
| polyoxin D | | x | x | G | u | 0d | NA |

(Continued)

Table 4-10. Peach Diseases - Summer Cover¹ (continued)

| Product And Formulation Active Ingredient | FRAC Code ² | Bacterial Spot/Shot Hole | Brown Rot | Peach Scab | Powdery Mildew/Peach Rusty Spot | REI ³ PHI ⁴ | Max Amt ⁵ Max App ⁶ |
|--|---------------------------|--------------------------------|-----------------|-----------------|---------------------------------------|--------------------------------------|--|
| Pristine | 7+11 | x | 10.5-14.5 oz. | 10.5-14.5 oz. | 10.5-14.5 oz. | 12h | 72.5 oz. |
| pyaclostrobin + boscalid | | x | G[r] | G-E | E[r] | 0d | 5 |
| Quadris Top | 11+3 | x | 12-14 fl. oz. | 12-14 fl. oz. | 12-14 fl. oz. | 12h | 56 fl. oz. |
| azoxystrobin + difenoconazole | | x | E | u | G | 0d | 4 |
| Quash | 3 | x | 2.5-3.5 fl. oz. | 2.5-3.5 fl. oz. | 3.5-4 fl. oz. | 12h | 12 fl. oz. |
| metconazole | | x | G-E | G | E | 14d | 3 |
| Quilt Xcel | 11+3 | x | 14 fl. oz. | 14 fl. oz. | 14 fl. oz. | 12h | 70 fl. oz. |
| azoxystrobin + propiconazole | | x | E | G | G | 0d | 5 |
| Quintec | 13 | x | x | x | 7 fl. oz. | 12h | 28 fl. oz. |
| quinoxifen | | x | x | x | E | 7d | 4 |
| Rally 40WSP | 3 | x | 2.5-6 oz. | x | 2.5-6 oz. | 24h | 3.25 lb. |
| myclobutanil | | x | G | x | G-E | 0h | NA |
| Scala (SC) | 9 | x | 9-18 fl. oz. | x | x | 12h | 54 fl. oz. |
| pyrimethanil | | x | G-E | x | x | 2d | 3 |
| Thiram Granuflo | M3 | x | 3.5 lb. | 3.5 lb. | x | 24h | 21.2 lb. |
| thiram | | x | G | G | x | 7d | NA |
| Tilt (EC) | 3 | x | 4 fl. oz. | x | 4 fl. oz. | 12h | 20 fl. oz. |
| propiconazole | | x | E | x | G | 0d | 5 |
| Topguard EQ | 3+11 | x | 6-8 fl. oz. | 6-8 fl. oz. | 6-8 fl. oz. | 12h | NA |
| flutriafol + azoxystrobin | | x | G | u | E | 7d | 4 |
| Topguard Specialty Crop | 3 | x | 14 fl. oz. | x | 14 fl. oz. | 12h | 56 fl. oz. |
| flutriafol | | x | E | s(G) | G | 7d | 4 |
| Topsin M WSB | 1 | x | 1-1.5 lb. | 1-1.5 lb. | 1-1.5 lb. | 48h | 4 lb. |
| thiophanate-methyl | | x | G[r] | G | G[r] | 1d | NA |
| Vanguard WG (75WG) | 9 | x | 5 oz. | x | x | 12h | 30 oz. |
| cyprodinil | | x | G-E | x | x | 2d | 4 |
| Ziram 76DF | M3 | x | 4.5-8 lb. | 4.5-8 lb. | x | 48h | 48.2 lb. |
| ziram | | x | G | G | x | 30d | 6 |

Footnotes: See how to read the spray schedule tables section at the beginning of the chapter.

Peach Summer Covers - Insects

Apply at 10- to 14-day intervals.

Table 4-11. Peach Insects Summer Cover¹

| Product And Formulation Active Ingredient | IRAC Code ² | Green June Beetle | Japanese Beetle | Oriental Fruit Moth | Plant Bug/ Stink Bug | REI ³ PHI ⁴ | Max Amt ⁵ Max App ⁶ |
|--|---------------------------|----------------------|--------------------|------------------------|-------------------------|--------------------------------------|--|
| Actara (25WDG) | 4A | x | x | x | 4.5-5.5 oz. | 12h | 11 oz. |
| thiamethoxam | | x | x | x | G | 14d | NA |
| Admire Pro (4.6F) | 4A | 1.4-2.8 fl. oz. | 1.4-2.8 fl. oz. | x | 2.8 fl. oz. | 12h | 8.4 fl. oz. |
| imidacloprid | | G | G | x | s | 0d | NA |

(Continued)

Table 4-11. Peach Insects - Summer Cover¹ (continued)

| Product And Formulation Active Ingredient | IRAC Code ² | Green June Beetle | Japanese Beetle | Oriental Fruit Moth | Plant Bug/ Stink Bug | REI ³ PHI ⁴ | Max Amt ⁵ Max App ⁶ |
|--|---------------------------|----------------------|----------------------|---------------------------|-------------------------|--------------------------------------|--|
| Altacor eVo | 28 | x | x | 1.5-2.2 oz. | x | 4h | 4.6 oz. |
| chlorantraniliprole | | x | x | E | x | 10d | 3 |
| Asana XL (0.66EC) (RUP) | 3A | x | x | 4.8-14.5 fl. oz. | 4.8-14.5 fl. oz. | 12h | 72.5 fl. oz. |
| esfenvalerate | | x | x | E[r] | G | 14d | NA |
| Assail 30SG | 4A | x | 5.3-8 oz. | 5.3 - 8 oz. | 5.3-8 oz. | 12h | 32 oz. |
| acetamiprid | | x | G | E | F | 7d | 4 |
| Avaunt eVo (30WDG) | 22 | x | x | 6 oz. | x | 12h | 24 oz. |
| indoxacarb | | x | x | G | x | 14d | 4 |
| Baythroid XL (1EC) (RUP) | 3A | x | x | 2-2.4 fl. oz. | 2-2.4 fl. oz. | 12h | 5.6 fl. oz. |
| beta-cyfluthrin | | x | x | E[r] | E | 7d | NA |
| BeetleGone! | 11 | 1-17.5 lb. | 1-17.5 lb. | x | x | 4h | NA |
| <i>B. thuringiensis</i> | | G | G | x | x | 0d | NA |
| Belay (2.13SC) | 4A | x | x | x | 6 fl. oz. | 12h | 12 fl. oz. |
| clothianidin | | x | x | x | E | 21d | NA |
| Beleaf 50SG | 29 | x | x | x | 2-2.8 oz. | 12h | 8.4 oz. |
| flonicamid | | x | x | x | G | 14d | 3 |
| Danitol 2.4EC (RUP) | 3A | x | 10.7-21.3 fl. oz | 10.7-21.3 fl. oz. | 10.7-21.3 fl. oz. | 24h | 42.7 fl. oz. |
| fenpropathrin | | x | E | E[r] | E | 3d | NA |
| Delegate WG (25WG) | 5 | x | x | 6-7 oz. | x | 4h | 28 oz. |
| spinetoram | | x | x | E | x | 1d | 4 |
| Diazinon AG 600WBC (RUP) | 1B | x | x | 12.75 fl. oz./100 gal. | x | 4d | 51 fl. oz. |
| diazinon | | x | x | G | x | 21d | 2 |
| Dimilin 2L (RUP) | 15 | x | x | 8-16 fl. oz. | x | 12h | 32 fl. oz. |
| diflubenzuron | | x | x | u | x | 14d | 2 |
| Entrust SC (2SC) | 5 | x | x | 4-8 fl. oz. | x | 4h | 29 fl. oz. |
| spinosad | | x | x | F | x | 7d | 3 |
| Esteem 35WP | 7C | x | x | 4-5 oz. | x | 12h | 15 oz. |
| pyriproxifen | | x | x | s | x | 14d | 3 |
| Exirel (0.83SE) | 28 | x | 13.5-20.5 fl. oz. | 10-20.5 fl. oz. | x | 12h | 61.5 fl. oz. |
| cyantraniliprole | | x | G | E | x | 3d | 3 |
| Imidan 70W | 1B | x | 2.1-4.25 lb. | 2.1-4.25 lb. | x | 4d | 17 lb. |
| phosmet | | x | G | E[r] | x | 14d | NA |
| Intrepid 2F | 18 | x | x | 10-16 fl. oz. | x | 4h | 64 fl. oz. |
| methoxyfenozide | | x | x | G | x | 7d | NA |
| Lannate LV | 1A | x | x | 3 pt. | 3 pt. | 4d | 18 pt. |
| methomyl | | x | x | F | G | 4d | 6 |
| Malathion 5EC | 1B | x | 2.5-4.8 pt. | 4.8 pt. | x | 24h | 4.8 pt. |
| malathion | | x | x | u | x | 7d | 3 |

(Continued)

Table 4-11. Peach Insects - Summer Cover¹ (continued)

| Product And Formulation Active Ingredient | IRAC Code ² | Green June Beetle | Japanese Beetle | Oriental Fruit Moth | Plant Bug/ Stink Bug | REI ³ PHI ⁴ | Max Amt ⁵ Max App ⁶ |
|--|---------------------------|----------------------|---------------------|------------------------|-------------------------|--------------------------------------|--|
| Mustang Maxx (0.83EC) (RUP) | 3A | x | x | 1.28-4 fl. oz. | 1.28-4 fl. oz. | 12h | 24 fl. oz. |
| zeta-cypermethrin | | x | x | E[r] | E | 14d | NA |
| Neemix 4.5 (0.39L) | UN | x | x | 7-16 fl. oz. | 7-16 fl. oz. | 4h | NA |
| azadirachtin | | x | x | u | E | 0d | NA |
| Pounce 25WP | 3A | x | x | 6.4-16 oz. | x | 12h | 48 oz. |
| permethrin | | x | x | E[r] | x | 14d | NA |
| Pyganic 5EC | 3A | x | 4.5-15.6 fl. oz. | x | 4.5-15.6 fl. oz. | 12h | 15.6 fl. oz. |
| pyrethrins | | x | u | x | u | 0d | 1 |
| Rimon 0.83EC | 15 | x | x | 20-40 fl. oz. | 20-40 fl. oz. | 12h | 150 fl. oz. |
| novaluron | | x | x | E | u | 8d | NA |
| Scorpion 35SL | 4A | x | x | x | 5.25-7 fl. oz. | 12h | 14.25 fl. oz. |
| dinotefuran | | x | x | x | E | 3/21d | NA |
| Sevin XLR Plus (4F) | 1A | x | 2-3 qt. | 2-3 qt. | 2-3 qt. | 12h | 14 qt. |
| carbaryl | | x | E | F | F | 3d | 3 |
| Surround WP (95WP) | UN | x | 25-50 lb. | 25-50 lb. | x | 4h | NA |
| kaolin | | x | s | s | x | 0d | NA |
| Venom (70SG) | 4A | x | x | x | 3-4 oz. | 12h | 6 oz. |
| dinotefuran | | x | x | x | E | 3d | NA |
| Verdepryn 100SL (0.83SL) | 28 | x | x | 5.5-11 fl. oz. | 5.5-11 fl. oz. | 4h | 33 fl. oz. |
| cyclaniliprole | | x | x | E | s | 7d | 3 |
| Warrior II (2.08CS) (RUP) | 3A | 1.2-2.5 fl. oz. | 1.2-2.5 fl. oz. | 1.2-2.5 fl. oz. | 1.2-2.5 fl. oz. | 24h | 12.8 fl. oz. |
| lambda-cyhalothrin | | E | E | G[r] | E | 14d | NA |

Footnotes: See how to read the spray schedule tables section at the beginning of the chapter.

Peach Preharvest - Diseases

Apply according to label directions beginning three weeks before harvest.

- Continued applications of fungicides may be necessary in orchards with moderate to high disease pressure (due to cultivar susceptibility, rainfall and/or fungicide resistance).

- Be aware of PHI of fungicides and bactericides (e.g., Mycoshield with PHI of 21 days).

Table 4-12. Peach Diseases - Preharvest¹

| Product And Formulation Active Ingredient | FRAC Code ² | Brown Rot | Peach Scab | Powdery Mildew/ Peach Rusty Spot | REI ³ PHI ⁴ | Max Amt ⁵ Max App ⁶ |
|--|---------------------------|-----------------|----------------|-------------------------------------|--------------------------------------|--|
| Abound (SC) | 11 | 12-15.5 fl. oz. | 9-15.5 fl. oz. | 12-15.5 fl. oz. | 4h | 90 fl. oz. |
| azoxystrobin | | F-G[r] | G | F | 0d | See label |
| Captan | M4 | 2.5-5 lb. | 2.5-5 lb. | 2.5-5 lb. | 24h | 40 lb. |
| captan | | G | G | F | 0d | NA |
| Cevya | 3 | 3-5 fl. oz. | 3-5 fl. oz. | 4-5 fl. oz. | 12h | 15 fl. oz. |
| mefentrifluconazole | | E | E | E | 0d | See label |

(Continued)

Table 4-12. Peach Diseases - Preharvest¹ (continued)

| Product And Formulation Active Ingredient | FRAC Code ² | Brown Rot | Peach Scab | Powdery Mildew/ Peach Rusty Spot | REI ³ PHI ⁴ | Max Amt ⁵ Max App ⁶ |
|--|---------------------------|-----------------|-----------------|-------------------------------------|--------------------------------------|--|
| Elevate 50 WDG | 17 | 1-1.5 lb. | x | x | 12h | 6 lb. |
| fenhexamid | | G-E | x | x | 0d | NA |
| Elite 45DF | 3 | 4-8 oz. | x | 4-8 oz. | 12h | 3 lb. |
| tebuconazole | | E | x | E | 0d | NA |
| Flint Extra | 11 | 2.5-3.8 fl. oz. | 2.5-3.8 fl. oz. | 2.5-3.8 fl. oz. | 12h | 15.2 fl. oz. |
| trifloxystrobin | | s (G) | E | E | 1d | 4 |
| Fontelis (SC) | 7 | 14-20 fl. oz. | 14-20 fl. oz. | 14-20 fl. oz. | 12h | 61 fl. oz. |
| penthiopyrad | | G-E | F-G | F-G | 0d | NA |
| Indar 2F | 3 | 6 fl. oz. | 6 fl. oz. | x | 12h | 48 fl. oz. |
| fenbuconazole | | E[r] | F | x | 0d | 8 |
| Inspire Super (EW) | 3+9 | 16-20 fl. oz. | 16-20 fl. oz. | 16-20 fl. oz. | 12h | 80 fl. oz. |
| difenoconazole + cyprodinil | | E | F-G | F-G | 2d | 4 |
| Kenja 400 SC | 7 | 12.5 fl. oz. | 12.5 fl. oz. | x | 12h | 37.5 fl. oz. |
| isofetamid | | E | G | x | 1d | 3 |
| Kocide 3000 | M | 3-5 lb. | x | x | 48h | 60 lb. |
| copper hydroxide | | F | x | x | 0d | NA |
| Luna Experience (SC) | 7+3 | 6-10 fl. oz. | 8-10 fl. oz. | 6-10 fl. oz. | 12h | 34 fl. oz. |
| fluopyram + tebuconazol | | G-E | E | G | 0d | NA |
| Luna Privilege | 7 | 4-6.8 fl. oz. | 4.8-6.8 fl. oz. | 4-6.84 fl. oz. | 12h | 13.7 fl. oz. |
| fluopyram | | E | F | G | 0d | NA |
| Luna Sensation (SC) | 7+11 | 5-7.6 fl. oz. | 5-7.6 fl. oz. | 5-7.6 fl. oz. | 12h | 27.1 fl. oz. |
| fluopyram + trifloxystrobin | | E | F | E-G | 1d | 4 |
| Merivon XBF | 7+11 | 4-6.7 fl. oz. | 4-6.7 fl. oz. | 4-6.7 fl. oz. | 12h | 20.1 fl. oz. |
| fluxapyroxad + pyraclostrobin | | E | E-G | E-G | 0d | 3 |
| Microthiol Disperss | M | 10-20 lb. | x | 10-20 lb. | 24h | NA |
| sulfur | | F-P | x | F-P | 0d | NA |
| Miravis | 7 | 3.4-5.1 fl. oz. | 3.4-5.1 fl. oz. | 3.4-5.1 fl. oz. | 4h | 20.4 fl. oz. |
| pydiflumetofen | | E | G-E | G-E | 0d | 4 |
| Ph-D | 19 | x | 6.2 oz. | 6.2 oz. | 4h | NA |
| polyoxin D | | x | G | u | 0d | NA |
| Pristine | 7+11 | 10.5-14.5 oz. | 10.5-14.5 oz. | 10.5-14.5 oz. | 12h | 72.5 oz. |
| pyraclostrobin + boscalid | | G[r] | G-E | E[r] | 0d | 5 |
| Quadris Top | 11+3 | 12-14 fl. oz. | 12-14 fl. oz. | 12-14 fl. oz. | 12h | 56 fl. oz. |
| azoxystrobin + difenoconazole | | E | u | G | 0d | 4 |
| Quash | 3 | 2.5-3.5 fl. oz. | 2.5-3.5 fl. oz. | 3.5-4 fl. oz. | 12h | 12 fl. oz. |
| metconazole | | E-G | G | E | 14d | 3 |
| Quilt Xcel | 11+3 | 14 fl. oz. | 14 fl. oz. | 14 fl. oz. | 12h | 70 fl. oz. |
| azoxystrobin + propiconazole | | E | G | G | 0d | 5 |
| Quintec | 13 | x | x | 7 fl. oz. | 12h | 28 fl. oz. |
| quinoxifen | | x | x | E | 7d | 4 |

(Continued)

Table 4-12. Peach Diseases - Preharvest¹ (continued)

| Product And Formulation Active Ingredient | FRAC Code ² | Brown Rot | Peach Scab | Powdery Mildew/ Peach Rusty Spot | REI ³ PHI ⁴ | Max Amt ⁵ Max App ⁶ |
|--|---------------------------|--------------|-------------|-------------------------------------|--------------------------------------|--|
| Rally 40WSP | 3 | 2.5-6 oz. | x | 2.5-6 oz. | 24h | 3.25 lb. |
| myclobutanil | | G | x | E-G | 0d | NA |
| Scala (SC) | 9 | 9-18 fl. oz. | x | x | 12h | 54 fl. oz. |
| pyrimethanil | | G-E | x | x | 2d | 3 |
| Thiram Granuflo | M3 | 3.5 lb. | 3.5 lb. | x | 24h | 21.2 lb. |
| thiram | | G | G | x | 7d | NA |
| Tilt (EC) | 3 | 4 fl. oz. | x | 4 fl. oz. | 12h | 20 fl. oz. |
| propiconazole | | E | x | G | 0d | 5 |
| Topguard EQ | 3+11 | 6-8 fl. oz. | 6-8 fl. oz. | 6-8 fl. oz. | 12h | NA |
| flutriafol + azoxystrobin | | G | u | E | 7d | 4 |
| Topguard Specialty Crop | 3 | 14 fl. oz. | x | 14 fl. oz. | 12h | 56 fl. oz. |
| flutriafol | | E | s(G) | G | 7d | 4 |
| Topsin M WSB | 1 | 1-1.5 lb. | 1-1.5 lb. | 1-1.5 lb. | 48h | 4 lb. |
| thiophanate-methyl | | G[r] | G | G[r] | 1d | NA |
| Vanguard WG (75WG) | 9 | 5 oz | x | x | 12h | 30 oz. |
| cyprodinil | | G-E | x | x | 2d | 4 |
| Ziram 76DF | M3 | 4.5-8 lb. | 4.5-8 lb. | x | 48h | 48.2 lb. |
| ziram | | G | G | x | 30d | 6 |

Footnotes: See how to read the spray schedule tables section at the beginning of the chapter.

Peach Preharvest - Insects

Apply any insecticides according to label directions beginning three weeks before harvest.

- If not using peachtree borer mating disruption, peach tree borer is best controlled by a trunk

drench at the time of peak moth flight, usually in early August. See Borers of Peach, Cherry, and Plum Trees, page 152.

Table 4-13. Peach Insects - Preharvest¹

| Product And Formulation Active Ingredient | IRAC Code ² | Borers | Japanese Beetle | Oriental Fruit Moth | Spotted-Wing Drosophila | REI ³ PHI ⁴ | Max Amt ⁵ Max App ⁶ |
|--|---------------------------|---------------------|--------------------|------------------------|----------------------------|--------------------------------------|--|
| Admire Pro (4.6F) | 4A | x | 1.4-2.8 fl. oz. | x | x | 12h | 8.4 fl. oz. |
| imidacloprid | | x | G | x | x | 0d | NA |
| Altacor eVo | 28 | 1.5-2.2 oz. | x | 1.5-2.2 oz. | x | 4h | 4.6 oz. |
| chlorantraniliprole | | u | x | E | x | 10d | 3 |
| Asana XL (0.66EC) (RUP) | 3A | 4.8-14.5 fl. oz. | x | 4.8-14.5 fl. oz. | x | 12h | 72.5 fl. oz. |
| esfenvalerate | | G | x | E[r] | x | 14d | NA |
| Assail 30SG | 4A | 5.3-8 oz. | 5.3-8 oz. | 5.3-8 oz. | x | 12h | 32 oz. |
| acetamiprid | | G | G | E | x | 7d | 4 |

(Continued)

Table 4-13. Peach Insects - Preharvest¹ (continued)

| Product And Formulation Active Ingredient | IRAC Code ² | Borers | Japanese Beetle | Oriental Fruit Moth | Spotted-Wing Drosophila | REI ³ PHI ⁴ | Max Amt ⁵ Max App ⁶ |
|--|---------------------------|------------------|--------------------|------------------------|----------------------------|--------------------------------------|--|
| Avaunt eVo (30WDG) | 22 | x | x | 6 oz. | x | 12h | 24 oz. |
| indoxacarb | | x | x | G | x | 14d | 4 |
| Baythroid XL (IEC) (RUP) | 3A | 1.4-2 fl. oz. | x | 2-2.4 fl. oz. | x | 12h | 5.6 fl. oz. |
| beta-cyfluthrin | | G | x | E[r] | x | 7d | NA |
| BeetleGone! | 11 | x | 1-17.5 lb. | x | x | 4h | NA |
| <i>B. thuringiensis</i> | | x | G | x | x | 0d | NA |
| Danitol 2.4EC (RUP) | 3A | x | 10.7-21.3 fl. oz. | 10.7-21.3 fl. oz. | 10.7-21.3 fl. oz. | 24h | 42.7 fl. oz. |
| fenpropathrin | | x | E | E[r] | E | 3d | NA |
| Delegate WG (25WG) | 5 | x | x | 6-7 oz. | 4.5-7 oz. | 4h | 28 oz. |
| spinetoram | | x | x | E | G | 1d | 4 |
| Dimilin 2L (RUP) | 15 | x | x | 8-16 fl. oz. | x | 12h | 32 fl. oz. |
| diflubenzuron | | x | x | u | x | 14d | 2 |
| Entrust SC (2SC) | 5 | x | x | 4-8 fl. oz. | x | 4h | 29 fl. oz. |
| spinosad | | x | x | F | x | 7d | 3 |
| Esteem 35WP | 7C | x | x | 4-5 oz. | x | 12h | 15 oz. |
| pyriproxifen | | x | x | s | x | 14d | 3 |
| Exirel (0.83SE) | 28 | x | 13.5-20.5 fl. oz. | 10-20.5 fl. oz. | 13.5-20.5 fl. oz. | 12h | 61.5 fl. oz. |
| cyantraniliprole | | x | G | E | E | 3d | 3 |
| Imidan 70W | 1B | x | 2.1-4.25 lb. | 2.1-4.25 lb. | 2.1-4.25 lb. | 4d | 17 lb. |
| phosmet | | x | G | E[r] | E | 14d | NA |
| Intrepid 2F | 18 | x | x | 10-16 fl. oz. | x | 4h | 64 fl. oz. |
| methoxyfenozide | | x | x | G | x | 7d | NA |
| Lannate LV | 1A | x | x | 3 pt. | x | 4d | 18 pt. |
| methomyl | | x | x | F | x | 4d | 6 |
| Malathion 5EC | 1B | x | 2.5-4.8 pt. | 4.8 pt. | x | 24h | 4.8 pt. |
| malathion | | x | u | u | x | 7d | 3 |
| Movento (2SC) | 23 | x | x | x | 6-9 fl. oz. | 24h | 15.3 fl. oz. |
| spirotetramat | | x | x | x | s | 7d | NA |
| Mustang Maxx (0.83EC) (RUP) | 3A | 1.28-4 fl. oz. | x | 1.28-4 fl. oz. | 4 fl. oz. | 12h | 24 fl. oz. |
| zeta-cypermethrin | | G | x | E[r] | E | 14d | NA |
| Neemix 4.5 (0.39L) | UN | 7-16 fl. oz. | x | 7-16 fl. oz. | x | 4h | NA |
| azadirachtin | | F | x | u | x | 0d | NA |
| Pounce 25WP | 3A | 6.4-16 oz. | x | 6.4-16 oz. | x | 12h | 48 oz. |
| permethrin | | F | x | E[r] | x | 14d | NA |
| Pyganic 5EC | 3A | 4.5-15.6 fl. oz. | 4.5-15.6 fl. oz. | x | 4.5-15.6 fl. oz. | 12h | 15.6 fl. oz. |
| pyrethrins | | u | u | x | F | 0d | 1 |
| Rimon 0.83EC | 15 | 20 fl. oz. | x | 20-40 fl. oz. | 20-40 fl. oz. | 12h | 150 fl. oz. |
| novaluron | | u | x | E | G | 8d | NA |

(Continued)

Table 4-13. Peach Insects - Preharvest¹ (continued)

| Product And Formulation Active Ingredient | IRAC Code ² | Borers | Japanese Beetle | Oriental Fruit Moth | Spotted-Wing Drosophila | REI ³ PHI ⁴ | Max Amt ⁵ Max App ⁶ |
|--|---------------------------|-----------------|--------------------|------------------------|----------------------------|--------------------------------------|--|
| Sevin XLR Plus (4F) | 1A | 2-3 qt. | 2-3 qt. | 2-3 qt. | x | 12h | 14 qt. |
| carbaryl | | u | E | F | x | 3d | 3 |
| Surround WP (95WP) | UN | x | 25-50 lb. | 25-50 lb. | x | 4h | NA |
| kaolin | | x | s | s | x | 0d | NA |
| Venom (70SG) | 4A | 3-4 oz. | x | x | x | 12h | 6 oz. |
| dinotefuran | | u | x | x | x | 3d | NA |
| Verdepryn 100SL (0.83SL) | 28 | x | x | 5.5-11 fl. oz. | 5.5-11 fl. oz. | 4h | 33 fl. oz. |
| cyclaniliprole | | x | x | E | u | 7d | 3 |
| Warrior II (2.08CS) (RUP) | 3A | 1.2-2.5 fl. oz. | 1.2-2.5 fl. oz. | 1.2-2.5 fl. oz. | x | 24h | 12.8 fl. oz. |
| lambda-cyhalothrin | | G | E | G[r] | x | 14d | NA |

Footnotes: See how to read the spray schedule tables section at the beginning of the chapter.

Special Notes on Peach Schedule

Spotted lanternfly

The spotted lanternfly is an invasive planthopper that has spread throughout the Midwest. This insect feeds on plant sap, causing wilting, dieback and even death.

Currently spotted lanternfly is believed to pose the greatest threat to the blueberry, grape, hops, stone fruit and hardwood industries. Know how to identify this pest and remain vigilant for its appearance in your orchard and vineyard systems.

5. PLUM

Plum Insect Pests

The shaded boxes represent the crop stages where common pests in the Midwest are active and action (scouting and preventative sprays) may be necessary/recommended. Weather and degree day accumulation will impact the exact timing of pest appearance in the orchard.

| Dormant | Pre-bloom | Full Bloom | Petal Fall | Shuck Split | First Cover | Second And Additional Cover | Pre-Harvest |
|-------------------|-----------|------------|---------------------------|-------------|-------------------|-----------------------------|-------------------------------|
| | | | | | | Japanese Beetle | |
| | OFM-MD | | Oriental Fruit Moth (OFM) | | | | |
| | PTB-MD | | Peachtree Borer | | | | |
| | | | Plum Curculio | | | | |
| European Red Mite | | | | | | | |
| | | | | | Periodical Cicada | | |
| San Jose Scale | | | | | | | |
| | | | | | | | Spotted Wing Drosophila (SWD) |

Plum Spray Schedule

How to read the spray schedule tables

Every plum growth stage has important notes on disease or insect management. In some cases, the reader will be directed to the special problems section at the end of the section or chapter. Please make sure to read thoroughly and contact your state Extension specialist with any specific questions.

Key to tables

E = excellent control

G = good control

F = fair control

[r] = fungicide/insecticide resistance possible

s = suppression only

i = ineffective

u = unknown efficacy

x = pest not on the label

¹ Efficacy data in this publication are based on trials conducted across various regions and does not necessarily reflect local efficacy differences or changes over time. Growers should contact their Extension specialist for the most recent or for state-specific information. The information on this publication is only a guide; the

authors and their institutions assume no liability for practices implemented based on this information. Always read and follow pesticide labels. The label is the law. Product registration may vary by state.

² FRAC/IRAC code represents the mode of action of the fungicide/insecticide.

³ PHI refers to the pre-harvest interval, which is the number of days before harvest that the product may not be applied.

⁴ All fungicides/insecticides have a Restricted-Entry Interval (REI). The restricted-entry interval is the time immediately after a pesticide application when entry into the treated area is limited. Check labels for REI. Restrictions in REI may prohibit the use of certain pesticides during harvest.

Applicators must abide by both maximum amount of product per season AND maximum number of applications.

⁵ Max amt refers to the product's maximum amount/acre/year.

⁶ Max app refers to the product's maximum number of applications per year.

RUP refers to restricted use pesticide.

Plum Dormant - Diseases

Notes on disease management

- Bacterial spot and copper pesticides:** Plum is on most, but not all, copper fungicide labels. Check label before use. Using copper at the dormant stage may also reduce the overwintering inoculum of the bacterium that cause bacterial canker, but have limited efficacy in suppressing bacterial spot, plum pockets and black knot. As season progresses, reduce the rate of copper applied to reduce the risk of phytotoxicity. Copper applied with tanks having a pH of less than 6.5 may cause phytotoxicity issues.
- Black knot:** Prune out and destroy knots during the dormant season with pruning cuts at least 8" below knots. Continue knot removal all season long, whenever they are observed. Remove nearby wild plums and cherry seedlings.
- Cytospora canker:** Paint trunks with whitewash to prevent winter injury and sunscald.
- Plum pockets (leaf curl):** Autumn application during leaf fall is the best time for application, followed by spring application at budswell. Unless disease pressure is severe, Luna Experience and Luna Sensation are better deployed later in the season for control of brown rot and powdery mildew.

Table 5-1. Plum Diseases - Dormant¹

| Product And Formulation Active Ingredient | FRAC Code ² | Bacterial Spot | Black Knot | Plum Pockets | REI ³ PHI ⁴ | Max Amt ⁵ Max App ⁶ |
|--|------------------------|----------------|--------------|--------------|--------------------------------------|--|
| Badge SC | M | 1.5-5 pt. | 1.5-5 pt. | 1.5-5 pt. | 24h | 18 lb. |
| copper sulfate + oxychloride | | F-G | x | s | 0d | NA |
| Bravo Weather Stik | M3 | 3-4 pt. | 3-4 pt. | 3-4 pt. | 12h | 18.8 lb. |
| chlorothalonil | | G | G | G | shuck-split | NA |
| C-O-C-S WDG | M | 1-2.9 lb. | x | 1-2.9 lb. | 48h | 35 lb. |
| copper oxychloride | | F-G | x | G | 21d | 3 |
| Champ FL | M | 5.3-8 pt. | x | 5.3-8 pt. | 48 h | 49.6 pt. |
| copper hydroxide | | s | x | s | NA | NA |
| Cuprofix Ultra 40 disperss | M | 5-8 lb. | 3-3.75 | 5-7.5 lb. | 48h | 45 lb. |
| copper sulfate | | F-G | F | G | 120d | NA |
| Kocide 3000 | M | 3.5-7 lb. | 1.75-3.5 lb. | 1.75-3.5 lb. | 48h | 60 lb. |
| copper hydroxide | | F-G | F | F-G | 0d | 4 |

Footnotes: See how to read the spray schedule tables section at the beginning of the chapter.

Plum Dormant - Insects

Notes on insect management

- European red mite and scale insects:** Apply dormant oil at a rate of 2 gal. per 100 gals. (2%). Check labels carefully for temperature restrictions (e.g., when temperatures are above 40F° or never during freezing weather).

Plum Pre-bloom - Diseases

Notes on disease management

- Copper pesticides:** Copper rates are tied to crop development, with rates reduced as the season progresses to minimize the risk of phytotoxicity. Do not apply during cooler conditions with extended dews or fog.
 Note: Copper has limited efficacy for control of bacterial spot. For more information see Bacterial Spot of Peach, Nectarine and Plum on pages 152-153.

Table 5-2. Plum Diseases Bloom¹

| Product And Formulation Active Ingredient | FRAC Code ² | Bacterial Spot | Black Knot | Plum Pockets | REI ³ PHI ⁴ | Max Amt ⁵ Max App ⁶ |
|--|------------------------|-------------------|--------------|------------------|--------------------------------------|--|
| Badge SC | M | 1.5-5 pt. | 1.5-5 pt. | 1.5-5 pt. | 24h | 18 lb. |
| copper sulfate + oxychloride | | F-G | F | s | 0d | NA |
| Bravo Weather Stik | M3 | 3-4 pt. | 3-4 pt. | 3-4 pt. | 12h | 18.8 lb. |
| chlorothalonil | | G | G | G | shuck-split | NA |
| C-O-C-S WDG | M | 1-2.9 lb. | x | x | 48h | 35 lb. |
| copper oxychloride | | F-G | x | x | 21d | 3 |
| Captan 80WDG | M | x | 3.75 lb. | 3.75 lb. | 24h | 40 lb. |
| captan | | x | s | F | 0d | NA |
| Champ FL | M | 4.2 pt. | x | x | 48 h | 49.6 pt. |
| copper hydroxide | | s | x | S | NA | NA |
| Cuprofix Ultra 40 disperss | M | 1-2.5 lb. | 3-3.75 | 5-7.5 lb. | 48h | 45 lb. |
| copper sulfate | | F-G | F | G | 120d | NA |
| Kocide 3000 | M | 0.25-0.5 lb. | 1.75-3.5 lb. | 1.75-3.5 lb. | 48h | 60 lb. |
| copper hydroxide | | F-G | F | F-G | 0d | 4 |
| Luna Experience (SC) | 7+3 | x | x | 6-10 fl. oz. | 12 h | 34 fl. oz. |
| fluopyram + tebuconazol | | x | x | u | 1d | varies |
| Luna Sensation (SC) | 7+11 | x | x | 5 to 7.6 fl. oz. | 12h | 271 fl. oz. |
| fluopyram + trifloxystrobin | | x | x | E | 1d | 4 |
| Pristine | 7+11 | x | x | 10.5-14.5 oz. | 12h | 72.5 oz. |
| pyaclostrobin + boscalid | | x | x | s | 0d | 9 |
| Topguard Specialty Crop | 3 | x | x | 13 fl. oz. | 12h | 56 fl. oz. |
| flutriafol | | x | x | G | 7d | 4 |

Footnotes: See how to read the spray schedule tables section at the beginning of the chapter.

Plum Pre-bloom - Insects

Notes on insect management

- **Oriental fruit moth and peachtree borer:** Pheromone traps for oriental fruit moth and peachtree borer should be deployed pre bloom. For mating disruption, see Mating Disruption for Peach Pests page 153.

Table 5-3. Plum Insects - Prebloom¹

| Product And Formulation Active Ingredient | IRAC Code ² | Aphids | European Red Mite | Leafroller | San Jose Scale | REI ³ PHI ⁴ | Max Amt ⁵ Max App ⁶ |
|--|------------------------|---------|----------------------|------------|-------------------|--------------------------------------|--|
| Acramite 50WS | 20D | x | 1 lb. | x | x | 12h | 1 lb. |
| bifenazate | | x | G | x | x | 3d | 1 |
| Actara (25WDG) | 4A | 3-4 oz. | x | x | x | 12h | 11 oz. |
| thiamethoxam | | E | x | x | x | 14d | NA |
| Agri-Mek SC (0.7SC) (RUP) | 6 | x | 2.25-4.25 fl. oz. | x | x | 12h | 8.5 fl. oz. |
| abamectin | | x | G | x | x | 21d | 2 |

(Continued)

Table 5-3. Plum Insects - Prebloom¹ (continued)

| Product And Formulation Active Ingredient | IRAC Code ² | Aphids | European Red Mite | Leafroller | San Jose Scale | REI ³ PHI ⁴ | Max Amt ⁵ Max App ⁶ |
|---|------------------------|---------------------------|-------------------------------|-------------------|---------------------------|--------------------------------------|--|
| Altacor (35WDG) | 28 | x | x | 3-4.5 oz. | x | 4h | 9 oz. |
| chlorantraniliprole | | x | x | E | x | 10d | NA |
| Apta (1.34SC) | 21A | 17-21 fl. oz. | x | 21-27 fl. oz. | x | 12h | 53.5 fl. oz. |
| tolfenpyrad | | u | x | G | x | 14d | 2 |
| Asana XL (0.66EC) (RUP) | 3A | 4.8-14.5 fl. oz. | x | 4.8-14.5 fl. oz. | x | 12h | 72.5 fl. oz. |
| esfenvalerate | | u | x | E | x | 14d | NA |
| Assail 30SG | 4A | 2.5-5.3 oz. | x | x | 5.3-8 oz. | 12h | 32 oz. |
| acetamiprid | | E | x | x | F | 7d | 4 |
| Bacillus thuringiensis (B.t.) (Dipel DF, etc.) | 11A | x | x | 0.5-2 lb. | x | 4h | NA |
| Bacillus thuringiensis | | x | x | u | x | 0d | NA |
| Baythroid XL (1EC) (RUP) | 3A | 2.4-2.8 fl. oz. | x | 2.4-2.8 fl. oz. | x | 12h | 5.6 fl. oz. |
| beta-cyfluthrin | | E | x | u | x | 7d | NA |
| Beleaf 50SG | 29 | 2-2.8 oz. | x | x | x | 12h | 8.4 fl. oz. |
| flonicamid | | E | x | x | x | 14d | 3 |
| Centaur WDG (70WDG) | 16 | x | x | x | 34.5 oz. | 12h | 69 oz. |
| buprofezin | | x | x | x | E | 14d | 2 |
| Closer SC (2SC) | 4C | 1.5-2.75 fl. oz. | x | x | 5.7 fl. oz. | 12h | 17 fl. oz. |
| sulfoxaflor | | E | x | x | s | 7d | 4 |
| Danitol 2.4EC (RUP) | 3A | 10.3-21.3 fl. oz. | 10.3-21.3 fl. oz. | 10.3-21.3 fl. oz. | x | 24h | 42.7 fl. oz. |
| fenpropathrin | | u | F | u | x | 3d | NA |
| Delegate WG (25WG) | 5 | x | x | 4.5-7 oz. | x | 4h | 28 oz. |
| spinetoram | | x | x | E | x | 1d | 4 |
| Diazinon AG600 WBC (RUP) | 1B | 12.75 fl. oz./100 gal. | 6.5-12.25 fl. oz./100 gal. | x | 12.75 fl. oz./100 gal. | 4d | 102 fl. oz. |
| diazinon | | u | u | x | F | 21d | 2 |
| Dimilin 2L (RUP) | 15 | x | x | 8-16 fl. oz. | x | 12h | 32 fl. oz. |
| diflubenzuron | | x | x | u | x | 14d | 2 |
| Entrust SC (2SC) | 5 | x | x | 4-8 fl. oz. | x | 4h | 29 fl. oz. |
| spinosad | | x | x | u | x | 1d | 3 |
| Envidor 2SC | 23 | x | 16-18 fl. oz. | x | x | 12h | 18 fl. oz. |
| spirodiclofen | | x | E | x | x | 7d | 1 |
| Esteem 35WP | 7C | x | x | x | 4-5 oz. | 12h | 15 oz. |
| pyriproxyfen | | x | x | x | E | 14d | 3 |
| Exirel (0.83SE) | 28 | 13.5-20.5 fl. oz. | x | 10-20.5 fl. oz. | x | 12h | 61.5 fl. oz. |
| cyantraniliprole | | E | x | E | x | 3d | 3 |
| Imidan 70W | 1B | x | x | 2.13-4.25 lb. | 2.13-4.25 lb. | 7d | 13 lb. |
| phosmet | | x | x | E | E | 7d | NA |
| Intrepid 2F | 18 | x | x | 8-16 fl. oz. | x | 4h | 64 fl. oz. |
| methoxyfenozide | | x | x | E | x | 7d | NA |

(Continued)

Table 5-3. Plum Insects - Prebloom¹ (continued)

| Product And Formulation Active Ingredient | IRAC Code ² | Aphids | European Red Mite | Leafroller | San Jose Scale | REI ³ PHI ⁴ | Max Amt ⁵ Max App ⁶ |
|--|------------------------|------------------|----------------------|------------------|-------------------|--------------------------------------|--|
| Magister SC (1.7SC) | 21A | x | 32-36 fl. oz. | x | x | 12h | 36 fl. oz. |
| fenazaquin | | x | E | x | x | 3d | 1 |
| Mustang Maxx (0.83EC) (RUP) | 3A | x | x | 1.28-4 fl. oz. | x | 12h | 24 fl. oz. |
| zeta-cypermethrin | | x | x | E | x | 14d | NA |
| Nexter (75WP) | 21 | x | 4.4-5.2 oz. | x | x | 12h | 21.3 oz. |
| pyridaben | | x | G | x | x | 7d | 2 |
| Oil (superior) | UN | x | 1.5-2% | x | 1.5-2% | 4h | NA |
| mineral oil | | x | E | x | G | 0d | NA |
| Onager (1EC) | 10A | x | 12-24 oz. | x | x | 12h | 24 fl. oz. |
| hexythiazox | | x | E | x | x | 7d | 1 |
| Portal XLO (0.4EC) | 21A | x | 2 pt. | x | x | 12h | 4 pt. |
| fenpyroximate | | x | E | x | x | 7d | 2 |
| PQZ (1.87SC) | 9B | 2.4-3.2 fl. oz. | x | x | x | 12h | 4.8 fl. oz. |
| pyrifluquinazon | | E | x | x | x | 7h | 2 |
| Proaxis (0.5EC) (RUP) | 3A | 2.5-5.1 fl. oz. | x | 2.5-5.1 fl. oz. | x | 24h | 1.6 pt. |
| gamma-cyhalothrin | | u | x | E | x | 14d | NA |
| Pyganic 5EC | 3A | 4.5-15.6 fl. oz. | 4.5-15.6 fl. oz. | 4.5-15.6 fl. oz. | 4.5-15.6 fl. oz. | 12h | NA |
| pyrethrins | | u | u | u | u | 0d | 10 |
| Rimon 0.83EC | 15 | x | x | 20-50 fl. oz. | x | 12h | 150 fl. oz. |
| novaluron | | x | x | E | x | 8d | NA |
| Savey 50DF | 10A | x | 3-6 oz. | x | x | 12h | 6 oz. |
| hexythiazox | | x | E | x | x | 28d | 1 |
| Sevin XLR Plus (4F) | 1A | 2-3 qt. | x | 2-3 qt. | 4-5 qt. | 12h | 14 qt. |
| carbaryl | | u | x | F | u | 3d | 3 |
| Sivanto Prime (1.67SC) | 4D | 7-14 fl. oz. | x | x | 10.5-14 fl. oz. | 4h | 28 fl. oz. |
| flupyradifurone | | E | x | x | G | 14d | NA |
| Surround WP (95WP) | UN | x | x | 25-50 lb. | x | 4h | NA |
| kaolin | | x | x | u | x | 0d | NA |
| Vendex 50WP (RUP) | 12B | x | 1-2 lb. | x | x | 2d | 3 lb. |
| fenbutatin-oxide | | x | G | x | x | 14d | 2 |
| Verdepryn 100SL (0.83SL) | 28 | x | x | 5.5-11 fl. oz. | x | 4h | 33 fl. oz. |
| cyclaniliprole | | x | x | E | x | 7d | 3 |
| Versys Inscalis (0.83DC) | 9D | 1.5 fl. oz. | x | x | x | 12h | 3 fl. oz. |
| afidopyropen | | G | x | x | x | 7d | NA |
| Warrior II (2.08CS) (RUP) | 3A | 1.2-2.5 fl. oz. | x | 1.2-2.5 fl. oz. | x | 24h | 12.8 fl. oz. |
| lambda-cyhalothrin | | u | x | u | x | 14d | NA |
| Zeal (72WP) | 10B | x | 2-3 oz. | x | x | 12h | 3 oz. |
| etoxazole | | x | E | x | x | 7d | 1 |

Footnotes: See how to read the spray schedule tables section at the beginning of the chapter.

Plum Full Bloom - Diseases

Notes on disease management

- Copper pesticides: Copper is not recommended after early bloom, to protect both flowers and bees.

Table 5-4. Plum Diseases - Full Bloom¹

| Product And Formulation Active Ingredient | FRAC Code ² | Brown Rot | Bacterial Spot | Black Knot | REI ³ PHI ⁴ | Max Amt ⁵ Max App ⁶ |
|--|------------------------|-----------------|----------------|--------------|--------------------------------------|--|
| Abound (SC) | 11 | 12-15.5 fl. oz. | x | x | 4h | 92.3 fl. oz. |
| azoxystrobin | | F-E[r] | x | x | 0d | 15 |
| Badge SC | M | 3.5-5 pt. | 1.5-5 pt. | 1.5-5 pt. | 24h | 18 lb. |
| copper sulfate + oxychloride | | F | F-G | s | 0d | NA |
| Bravo Weather Stik | M3 | 3.1-4.1 pt. | x | 3-4 pt. | 12h | 18.8 lb. |
| chlorothalonil | | G | G | G | shuck split | NA |
| C-O-C-S WDG | M | 1-2.9 lb. | 1-2.9 lb. | x | 48h | 35 lb. |
| copper oxychloride | | F | F-G | x | 21d | 3 |
| Captan 80WDG | M | 3.7 lb. | x | x | 24h | 40 lb. |
| captan | | G | x | s | 0d | NA |
| Cevya | 3 | 5 oz. | x | x | 12h | 15 fl. oz. |
| mefentrifluconazole | | E | x | x | 0d | 5 |
| Champ FL | M | 4.2 pt. | 4.2 pt. | 4.2 pt. | 48 h | 49.6 pt. |
| copper hydroxide | | F | s | s | NA | NA |
| Cuprofix Ultra 40 Disperss | M | 3.75 lb. | 1-2.5 lb. | 3-3.75 | 48h | 45 lb. |
| copper sulfate | | F | F-G | F | 120d | NA |
| Elevate 50 WDG | 17 | 1-1.5 lb. | x | x | 12h | 6 lb. |
| fenhexamid | | G-E | x | x | 0d | NA |
| Elite 45DF | 3 | 2 oz. | x | x | 12h | 3 lb. |
| tebuconazole | | E | x | x | 0d | NA |
| Flint Extra | 11 | 2.5-3.8 oz. | x | x | 12h | 15.2 fl. oz. |
| trifloxystrobin | | G-E | x | x | 1d | 3 |
| Fontelis (SC) | 7 | 14-20 fl. oz. | x | x | 12h | 61 fl. oz. |
| penthiopyrad | | E-G | x | x | 0d | 4 |
| Indar 2F | 3 | 6 fl. oz. | x | x | 12h | 48 fl. oz. |
| fenbuconazole | | E[r] | x | x | 0d | NA |
| Inspire Super (EW) | 3+9 | 16-20 fl. oz. | x | x | 12h | 80 fl. oz. |
| difenoconazole + cyprodinil | | E | x | x | 2d | 8 |
| Kenja 400 SC | 7 | 12.5 fl. oz. | x | x | 12h | 37.5 fl. oz. |
| isofetamid | | E | x | x | 1 day | 3 |
| Kocide 3000 | M | 3.5-5 lb. | 0.25-0.5 lb. | 1.75-3.5 lb. | 48h | 60 lb. |
| copper hydroxide | | F | F-G | s | 0d | 4 |
| Luna Experience (SC) | 7+3 | 6-10 oz. | x | x | 12h | 34 fl. oz. |
| fluopyram + tebuconazol | | G-E | x | x | 1d | varies |

(Continued)

Table 5-4. Plum Diseases - Full Bloom¹ (continued)

| Product And Formulation Active Ingredient | FRAC Code ² | Brown Rot | Bacterial Spot | Black Knot | REI ³ PHI ⁴ | Max Amt ⁵ Max App ⁶ |
|--|------------------------|-----------------|-------------------|------------|--------------------------------------|--|
| Luna Privilege | 7 | 4-6.8 fl. oz. | x | x | 12h | 13.7 fl. oz. |
| fluopyram | | E | x | x | 0d | 6 |
| Luna Sensation (SC) | 7+11 | 6-10 fl. oz. | x | x | 12h | 271 fl. oz. |
| fluopyram + trifloxystrobin | | E | x | x | 1d | 4 |
| Merivon XBF | 7+11 | 4-6.7 fl. oz. | x | x | 12h | 20.1 fl. oz. |
| fluxapyroxad + pyraclostrobin | | E | x | x | 0d | 4 |
| Microthiol Disperss | M | 10-20 lb. | x | x | 24h | NA |
| sulfur | | F | x | x | 0d | 3 |
| Miravis | 7 | 3.4-5.1 fl. oz. | x | x | 4 hr | 20.4 fl. oz. |
| pydiflumetofen | | G-E | x | x | 0d | 4 |
| Ph-D | 19 | 6.2 oz. | x | x | 0d | NA |
| polyoxin D | | G | x | x | NA | NA |
| Pristine | 7+11 | 10.5-14.5 oz. | x | x | 12h | 72.5 oz. |
| pyraclostrobin + boscalid | | G-E[r] | x | x | 0d | 9 |
| Quadris Top | 11+3 | 12-14 fl. oz. | x | x | 12h | 56 fl. oz. |
| azoxystrobin + difenoconazole | | E | x | x | 0d | 5 |
| Quash | 3 | 2.5-3.5 oz. | x | x | 12h | 12 oz. |
| metconazole | | E-G | x | x | 14d | 3 |
| Quilt Xcel | 11+3 | 14 fl. oz. | x | x | 12h | 70 fl. oz. |
| azoxystrobin + propiconazole | | E | x | x | 0d | 5 |
| Rally 40WSP | 3 | 2.5-6 oz. | x | x | 24h | 2.75 lb. |
| myclobutanil | | G | x | x | 1d | NA |
| Rovral 4F | 2 | 1-2 pt. | x | x | 24h | 4 pt. |
| iprodione | | E | x | x | N/A | NA |
| Scala (SC) | 9 | 9-18 fl. oz. | x | x | 12h | 54 fl. oz. |
| pyrimethanil | | G-E | x | x | 2d | 2 |
| Tilt (EC) | 3 | 4 fl. oz. | x | x | 12h | 20 fl. oz. |
| propiconazole | | E | x | x | 0d | 5 |
| Topguard EQ | 3+11 | 6-8 oz. | x | x | 12h | NA |
| flutriafol + azoxystrobin | | G | x | x | 7d | 4 |
| Topguard Specialty Crop | 3 | 13 fl. oz. | x | x | 12h | 56 fl. oz. |
| flutriafol | | E | x | x | 7d | 4 |
| Topsin M WSB | 1 | 1-1.5 lb. | x | 1-1.5 lb. | 48h | 4 lb. |
| thiophanate-methyl | | E[r] | x | E[r] | 1d | NA |
| Vanguard WG (75WG) | 9 | 5 oz. | x | x | 12h | 30 oz. |
| cyprodinil | | G-E | x | x | 2d | 4 |

Footnotes: See how to read the spray schedule tables section at the beginning of the chapter.

Plum Full Bloom - Insects

Notes on insect management

- Protect pollinators! Do not apply insecticides during bloom.

Plum Petal Fall To Shuck Split - Diseases

Notes on disease management

Brown rot

- Failure to control plum curculio and/or brown marmorated stink bug may result in an increase in brown rot.

- When using coppers or captan post-bloom, drying conditions should be excellent to minimize the risk of phytotoxicity. Applications of captan from shuck split through early July can cause shot-holing of leaves and spotting of fruit of some European and Japanese plums.
- Rovral cannot be applied after petal fall.
- The last application of Bravo is at shuck split.

Table 5-5. Plum Diseases - Petal Fall Through Shuck Split¹

| Product And Formulation Active Ingredient | FRAC Code ² | Brown Rot | Bacterial Spot | Black Knot | REI ³ PHI ⁴ | Max Amt ⁵ Max App ⁶ |
|--|------------------------|-----------------|----------------|------------|--------------------------------------|--|
| Abound (SC) | 11 | 12-15.5 fl. oz. | x | x | 4h | 92.3 fl. oz. |
| azoxystrobin | | F-E[r] | x | x | 0d | 15 |
| Bravo Weather Stik | M3 | 3.1-4.1 pt. | x | 3-4 pt. | 12h | 18.8 lb. |
| chlorothalonil | | G | G | G | shuck split | NA |
| Captan 80WDG | M | 3.7 lb. | x | x | 24h | 40 lb. |
| captan | | G | x | x | 0d | NA |
| Cevya | 3 | 5 oz. | x | x | 12h | 15 fl. oz. |
| mefentrifluconazole | | E | x | x | 0d | 5 |
| Elevate 50 WDG | 17 | 1-1.5 lb. | x | x | 12h | 6 lb. |
| fenhexamid | | G-E | x | x | 0d | NA |
| Elite 45DF | 3 | 2 oz. | x | x | 12h | 3 lb. |
| tebuconazole | | E | x | x | 0d | NA |
| Flint Extra | 11 | 2.5-3.8 fl. oz. | x | x | 12h | 15.2 fl. oz. |
| trifloxystrobin | | G-E | x | x | 1d | 3 |
| Fontelis (SC) | 7 | 14-20 fl. oz. | x | x | 12h | 61 fl. oz. |
| penthiopyrad | | E-G | x | x | 0d | 4 |
| Indar 2F | 3 | 6 fl. oz. | x | x | 12h | 48 fl. oz. |
| fenbuconazole | | E[r] | x | x | 0d | NA |
| Inspire Super (EW) | 3+9 | 16-20 fl. oz. | x | x | 12h | 80 fl. oz. |
| difenoconazole + cyprodinil | | E | x | x | 2d | 8 |
| Kenja 400 SC | 7 | 12.5 fl. oz. | x | x | 12h | 37.5 fl. oz. |
| isofetamid | | E | x | x | 1 day | 3 |
| Luna Experience (SC) | 7+3 | 6-10 fl. oz. | x | x | 12h | 34 fl. oz. |
| fluopyram + tebuconazol | | G-E | x | x | 1d | NA |
| Luna Privilege | 7 | 4-6.8 fl. oz. | x | x | 12h | 13.7 fl. oz. |
| fluopyram | | E | x | x | 0d | 6 |

(Continued)

Table 5-5. Plum Diseases - Petal Fall Through Shuck Split¹ (continued)

| Product And Formulation Active Ingredient | FRAC Code ² | Brown Rot | Bacterial Spot | Black Knot | REI ³ PHI ⁴ | Max Amt ⁵ Max App ⁶ |
|--|------------------------|-----------------|----------------|------------|--------------------------------------|--|
| Luna Sensation (SC) | 7+11 | 6-10 fl. oz. | x | x | 12h | 271 fl. oz. |
| fluopyram + trifloxystrobin | | E | x | x | 1d | 4 |
| Merivon XBF | 7+11 | 4-6.7 fl. oz. | x | x | 12h | 20.1 fl. oz. |
| fluxapyroxad + pyraclostrobin | | E | x | x | 0d | 4 |
| Microthiol Disperss | M | 10-20 lb. | x | x | 24h | NA |
| sulfur | | F | x | x | 0d | 3 |
| Miravis | 7 | 3.4-5.1 fl. oz. | x | x | 4 hr | 20.4 fl. oz. |
| pydiflumetofen | | E-G | x | x | 0d | 4 |
| Ph-D | 19 | 6.2 oz. | x | x | 0d | NA |
| polyoxin D | | G | x | x | NA | NA |
| Pristine | 7+11 | 10.5-14.5 oz. | x | x | 12h | 72.5 oz. |
| pyraclostrobin + boscalid | | G-E[r] | x | x | 0d | 9 |
| Quadris Top | 11+3 | 12-14 fl. oz. | x | x | 12h | 56 fl. oz. |
| azoxystrobin + difenoconazole | | E | x | x | 0d | 5 |
| Quash | 3 | 2.5-3.5 oz. | x | x | 12h | 12 oz. |
| metconazole | | E-G | x | x | 14d | 3 |
| Quilt Xcel | 11+3 | 14 fl. oz. | x | x | 12h | 70 fl. oz. |
| azoxystrobin + propiconazole | | E | x | x | 0d | 5 |
| Rally 40WSP | 3 | 2.5-6 oz. | x | x | 24h | 2.75 lb. |
| myclobutanil | | G | x | x | 1d | NA |
| Rovral 4F | 2 | 1-2 pt. | x | x | 24h | 4 pt. |
| iprodione | | E | x | x | N/A | NA |
| Scala (SC) | 9 | 9-18 fl. oz. | x | x | 12h | 54 fl. oz. |
| pyrimethanil | | G-E | x | x | 2d | 2 |
| Tilt (EC) | 3 | 4 fl. oz. | x | x | 12h | 20 fl. oz. |
| propiconazole | | E | x | x | 0d | 5 |
| Topguard EQ | 3+11 | 6-8 oz. | x | x | 12h | NA |
| flutriafol + azoxystrobin | | G | x | x | 7d | 4 |
| Topguard Specialty Crop | 3 | 13 fl. oz. | x | x | 12h | 56 fl. oz. |
| flutriafol | | E | x | x | 7d | 4 |
| Topsin M WSB | 1 | 1-1.5 lb. | x | 1-1.5 lb. | 48h | 4 lb. |
| thiophanate-methyl | | E[r] | x | E[r] | 1d | NA |
| Vanguard WG (75WG) | 9 | 5 oz. | x | x | 12h | 30 oz. |
| cyprodinil | | G-E | x | x | 2d | 4 |

Footnotes: See how to read the spray schedule tables section at the beginning of the chapter.

Plum Petal Fall To Second Cover - Insects

Notes on insect management

- **San Jose scale:** Insecticides are best applied when scale crawler nymphs are active.
 - Do not apply Movento until PF is complete.

- **Lesser peachtree borer:** Control of the first generation of lesser peachtree borer with trunk sprays is during the time of peak moth flight, generally from early May to early June, See Borers of Peach, Cherry, and Plum, page 152.

Table 5-6. Plum Insects - Petal Fall Through Second Cover¹

| Product And Formulation Active Ingredient | IRAC Code ² | European Red Mite | Plum Curculio | Japanese Beetle | San Jose Scale | Leaf-Roller | Oriental Fruit Moth | Peach Tree Borers | Periodical Cicada | REI ³ PHI ⁴ | Max Amt ⁵ Max App ⁶ |
|--|------------------------|-------------------|------------------|-----------------|-----------------|------------------|---------------------|-------------------|-------------------|-----------------------------------|---|
| Acramite 50WS | 20D | 1 lb. | x | x | x | x | x | x | x | 12h | 1 lb. |
| bifenazate | | G | x | x | x | x | x | x | x | 3d | 1 |
| Actara (25WDG) | 4A | x | 4.5-5.5 oz. | x | x | x | x | x | x | 12h | 11 oz. |
| thiamethoxam | | x | G | x | x | x | x | x | x | 14d | NA |
| Admire Pro (4.6F) | 4A | x | 2.8 fl. oz. | 1.4-2.8 fl. oz. | 1.4-2.8 fl. oz. | x | x | x | x | 12h | 10.5/14 fl. oz. |
| imidacloprid | | x | s | G | F | x | x | x | x | 0-21d | NA |
| Agri-Mek SC (0.7SC) (RUP) | 6 | 2.25-4.25 fl. oz. | x | x | x | x | x | x | x | 12h | 8.5 fl. oz. |
| abamectin | | G | x | x | x | x | x | x | x | 21d | 2 |
| Altacor (35WDG) | 28 | x | x | x | x | 3-4.5 oz. | 3-4.5 oz. | x | x | 4h | 9 oz. |
| chlorantraniliprole | | x | x | x | x | E | E | x | x | 10d | NA |
| Apta (1.34SC) | 21A | x | 21-27 fl. oz. | x | x | 21-27 fl. oz. | x | x | x | 12h | 53.5 fl. oz. |
| tolfenpyrad | | x | G | x | x | G | x | x | x | 14d | 2 |
| Asana XL (0.66EC) (RUP) | 3A | x | 4.8-14.5 fl. oz. | x | x | 4.8-14.5 fl. oz. | 4.8-14.5 fl. oz. | 4.8-14.5 fl. oz. | 4.8-14.5 fl. oz. | 12h | 72.5 fl. oz. |
| esfenvalerate | | x | G | x | x | E | E | G | E | 14d | NA |
| Assail 30SG | 4A | x | 5-8 oz. | 5.3-8 oz. | 5.3-8 oz. | x | 5.3-8 oz. | x | x | 12h | 32 oz. |
| acetamiprid | | x | E | F | F | x | E | x | x | 7d | 4 |
| Avaunt (30WDG) | 22 | x | 5-6 oz. | x | x | x | 6 oz. | x | x | 12h | 24 oz. |
| indoxacarb | | x | G | x | x | x | F | x | x | 14d | 4 |
| <i>Bacillus thuringiensis</i> (B.t.) (Dipel DF, etc) | 11A | x | x | x | x | 0.5-2 lb. | 0.5-2 lb. | x | x | 4h | NA |
| <i>Bacillus thuringiensis</i> | | x | x | x | x | u | u | x | x | 0d | NA |
| Baythroid XL (IEC) (RUP) | 3A | x | 2.4-2.8 fl. oz. | x | x | 2.4-2.8 fl. oz. | 2-2.4 fl. oz. | 1.4-2 fl. oz. | 2.4-2.8 fl. oz. | 12h | 5.6 fl. oz. |
| beta-cyfluthrin | | x | G | x | x | u | E | G | E | 7d | NA |
| BeetleGone! | 11A | x | x | 1-17.5 lb. | x | x | x | x | x | 4h | NA |
| <i>Bacillus thuringiensis</i> | | x | x | G | x | x | x | x | x | 0h | NA |
| Centaur WDG (70WDG) | 16 | x | x | x | 34.5 oz. | x | x | x | x | 12h | 69 oz. |
| buprofezin | | x | x | x | E | x | x | x | x | 14d | 2 |

(Continued)

Table 5-6. Plum Insects - Petal Fall Through Second Cover¹ (continued)

| Product And Formulation Active Ingredient | IRAC Code ² | European Red Mite | Plum Curculio | Japanese Beetle | San Jose Scale | Leaf-Roller | Oriental Fruit Moth | Peach Tree Borers | Periodical Cicada | REI ³ PHI ⁴ | Max Amt ⁵ Max App ⁶ |
|---|------------------------|----------------------------|-------------------|-------------------|----------------|-------------------|---------------------|-------------------|-------------------|-----------------------------------|---|
| Closer SC (2SC) | 4C | x | x | x | 5.7 fl. oz. | x | x | x | x | 12h | 17 fl. oz. |
| sulfoxaflor | | x | x | x | s | x | x | x | x | 7d | 4 |
| Danitol 2.4EC (RUP) | 3A | 10.3-21.3 fl. oz. | 10.3-21.3 fl. oz. | 10.3-21.3 fl. oz. | x | 10.3-21.3 fl. oz. | 10.3-21.3 fl. oz. | 10.7-21.3 fl. oz. | x | 24h | 42.7 fl. oz. |
| fenpropathrin | | F | G | E | x | u | E | G | x | 3d | NA |
| Delegate WG (25WG) | 5 | x | 6-7 oz. | x | x | 4.5-7 oz. | 6-7 oz. | x | x | 4h | 28 oz. |
| spinetoram | | x | s | x | x | E | E | x | x | 1d | 4 |
| Diazinon AG600 WBC (RUP) | 1B | 6.5-12.25 fl. oz./100 gal. | x | x | x | x | x | x | x | 4d | 102 fl. oz. |
| diazinon | | u | x | x | x | x | x | x | x | 21d | 2 |
| Dimilin 2L (RUP) | 15 | x | 8-16 fl. oz. | x | x | 8-16 fl. oz. | 8-16 fl. oz. | x | x | 12h | 32 fl. oz. |
| diflubenzuron | | x | u | x | x | u | u | x | x | 14d | 2 |
| Entrust SC (2SC) | 5 | x | x | x | x | 4-8 fl. oz. | 4-8 fl. oz. | x | x | 4h | 29 fl. oz. |
| spinosad | | x | x | x | x | u | F | x | x | 1d | 3 |
| Envidor 2SC | 23 | 16-18 fl. oz. | x | x | x | x | x | x | x | 12h | 18fl. oz. |
| spiroticlofen | | E | x | x | x | x | x | x | x | 7d | 1 |
| Esteem 35WP | 7C | x | x | x | 4-5 oz. | x | 4-5 oz. | x | x | 12h | 15 oz. |
| pyriproxyfen | | x | x | x | E | x | s | x | x | 14d | 3 |
| Exirel (0.83SE) | 28 | x | 13.5-20.5 fl. oz. | 13.5-20.5 fl. oz. | x | 10-20.5 fl. oz. | 10-20.5 fl. oz. | x | x | 12h | 61.5 fl. oz. |
| cyantraniliprole | | x | G | G | x | E | E | x | x | 3d | 3 |
| Imidan 70W | 1B | x | 2.13-4.25 lb. | 2.13-4.25 lb. | 2.13-4.25 lb. | 2.13-4.25 lb. | 2.13-4.25 lb. | x | x | 7d | 13 lb. |
| phosmet | | x | G | G | E | E | E | x | x | 7d | NA |
| Intrepid 2F | 18 | x | x | x | x | 8-16 fl. oz. | 10-16 fl. oz. | x | x | 4h | 64 fl. oz. |
| methoxyfenozide | | x | x | x | x | E | G | x | x | 7d | NA |
| Magister SC (1.7SC) | 21A | 32-36 fl. oz. | x | x | x | x | x | x | x | 12h | 36 fl. oz. |
| fenazaquin | | E | x | x | x | x | x | x | x | 3d | 1 |
| Movento (2SC) | 23 | 6-9 fl. oz. | x | x | 6-9 fl. oz. | x | x | x | x | 24h | 15.3 fl. oz. |
| spirotetramat | | s | x | x | G | x | x | x | x | 7d | NA |
| Mustang Maxx (0.83EC) (RUP) | 3A | x | 1.2-4 fl. oz. | x | x | 1.2-4 fl. oz. | 1.2-4 fl. oz. | 1.2-4 fl. oz. | x | 12h | 24 fl. oz. |
| zeta-cypermethrin | | x | E | x | x | E | E | G | x | 14d | NA |

(Continued)

Table 5-6. Plum Insects - Petal Fall Through Second Cover¹ (continued)

| Product And Formulation Active Ingredient | IRAC Code ² | European Red Mite | Plum Curculio | Japanese Beetle | San Jose Scale | Leaf-Roller | Oriental Fruit Moth | Peach Tree Borers | Periodical Cicada | REI ³ PHI ⁴ | Max Amt ⁵ Max App ⁶ |
|---|------------------------|-------------------|------------------|------------------|------------------|------------------|---------------------|-------------------|-------------------|-----------------------------------|---|
| Nexter (75WP) | 21 | 4.4-5.2 oz. | x | x | x | x | x | x | x | 12h | 21.3 oz. |
| pyridaben | | G | x | x | x | x | x | x | x | 7d | 2 |
| Oil (superior) | UN | 1.5-2% | x | x | 1.5-2% | x | x | x | x | 4h | NA |
| mineral oil | | E | x | x | G | x | x | x | x | 0d | NA |
| Onager (1EC) | 10A | 12-24 oz. | x | x | x | x | x | x | x | 12h | 24 fl. oz. |
| hexythiazox | | E | x | x | x | x | x | x | x | 7d | 1 |
| Portal XLO (0.4EC) | 21A | 2 pt. | x | x | x | x | x | x | x | 12h | 4 pt. |
| fenpyroximate | | E | x | x | x | x | x | x | x | 7d | 2 |
| Proaxis (0.5EC) (RUP) | 3A | x | 2.5-5.1 fl. oz. | 2.5-5.1 fl. oz. | x | 2.5-5.1 fl. oz. | 2.5-5.1 fl. oz. | 2.5-5.1 fl. oz. | 2.5-5.1 fl. oz. | 24h | 1.6 pt. |
| gamma-cyhalothrin | | x | G | E | x | E | G | G | E | 14d | NA |
| Pyganic 5EC | 3A | 4.5-15.6 fl. oz. | 4.5-15.6 fl. oz. | 4.5-15.6 fl. oz. | 4.5-15.6 fl. oz. | 4.5-15.6 fl. oz. | x | x | x | 12h | NA |
| pyrethrins | | u | i | F | u | u | x | x | x | 0d | 10 |
| Rimon 0.83EC | 15 | x | x | x | x | 20-50 fl. oz. | 20-40 fl. oz. | 20 fl. oz. | x | 12h | 150 fl. oz. |
| novaluron | | x | x | x | x | E | E | u | x | 8d | NA |
| Savey 50DF | 10A | 3-6 oz. | x | x | x | x | x | x | x | 12h | 6 oz. |
| hexythiazox | | E | x | x | x | x | x | x | x | 28d | 1 |
| Sevin XLR Plus (4F) | 1A | x | 2-3 qt. | 2-3 qt. | 4-5 qt. | 2-3 qt. | 2-3 qt. | 2-3 qt. | 2-3 qt. | 12h | 14 qt. |
| carbaryl | | x | F | E | u | F | F | i | G | 3d | 3 |
| Sivanto Prime (1.67SC) | 4D | x | x | x | 10.5-14 fl. oz. | x | x | x | x | 4h | 28 fl. oz. |
| flupyradifurone | | x | x | x | G | x | x | x | x | 14d | NA |
| Surround WP (95WP) | UN | x | 25-50 lb. | 25-50 lb. | x | 25-50 lb. | 25-50 lb. | 25-50 lb. | x | 4h | NA |
| kaolin | | x | F | F | x | u | u | s | x | 0d | NA |
| Vendex 50WP (RUP) | 12B | 1-2 lb. | x | x | x | x | x | x | x | 2d | 3 lb. |
| fenbutatin-oxide | | G | x | x | x | x | x | x | x | 14d | 2 |
| Verdepryn 100SL (0.83SL) | 28 | x | 5.5-11 fl. oz. | 5.5-11 fl. oz. | x | 5.5-11 fl. oz. | 5.5-11 fl. oz. | x | x | 4h | 33 fl. oz. |
| cyclaniliprole | | x | G | u | x | E | E | x | x | 7d | 3 |
| Warrior II (2.08CS) (RUP) | 3A | x | 1.2-2.5 fl. oz. | 1.2-2.5 fl. oz. | x | 1.2-2.5 fl. oz. | 1.2-2.5 fl. oz. | 1.2-2.5 fl. oz. | 1.2-2.5 fl. oz. | 24h | 12.8 fl. oz. |
| lambda-cyhalothrin | | x | G | E | x | u | G | G | E | 14d | NA |
| Zeal (72WP) | 10B | 2-3 oz. | x | x | x | x | x | x | x | 12h | 3 oz. |
| etoxazole | | E | x | x | x | x | x | x | x | 7d | 1 |

Footnotes: See how to read the spray schedule tables section at the beginning of the chapter.

Plum Summer Cover Sprays - Diseases

7-10 days after shuck split

Notes on disease management

Bacterial spot

- Copper pesticides are not recommended after first cover, to reduce the risk of phytotoxicity.

Brown rot

- Although captan is still labeled for use for brown rot control, it is phytotoxic to some varieties of European, hybrid and Japanese plums. Symptoms of phytotoxicity include shot-holes of leaves and

spotting of fruit, and may be mistaken for bacterial leaf spot.

- To prevent brown rot at harvest, continue sprays 21 days before harvest. Depending on rainfall, repeat applications every 5-10 days, if needed. Merivon, Luna Sensation and Pristine all have PHIs of 1 day or less.

Powdery mildew

- Stanley plums: **Do not apply Quash.** Do not apply Quilt Xcel to Stanley plums earlier than 21 days prior to harvest, which has been implicated in reduced size and shape issues.

Table 5-7. Plum Diseases - Summer Cover¹

| Product And Formulation | Active Ingredient | FRAC Code ² | Brown Rot | Powdery Mildew | REI ³ PHI ⁴ | Max Amt ⁵ Max App ⁶ |
|-------------------------|-------------------------------|------------------------|-----------------|-----------------|--------------------------------------|--|
| Abound (SC) | | 11 | 12-15.5 fl. oz. | 12-15.5 fl. oz. | 4h | 92.3 fl. oz. |
| | azoxystrobin | | F-E[r] | F | 0d | 15 |
| Captan 80WDG | | M | 3.7 lb. | x | 24h | 40 lb. |
| | captan | | G | x | 0d | NA |
| Cevya | | 3 | 5 oz. | 3-5 fl. oz. | 12h | 15 fl. oz. |
| | mefentrifluconazole | | E | E | 0d | 5 |
| Elevate 50 WDG | | 17 | 1-1.5 lb. | x | 12h | 6 lb. |
| | fenhexamid | | G-E | x | 0d | NA |
| Elite 45DF | | 3 | 2 oz. | x | 12h | 3 lb. |
| | tebuconazole | | E | x | 0d | NA |
| Flint Extra | | 11 | 2.5-3.8 fl. oz. | 2.5-3.8 fl. oz. | 12h | 15.2 fl. oz. |
| | trifloxystrobin | | G-E | E | 1d | 3 |
| Fontelis (SC) | | 7 | 14-20 fl. oz. | 14-20 fl. oz. | 12h | 61 fl. oz. |
| | penthiopyrad | | G-E | F-G | 0d | 4 |
| Indar 2F | | 3 | 6 fl. oz. | x | 12h | 48 fl. oz. |
| | fenbuconazole | | E[r] | G-E | 0d | NA |
| Inspire Super (EW) | | 3+9 | 16-20 fl. oz. | 16-20 fl. oz. | 12h | 80 fl. oz. |
| | difenoconazole + cyprodinil | | E | G-F | 2d | 8 |
| Kenja 400 SC | | 7 | 12.5 fl. oz. | x | 12h | 37.5 fl. oz. |
| | isofetamid | | E | S | 1 day | 3 |
| Luna Experience (SC) | | 7+3 | 6-10 fl. oz. | 6-10 fl. oz. | 12h | 34 fl. oz. |
| | fluopyram + tebuconazol | | G-E | E | 1d | NA |
| Luna Privilege | | 7 | 4-6.8 fl. oz. | 4-6.8 fl. oz. | 12h | 13.7 fl. oz. |
| | fluopyram | | E | G | 0d | 6 |
| Luna Sensation (SC) | | 7+11 | 6-10 fl. oz. | 6-10 fl. oz. | 12h | 27.1 fl. oz. |
| | fluopyram + trifloxystrobin | | E | G-E | 1d | 4 |
| Merivon XBF | | 7+11 | 4-6.7 fl. oz. | 4-6.7 fl. oz. | 12h | 20.1 fl. oz. |
| | fluxapyroxad + pyraclostrobin | | E | E-G | 0d | 4 |

(Continued)

Table 5-7. Plum Diseases - Summer Cover¹ (continued)

| Product And Formulation Active Ingredient | FRAC Code ² | Brown Rot | Powdery Mildew | REI ³ PHI ⁴ | Max Amt ⁵ Max App ⁶ |
|--|------------------------|-----------------|-------------------|--------------------------------------|--|
| Microthiol Disperss | M | 10-20 lb. | 10-20 lb. | 24h | NA |
| sulfur | | F | G | 0d | 3 |
| Miravis | 7 | 3.4-5.1 fl. oz. | 3.4-5.1 fl. oz. | 4 hr | 20.4 fl. oz. |
| pydiflumetofen | | G-E | G-E | 0d | 4 |
| Ph-D | 19 | 6.2 oz. | 6.2 oz. | 0d | NA |
| polyoxin D | | G | G | NA | NA |
| Pristine | 7+11 | 10.5-14.5 oz. | 10.5-14.5 oz. | 12h | 72.5 oz. |
| pyaclostrobin + boscalid | | E-G[r] | E[r] | 0d | 9 |
| Quadris Top | 11+3 | 12-14 fl. oz. | 12-14 fl. oz. | 12h | 56 fl. oz. |
| azoxystrobin + difenoconazole | | E | G | 0d | 5 |
| Quash | 3 | 2.5-3.5 oz. | 3.5-4 oz. | 12h | 12 oz. |
| metconazole | | E-G | E | 14d | 3 |
| Quilt Xcel | 11+3 | 14 fl. oz. | 14 fl. oz. | 12h | 70 fl. oz. |
| azoxystrobin + propiconazole | | E | G | 0d | 5 |
| Quintec | 13 | x | 7 fl. oz. | 12h | 28 fl. oz. |
| quinoxifen | | x | E | 7d | 4 |
| Rally 40WSP | 3 | 2.5-6 oz. | 2.5-6 oz. | 24h | 2.75 lb. |
| myclobutanil | | G | G-E | 1d | NA |
| Rovral 4F | 2 | 1-2 pt. | x | 24h | 4 pt. |
| iprodione | | E | x | N/A | NA |
| Scala (SC) | 9 | 9-18 fl. oz. | x | 12h | 54 fl. oz. |
| pyrimethanil | | G-E | x | 2d | 2 |
| Tilt (EC) | 3 | 4 fl. oz. | 4 fl. oz. | 12h | 20 fl. oz. |
| propiconazole | | E | G | 0d | 5 |
| Topguard EQ | 3+11 | 6-8 oz. | 6-8 oz. | 12h | NA |
| flutriafol + azoxystrobin | | G | E | 7d | 4 |
| Topguard Specialty Crop | 3 | 14 oz. | 14 oz. | 12h | 56 fl. oz. |
| flutriafol | | G | E | 7d | 4 |

Footnotes: See how to read the spray schedule tables section at the beginning of the chapter.

Plum Second To Summer Cover Sprays - Insects

Two weeks after first cover spray and 10- to 14-day intervals thereafter

Notes on insect management

- **Plum curculio** can stay active thorough second cover. Failure to control plum curculio may result in an increase in brown rot.
- **Peachtree borer:** Best controlled by a trunk drench at the time of peak moth flight in, usually

in early August. See Borers of Peach, Cherry and Plum, page 152.

- **Japanese beetle:** Begin applications for Japanese beetle as soon as observed. Multiple applications may be needed.
- **Cherry fruit flies:** Adults emerge from late May to early July and lay their eggs in the fruits. Sprays need to target the adults before egg laying begins, generally 5 to 6 days after adults emerge.

Table 5-8. Plum Insects - Second Through Summer Cover¹

| Product And Formulation Active Ingredient | IRAC Code ² | European Red Mite | Cherry Fruit Fly | Japanese Beetle | Leaf- roller | Oriental Fruit Moth | Spotted Wing Drosophila | REI ³ PHI ⁴ | Max Amt ⁵ Max App ⁶ |
|---|---------------------------|----------------------------------|---------------------|----------------------|----------------------|---------------------------|-------------------------------|--------------------------------------|--|
| Acramite 50WS | 20D | 1 lb. | x | x | x | x | x | 12h | 1 lb. |
| bifenazate | | G | x | x | x | x | x | 3d | 1 |
| Actara (25WDG) | 4A | x | 4.5-5.5 oz. | x | x | x | x | 12h | 11 oz. |
| thiamethoxam | | x | G | x | x | x | x | 14d | NA |
| Admire Pro (4.6F) | 4A | x | 2-2.8 fl. oz. | 1.4-2.8 fl. oz. | x | x | x | 12h | 10.5/14 fl. oz. |
| imidacloprid | | x | G | G | x | x | x | 0-21d | NA |
| Agri-Mek SC (0.7SC) (RUP) | 6 | 2.25-4.25 fl. oz. | x | x | x | x | x | 12h | 8.5 fl. oz. |
| abamectin | | G | x | x | x | x | x | 21d | 2 |
| Altacor (35WDG) | 28 | x | 3-4.5 oz. | x | 3-4.5 oz. | 3-4.5 oz. | x | 4h | 9 oz. |
| chlorantraniliprole | | x | s | x | E | E | x | 10d | NA |
| Apta (1.34SC) | 21A | x | 14-27 fl. oz. | x | 21-27 fl. oz. | x | 21-27 fl. oz. | 12h | 53.5 fl. oz. |
| tolfenpyrad | | x | F | x | G | x | s | 14d | 2 |
| Asana XL (0.66EC) (RUP) | 3A | x | 4.8-14.5 fl. oz. | x | 4.8-14.5 fl. oz. | 4.8-14.5 fl. oz. | x | 12h | 72.5 fl. oz. |
| esfenvalerate | | x | G | x | E | E | x | 14d | NA |
| Assail 30SG | 4A | x | 5.3-8 oz. | 5.3-8 oz. | x | 5.3-8 oz. | x | 12h | 32 oz. |
| acetamiprid | | x | G | F | x | E | x | 7d | 4 |
| Avaunt (30WDG) | 22 | x | x | x | x | 6 oz. | x | 12h | 24 oz. |
| indoxacarb | | x | x | x | x | F | x | 14d | 4 |
| <i>Bacillus thuringiensis</i> (B.t.) (Dipel DF, etc) | 11A | x | x | x | 0.5-2 lb. | 0.5-2 lb. | x | 4h | NA |
| <i>Bacillus thuringiensis</i> | | x | x | x | u | u | x | 7d | NA |
| Baythroid XL (1EC) (RUP) | 3A | x | 2.4-2.8 fl. oz. | x | 2.4-2.8 fl. oz. | 2-2.4 fl. oz. | x | 12h | 5.6 fl. oz. |
| beta-cyfluthrin | | x | G | x | u | E | x | 7d | NA |
| BeetleGone! | 11A | x | x | 1-17.5 lb. | x | x | x | 4h | NA |
| <i>Bacillus thuringiensis</i> | | x | x | G | x | x | x | 0d | NA |
| Danitol 2.4EC (RUP) | 3A | 10.3-21.3 fl. oz. | 16-21.3 fl. oz. | 10.3-21.3 fl. oz. | 10.3-21.3 fl. oz. | 10.3-21.3 fl. oz. | 10.7-21.3 fl. oz. | 24h | 42.7 fl. oz. |
| fenpropathrin | | F | G | E | u | E | E | 3d | NA |
| Delegate WG (25WG) | 5 | x | x | x | 4.5-7 oz. | 6-7 oz. | 4.5-7 oz. | 4h | 28 oz. |
| spinetoram | | x | x | x | E | E | E | 1d | 4 |
| Diazinon AG600 WBC (RUP) | 1B | 6.5-12.25 fl. oz./100 gal. | x | x | x | x | x | 4d | 102 fl. oz. |
| diazinon | | u | x | x | x | x | x | 21d | 2 |
| Dimilin 2L (RUP) | 15 | x | x | x | 8-16 fl. oz. | 8-16 fl. oz. | x | 12h | 32 fl. oz. |
| diflubenzuron | | x | x | x | u | u | x | 14d | 2 |

(Continued)

Table 5-8. Plum Insects - Second Through Summer Cover¹ (continued)

| Product And Formulation Active Ingredient | IRAC Code ² | European Red Mite | Cherry Fruit Fly | Japanese Beetle | Leaf- roller | Oriental Fruit Moth | Spotted Wing Drosophila | REI ³ PHI ⁴ | Max Amt ⁵ Max App ⁶ |
|--|---------------------------|----------------------|---------------------|----------------------|---------------------|---------------------------|-------------------------------|--------------------------------------|--|
| Entrust SC (2SC) | 5 | x | 4-8 fl. oz. | x | 4-8 fl. oz. | 4-8 fl. oz. | x | 4h | 29 fl. oz. |
| spinosad | | x | G | x | u | F | x | 1d | 3 |
| Envidor 2SC | 23 | 16-18 fl. oz. | x | x | x | x | x | 12h | 18 fl. oz. |
| spirodiclofen | | E | x | x | x | x | x | 7d | 1 |
| Esteem 35WP | 7C | x | x | x | x | 4-5 oz. | x | 12h | 15 oz. |
| pyriproxyfen | | x | x | x | x | s | x | 14d | 3 |
| Exirel (0.83SE) | 28 | x | 10-17 fl. oz. | 13.5-20.5 fl. oz. | 10-20.5 fl. oz. | 10-20.5 fl. oz. | 13.5-20.5 fl. oz. | 12h | 61.5 fl. oz. |
| cyantraniliprole | | x | E | G | E | E | G | 3d | 3 |
| Imidan 70W | 1B | x | x | 2.1-4.25 lb. | 2.1-4.25 lb. | 2.1-4.25 lb. | x | 7d | 13 lb. |
| phosmet | | x | x | G | E | E | x | 7d | NA |
| Intrepid 2F | 18 | x | x | x | 8-16 fl. oz. | 10-16 fl. oz. | x | 4h | 64 fl. oz. |
| methoxyfenozide | | x | x | x | E | G | x | 7d | NA |
| Magister SC (1.7SC) | 21A | 32-36 fl. oz. | x | x | x | x | x | 12h | 36 fl. oz. |
| fenazaquin | | E | x | x | x | x | x | 3d | 1 |
| Movento (2SC) | 23 | 6-9 fl. oz. | 6-9 fl. oz. | x | x | x | 6-9 fl. oz. | 24h | 15.3 fl. oz. |
| spirotetramat | | s | s | x | x | x | s | 7d | NA |
| Mustang Maxx (0.83EC) (RUP) | 3A | x | 1.28-4 fl. oz. | x | 1.28-4 fl. oz. | 1.28-4 fl. oz. | 4 fl. oz. | 12h | 24 fl. oz. |
| zeta-cypermethrin | | x | F | x | E | E | E | 14d | NA |
| Nexter (75WP) | 21 | 4.4-5.2 oz. | x | x | x | x | x | 12h | 21.3 oz. |
| pyridaben | | G | x | x | x | x | x | 7d | 2 |
| Oil (superior) | UN | see label | x | x | x | x | x | 4h | NA |
| mineral oil | | E | x | x | x | x | x | 0d | NA |
| Onager (1EC) | 10A | 12-24 oz. | x | x | x | x | x | 12h | 24 fl. oz. |
| hexythiazox | | E | x | x | x | x | x | 7d | 1 |
| Portal XLO (0.4EC) | 21A | 2 pt. | x | x | x | x | x | 12h | 4 pt. |
| fenpyroximate | | E | x | x | x | x | x | 7d | 2 |
| Proaxis (0.5EC) (RUP) | 3A | x | 2.5-5.1 fl. oz. | 2.5-5.1 fl. oz. | 2.5-5.1 fl. oz. | 2.5-5.1 fl. oz. | x | 24h | 1.6 pt. |
| gamma-cyhalothrin | | x | G | E | E | G | x | 14d | NA |
| Pyganic 5EC | 3A | 4.5-15.6 fl. oz. | 4.5-15.6 fl. oz. | 4.5-15.6 fl. oz. | 4.5-15.6 fl. oz. | x | 4.5-15.6 fl. oz. | 12h | NA |
| pyrethrins | | u | u | F | u | x | i | 0d | 10 |
| Rimon 0.83EC | 15 | x | 20-40 fl. oz. | x | 20-50 fl. oz. | 20-40 fl. oz. | x | 12h | 150 fl. oz. |
| novaluron | | x | G | x | E | E | x | 8d | NA |

(Continued)

Table 5-8. Plum Insects - Second Through Summer Cover¹ (continued)

| Product And Formulation Active Ingredient | IRAC Code ² | European Red Mite | Cherry Fruit Fly | Japanese Beetle | Leaf- roller | Oriental Fruit Moth | Spotted Wing Drosophila | REI ³ PHI ⁴ | Max Amt ⁵ Max App ⁶ |
|--|---------------------------|----------------------|---------------------|--------------------|--------------------|---------------------------|-------------------------------|--------------------------------------|--|
| Savey 50DF | 10A | 3-6 oz. | x | x | x | x | x | 12h | 6 oz. |
| hexythiazox | | E | x | x | x | x | x | 28d | 1 |
| Sevin XLR Plus (4F) | 1A | x | 2-3 qt. | 2-3 qt. | 2-3 qt. | 2-3 qt. | x | 12h | 14 qt. |
| carbaryl | | x | G | E | F | F | x | 3d | 3 |
| Surround WP (95WP) | UN | x | 25-50 lb. | 25-50 lb. | 25-50 lb. | 25-50 lb. | x | 4h | NA |
| kaolin | | x | G | F | u | u | x | 0d | NA |
| Vendex 50WP (RUP) | 12B | 1-2 lb | x | x | x | x | x | 2d | 3 lb. |
| fenbutatin-oxide | | G | x | x | x | x | x | 14d | 2 |
| Verdepryn 100SL (0.83SL) | 28 | x | 5.5-11 fl. oz. | 5.5-11 fl. oz. | 5.5-11 fl. oz. | 5.5-11 fl. oz. | 5.5-11 fl. oz. | 4h | 33 fl. oz. |
| cyclaniliprole | | x | G | u | E | E | u | 7d | 3 |
| Warrior II (2.08CS) (RUP) | 3A | x | 1.2-2.5 fl. oz. | 1.2-2.5 fl. oz. | 1.2-2.5 fl. oz. | 1.2-2.5 fl. oz. | x | 24h | 12.8 fl. oz. |
| lambda-cyhalothrin | | x | G | E | u | G | x | 14d | NA |
| Zeal (72WP) | 10B | 2-3 oz. | x | x | x | x | x | 12h | 3 oz. |
| etoxazole | | E | x | x | x | x | x | 7d | 1 |

Footnotes: See how to read the spray schedule tables section at the beginning of the chapter.

Special Problems of Stone Fruit

Borers of peach, cherry, and plum trees

The peachtree borer, lesser peachtree borer and shot-hole borer often infest peach, apricot, cherry and plum trees. Peachtree borers infest the trunk at the soil line, while lesser peachtree borers infest scaffold limbs and the upper trunk.

The peachtree borer is primarily a pest of young trees, whereas the lesser peachtree borer is a pest of older trees. The shothole borer is often found in trees of low vigor with dead and/or diseased limbs. Moths of the two peachtree borers lay their eggs on the surface of bark; shothole beetles lay their eggs in the inner bark.

Some of the regularly applied cover sprays help control borers; however, specific trunk and scaffold branch spray are often required. Pheromone traps are available to monitor emergence of the adult stage (moth) of lesser peachtree borer and peachtree borer. Knowing the time of emergence can help you properly time insecticide applications, because insecticides target the hatching eggs laid by the newly emerged moths.

Bacterial canker of sweet cherry

Bacterial canker is a sporadic but serious problem on sweet cherries. It is generally less severe on tart cherries, plums and prunes.

The disease is favored by cold, wet conditions during and shortly after bloom. Copper compounds are moderately effective in reducing pathogen populations and controlling the disease. Apply copper compounds according to the product label in the spring while trees are dormant.

If favorable conditions for the disease persist, apply reduced-rate applications (25-35 percent of dormant rate) after bud break, but before bloom. You can add hydrated lime (6-9 lb./acre) to reduce the phytotoxicity that can occur when copper compounds are applied in cool, wet conditions.

Bacterial spot of peach, nectarine and plum

Bacterial spot of stone fruit can be a serious problem in certain varieties, areas and years. The disease is favored by stormy, rainy weather from May through August. It causes the most damage in areas where the soil is sandy and where strong winds blow the sand.

Planting cultivars resistant to bacterial spot provides the best control. In the past, various control programs have been tried using foliar sprays of zinc sulfate plus lime, or fall applications of copper with or without lime. None of these programs offered reliable control and, in some cases, caused foliar and twig damage.

For peaches, the antibiotic oxytetracycline (My-coshield or FireLine) provides good control when properly applied. For best results, use oxytetracycline at 12 oz. per 100 gal. of dilute spray. Use dilute or 2x; higher concentrates are not effective and may be phytotoxic. These products are no longer labeled for plums.

Spraying the entire tree once per week is essential. If you spray only one side of the tree (alternate row middle), make certain to spray the other side of the tree within 3-4 days. Begin sprays at shuck split and continue at 7-day intervals until 21 days before harvest. Copper sprays, applied for peach leaf curl at leaf drop, also may help control bacterial spot. The rate of copper decreases over the growing season. On peaches, copper can cause injury to leaves and appears as reddish spots and shot-holes with some very mild defoliation when using an effective rate of copper. Because of risk for foliar injury, most copper materials are not registered for use post-bloom or in more than two applications post-bloom, and then only at low rates (always read and follow label instructions).

For more information, see Learning from Peach Bacterial Spot Epidemics: Potential Strategies for Reducing Fruit Losses (David Ritchie, North Carolina State University), <https://plantpathology.ces.ncsu.edu/wp-content/uploads/2013/06/Learning-from-Peach-Bacterial-Spot-Epidemics.pdf?fwd=no>

Phytophthora root, crown and collar rots

Peach rootstocks are highly susceptible to *Phytophthora* root, crown and collar rots. The main defense against these diseases is providing good soil drainage through proper site selection and tiling.

However, Ridomil Gold SL provides additional protection in wet years, on marginal sites, or in wetter sections of the orchard. Make applications just before growth starts in the spring and at two- to three-month intervals thereafter if soil is very wet. Apply to the soil beneath the tree canopy in sufficient water to ensure good coverage. (Subsequent rain or irrigation moves material into the soil.)

Ridomil Gold SL is also registered for use on cherries (sweet and tart), nectarines, plums and prunes. See label for further information and use rates. See page 133 for additional information on phosphorous acid, phosphonates and phosphites.

Periodical cicadas

See Periodical Cicadas in the Insect Management Notes section for apples, page 52.

6. GRAPE

Grape Insect Pests

The shaded/colored boxes represent the crop stages where common pests in the Midwest are active. Scouting and/or preventative sprays may be necessary or recommended.

| Grape Growth Stage | | | | | | | | |
|-----------------------------------|------------------|----------------------|-----------------------------|-------|-----------------------|---------------------|---------------------------------|-------------|
| Delayed Dormant Through Bud Swell | Bud Break | 4- To 10-inch Shoots | Pre-bloom Through Bloom | Bloom | Shatter | Shatter To Veraison | Veraison To Harvest | Postharvest |
| Grape Flea Beetle | | | | | | | | |
| | Grape Phylloxera | | | | | | | |
| | | Rose Chafer | | | Rose Chafer | | | |
| | | | | | Grape Berry Moth | | | |
| | | | | | Japanese Beetle | | | |
| | | | | | | | Spotted-Wing Drosophila | |
| | | | | | | | Multi-Colored Asian Lady Beetle | |
| | | | | | | | Green June Beetle | |
| | | | | | | | Grape Root Borer | |
| Climbing Cutworm | | | | | | | | |
| Spider Mites | | Spider Mites | | | | | | |
| Grape Scale | | | Grape Scale | | | | | |
| Grape Mealy Bug | | | | | Grape Mealybug | | | |
| | | Redbanded Leafroller | | | Red-Banded Leafroller | | | |
| | | | Eight-Spot- ted Forester | | | | | |
| | | | Grape Cane Girdler | | | | | |
| | | | Grape Cane Gallmaker | | | | | |
| | | | | | Grape Rootworm | | | |
| | | | | | | | Stink Bug | |
| | | | | | Spotted Lanternfly | | | |

Grape Spray Schedule

How to read the spray schedule tables

Every grape growth stage has important notes on disease or insect management. In some cases, the

reader will be directed to the special problems section at the end of the chapter. Please make sure to read thoroughly and contact your local Extension Specialist with questions.

Key to tables

E = excellent control

G = good control

F = fair control

[r] = Fungicide/Insecticide resistance possible

s = suppression only

i = ineffective

u = unknown efficacy

x = pest not on the label

¹ Efficacy data in this publication are based on trials conducted across various regions and does not necessarily reflect local efficacy differences or changes over time. Growers should contact their Extension Specialist for the most recent or for state-specific information. The information in this publication is only a guide; the authors and their institutions assume no liability for practices implemented based on this information. Always read and follow pesticide labels. The label is the law. Product registration may vary by state.

² F/I-RAC code represents the mode of action of the fungicide/insecticide.

³ PHI refers to the pre-harvest interval, which is the number of days before harvest that the product may not be applied.

⁴ All fungicides/insecticides have a Restricted-Entry Interval (REI). The restricted-entry interval is the time immediately after a pesticide application when entry into the treated area is limited. Check labels for REI as restrictions may prohibit the use of certain pesticides during harvest.

⁵ Max amt refers to the product's maximum amount/acre/year. Applicators must abide by both maximum amount of product per season AND maximum number of applications.

⁶ Max app refers to the product's maximum number of applications per year. Applicators must abide by both maximum amount of product per season AND maximum number of applications.

⁷ For treated wine grapes, the REI is 5 days when conducting cane tying, turning, or girdling. The REI is 12 hours for all other activities in wine grapes.

⁸ For table grapes, the REI for cane tying, turning, or girdling is 7 days.

RUP refers to restricted use pesticide.

Notes on disease control recommendations

The following information is intended to provide general guidelines for use in developing a fungicide spray program for grapes in the Midwest. This spray schedule presents various fungicide options that growers

can consider. The major grape diseases that generally require at least some fungicide application for control on an annual basis include black rot, powdery mildew, downy mildew, and Phomopsis blight. Several recommendations in this guide include tank mixes of different fungicides that are intended to provide a program to control all these diseases simultaneously.

Phototoxicity Alerts

It is extremely important to read fungicide labels carefully as many fungicides can cause crop injury when used alone or combined with insecticides or horticultural oils.

- All fungicides with difenoconazole labeled for grapes have the following precaution: "On *V. labrusca*, *V. labrusca* hybrids, and other non-vinifera hybrids where sensitivity is not known, the use of Inspire Super, Quadris Top, or Revus Top by itself or in tank mixes with materials that may increase uptake (adjuvants, foliar fertilizers) may result in leaf burning or other phytotoxic effects."
- Concord and other American-type grapes: Crop injury may occur if Flint Extra, Inspire Super, Flint Extra, Intuity, Luna Experience, Merivon, Pristine, Quadris Top, REGEV, or Revus Top are used on Concord grapes.
- Pristine should not be applied to Concord or Noiret grapes.
- Merivon should not be applied to Concord, Noiret, or NY73.0136.17
- Do not apply sulfur or captan at the same time as an oil or within two weeks of an oil application.
- Do not tank mix captan with the insecticide Sevin XLR.

Grape Delayed Dormant To Bud Swell - Diseases

Apply just as buds are beginning to swell but before they show green.

Notes on disease management

- **Anthracnose, Black rot, Powdery mildew and Phomopsis:** A delayed dormant application of lime-sulfur or Sulforix (calcium polysulfide) is recommended to reduce overwintering inoculum of the fungi that cause these four diseases.

Table 6-1. Grape Diseases - Delayed Dormant Through Bud Swell¹

| Product And Formulation Active Ingredient | FRAC Code ² | Anthraco | Phomopsis | Powdery Mildew | REI ³ PHI ⁴ | Max Amt ⁵ Max App ⁶ |
|--|---------------------------|----------|------------|----------------|--------------------------------------|--|
| Sulfurix | M | x | See Label | See Label | 48h | 16 gal. |
| calcium polysulfide | | x | E | s(E) | NA | 8 |
| Lime Sulfur | M | x | 15-20 gal. | 4-10 gal. | 48 h | 46 gal. |
| calcium polysulfide | | x | E | s | 0 day | 16 |

Footnotes: See How to read the spray schedule tables section at the beginning of the chapter.

**Grape Delayed Dormant To Bud Swell -
Insects**

Apply just as buds are beginning to swell but before they show green.

Notes on insect pest management

- **Flea beetle (adults) and climbing cutworms:**
Scout at least weekly as bud swell occurs.

Table 6-2. Grape Insects - Delayed Dormant Through Bud Swell¹

| Product And Formulation Active Ingredient | IRAC Code ² | Climbing Cutworm | Grape Flea Beetle | Mites: European Red, Spider Mite | Scale | REI ³ PHI ⁴ | Max Amt ⁵ Max App ⁶ |
|--|---------------------------|---------------------|----------------------|--|------------------------|--------------------------------------|--|
| Admire Pro (4.6F) | 4A | x | x | x | 7-14 fl. oz. (soil) | 12h | 2.8/14 fl. oz. |
| imidacloprid | | x | x | x | G | 0/30d | NA |
| Agri-Mek SC (0.7SC) (RUP) | 6 | x | x | 1.75-3.5 fl. oz. | x | 12h/4d | 7 fl. oz. |
| abamectin | | x | x | G | x | 28d | 2 |
| Altacor eVo (35WDG) | 28 | 3-4.5 oz. | x | x | x | 4h | 4.6 oz. |
| chlorantraniliprole | | G | x | x | x | 1d | 3 |
| Apollo SC (1SC) | 10A | x | x | 4-8 oz. | x | 12h/14d | NA |
| clofentezine | | x | x | E | x | 21d | 1 |
| Azera 0.21EC | 3A | 16-56 fl. oz. | 16-56 fl. oz. | 16-56 fl. oz. | 16-56 fl. oz. | 12h | NA |
| azadirachtin + pyrethrins | | u | u | u | u | 0d | 10 |
| Baythroid XL (1EC) (RUP) | 3A | 2.4-3.2 fl. oz. | 2.4-3.2 fl. oz. | x | x | 12h | 12.8 fl. oz. |
| beta-cyfluthrin | | E | G | x | x | 3d | NA |
| Brigade 2EC (RUP) | 3A | 3.2-6.4 fl. oz. | x | 6.4 fl. oz. | x | 12h | 7 fl. oz. |
| bifenthrin | | G | x | u | x | 30d | NA |
| Brigade WSB (10WP) (RUP) | 3A | 8-16 oz. | x | 16 oz. | x | 12h | 16 oz. |
| bifenthrin | | G | x | u | x | 30d | NA |
| B.t. (Agree, Dipel, etc.) | 11A | x | x | x | x | 4h | NA |
| <i>Bacillus thuringiensis</i> | | F | x | x | x | 0d | NA |
| Danitol 2.4EC (RUP) | 3A | 10.6-21.3 fl. oz. | 5.3-10.6 fl. oz. | 10.6-21.3 fl. oz. | x | 24h | 42.6 fl. oz. |
| fenpropathrin | | G | E | G | x | 21d | NA |
| Delegate WG (25WG) | 5 | 3-5 oz. | x | x | x | 4h | 19.5 oz. |
| spinetoram | | G | x | x | x | 3d | 5 |
| Entrust SC (2SC) | 5 | 4-8 fl. oz. | x | x | x | 4h | 23 fl. oz. |
| spinosad | | G | x | x | x | 3d | 5 |

(Continued)

Table 6-2. Grape Insects - Delayed Dormant Through Bud Swell¹ (continued)

| Product And Formulation Active Ingredient | IRAC Code ² | Climbing Cutworm | Grape Flea Beetle | Mites: European Red, Spider Mite | Scale | REI ³ PHI ⁴ | Max Amt ⁵ Max App ⁶ |
|--|---------------------------|---------------------|----------------------|--|------------------|--------------------------------------|--|
| Envidor 2SC | 23 | x | x | 16-34 fl. oz. | x | 12h | 34 fl. oz. |
| spirodiclofen | | x | x | E | x | 14d | 1 |
| Imidan 70W | 1B | x | 1.3-2.1 lb. | x | x | 14d | 6.5 lb. |
| phosmet | | x | F | x | x | 7/14d | NA |
| Kanemite 15SC | 20B | x | x | 21-31 fl. oz. | x | 12h | 62 fl. oz. |
| acequinocyl | | x | x | F | x | 7d | 2 |
| Magister SC (1.7SC) | 21A | x | x | 32-36 fl. oz. | x | 12h | 36 fl. oz. |
| fenazaquin | | x | x | u | x | 7d | 1 |
| Malathion 5EC | 1B | x | x | 3 pt. | 3 pt. | 72h | NA |
| malathion | | x | x | u | u | 3d | 2 |
| Movento (2SC) | 23 | x | x | 6-8 fl. oz. | 6-8 fl. oz. | 24h | 12.5 fl. oz. |
| spirotetramat | | x | x | s | s | 7d | NA |
| Mustang Maxx (0.83EC) (RUP) | 3A | 2-4 fl. oz. | x | x | x | 12h | 24 fl. oz. |
| zeta-cypermethrin | | E | x | x | x | 1d | NA |
| Nealta (1.67SC) | 25 | x | x | 13.7 fl. oz. | x | 12h | 28 fl. oz. |
| cyflumetofen | | x | x | G | x | 14d | 2 |
| Nexter SC (3.75SC) | 21A | x | x | 7.5-17 fl. oz. | x | 12h | NA |
| pyridaben | | x | x | G | x | 7d | 2 |
| Onager (1EC) | 10A | x | x | 12-24 oz. | x | 12h | 24 oz. |
| hexythiazox | | x | x | E | x | 7d | 1 |
| Portal XLO (0.4EC) | 21A | x | x | 2 pt. | x | 12h | 2 pt. |
| fenpyroximate | | x | x | E | x | 14d | 2 |
| Pyganic 5%EC | 3A | x | 4.5-15.6 fl. oz. | 4.5-15.6 fl. oz. | 4.5-15.6 fl. oz. | 12h | NA |
| pyrethrins | | x | F | u | u | 0d | 10 |
| Sevin XLR Plus (4F) | 1A | 2 qt. | 1-2 qt. | x | 1-2 qt. | 2d/6d | 10 qt. |
| carbaryl | | E | E | x | u | 7d | 5 |
| Vendex 50WP (RUP) | 12B | x | x | 1-2.5 lb. | x | 48h | 4 lb. |
| fenbutatin-oxide | | x | x | F | x | 28d | 2 |
| Verdepryn 100SL (0.83SL) | 28 | 8.2-11 fl. oz. | x | x | x | 4h | 33 fl. oz. |
| cyclaniliprole | | u | x | x | x | 7d | 3 |
| Zeal (72WP) | 10B | x | x | 2-3 oz. | x | 12h | 3 oz. |
| etoxazole | | x | x | E | x | 14d | 1 |

Footnotes: See how to read the spray schedule tables section at the beginning of the chapter.

Grape Bud Break to Prebloom - Diseases

Notes on disease management

- Begin fungicide applications at 1-3 inch new shoot growth; repeat at 7-10 day intervals or according to label instructions and environmental conditions.
- Powdery mildew:** Primary infections of powdery mildew can occur during this period, and it is critical to protect all new shoot growth. Sulfur is sufficient to control powdery mildew during this period unless varieties are sulfur sensitive. For sulfur sensitive varieties alternate FRAC 3 (Cevya, Mettle, Procure, Rally, TebuStar), FRAC 7 (Aprovia, Luna Privilege, Endura) and FRAC U13 (Gatten) fungicides. Using these fungicides in the third or fourth spray during this time period will also improve control of black rot.

Fungicide resistance alert

- The downy and powdery mildew pathogens are especially prone to fungicide resistance. Avoid

back-to-back applications of any one systemic fungicide class. See Fungicide Resistance Management, page 180, for information about fungicide resistance development in powdery and downy mildews. See generic fungicides table for product with the same active ingredient, page 251.

- Avoid using fungicides in FRAC 11 during this period.

Foundation program: This program contains products that are at a lower risk of resistance and serves as foundation for a grape disease management program.

Supplemental sprays: These fungicides provide greater activity against specific diseases and should be applied as a tank mix or in rotation with the foundation program when specific disease pressures are higher. As more green tissue develops, systemic fungicides have greater efficacy against specific pathogens.

Table 6-3. Grape Diseases - Bud Break Through Prebloom¹

| Product And Formulation Active Ingredient | FRAC Code ² | Anthraco- nose Foliar | Black Rot | Downy Mildew | Phomopsis Blight | Powdery Mildew | REI ³ PHI ⁴ | Max Amt ⁵ Max App ⁶ |
|--|---------------------------|-----------------------------|---------------------|--------------------|---------------------|-------------------|--------------------------------------|--|
| Abound (SC) | 11 | x | 10-15.5 fl. oz. | 10-15.5 fl. oz. | 10-15.5 fl. oz. | 10-15.5 fl. oz. | 4h | 90 fl. oz. |
| azoxystrobin | | x | E | E [r] | F | E [r] | 14d | varies |
| Aliette WDG | P07 (33) | x | x | 3-5 lb. | x | x | 12h | NA |
| aluminum tris | | x | x | E | x | x | 15d | 3 |
| Aprovia (EC) | 7 | 8.6-10.5 fl. oz. | 8.6-10.5 fl. oz. | x | 8.6-10.5 fl. oz. | 8.6-10.5 fl. oz. | 12h | 32 fl. oz. |
| benzovindiflupyr | | i | G-E | x | i | G-E | 21d | 3 |
| Captan 80 WDG | M | 1.25-2.5 lb. | 1.25-2.5 lb. | 1.25-2.5 lb. | 1.2-2.5 lb. | 1.2-2.5 lb. | 48h | 15 lb. |
| captan | | s(G) | s(F) | G | E | i | 0d | NA |
| Cevya | 3 | x | 4 fl. oz. | x | 3-4 fl. oz. | 3-4 fl. oz. | 12h | 8 fl. oz. |
| mefentrifluconazole | | x | E | x | G | G-E | 14d | 2 |
| Cuprofix Ultra 40D | M | x | 1.25-3 lb. | 1.2-3 lb. | 1.2-3 lb. | 1.2-3 lb. | 48 h | 50 lb. |
| copper sulfate | | x | F | F | F | F-i | NA | NA |
| Endura (70WG) | 7 | x | x | x | x | 4.5 oz. | 12h | 24 oz. |
| boscalid | | x | x | x | x | E | 14d | varies |
| Fervent 475SC | 3+7 | x | 8.5 fl. oz. | x | 8.5 fl. oz. | 8.5 fl. oz. | 12h | 26 fl. oz. |
| isofetamid+tebuconazole | | x | u | x | u | E | 14d | NA |
| Flint Extra | 11 | x | 3.5-3.8 fl. oz. | 3.8 fl. oz. | 3.5-3.8 fl. oz. | 3-3.5 fl. oz. | 12h | 23 fl. oz. |
| trifloxystrobin | | x | E | s | F | E | 14d | 6 |
| Forum | 40 | x | x | 6 oz. | x | x | 12h | 24 oz. |
| dimethomorph | | x | x | E | x | x | 14d | 4 |

(Continued)

Table 6-3. Grape Diseases - Bug Break Through Prebloom¹ (continued)

| Product And Formulation Active Ingredient | FRAC Code ² | Anthraco- nose Foliar | Black Rot | Downy Mildew | Phomopsis Blight | Powdery Mildew | REI ³ PHI ⁴ | Max Amt ⁵ Max App ⁶ |
|--|---------------------------|-----------------------------|---------------------|-----------------|---------------------|----------------------|--------------------------------------|--|
| Fracture | M | x | x | x | x | 20.5-24.4 fl. oz. | 4h | NA |
| Banda de Lupinus albus doce (BLAD) | | x | x | x | x | E | 1d | 5 |
| Gatten | U13 | x | x | x | x | 6.4 fl. oz. | 12h | 1 lb. |
| flutianil | | x | x | x | x | G-E | 14d | 4 |
| Inspire Super (EW) | 3+9 | 16-20 fl. oz. | 16-20 fl. oz. | x | x | 16-20 fl. oz. | 12h | 80 fl. oz. |
| difenoconazole + cyprodinil | | E | E | x | x | G | 14d | NA |
| Kenja 400SC | 7 | 20-22 fl. oz. | x | x | x | 20-22 fl. oz. | 12h | 66 fl. oz. |
| isofetamid | | G | x | x | x | F | 14d | NA |
| Lifegard WG | M | x | 4.5 oz. | 4.5 oz. | 4.5 oz. | 4.5 oz. | 4h | NA |
| <i>Bacillus mycooides</i> isolate J | | x | x | F | x | F | 0d | NA |
| Luna Experience (SC) | 7+3 | x | 8-8.6 fl. oz. | x | 8.6 fl. oz. | 6-8.6 fl. oz. | 12h/5d6 | 34 fl. oz. |
| fluopyram + tebuconazol | | x | G | x | s | E | 14d | NA |
| Luna Privilege (SC) | 7 | x | 6-6.8 fl. oz. | x | x | 3.2-6.8 fl. oz. | 12h | 14 fl. oz. |
| fluopyram | | x | G | x | x | G | 7d | 2 |
| Luna Sensation (SC) | 7+11 | x | 5-7.6 fl. oz. | 7.6 fl. oz. | 5-7.6 fl. oz. | 4-7.6 fl. oz. | 12h | 28 fl. oz. |
| fluopyram + trifloxystrobin | | x | G | s | F-G | E | 14d | 6 |
| Merivon (2.09SC) | 7+11 | 4-5.5 fl. oz. | 4-5.5 fl. oz. | 4-5.5 fl. oz. | 4-5.5 fl. oz. | 4-5.5 fl. oz. | 12h | 33 fl. oz. |
| fluxapyroxad + pyraclostrobin | | E | G-E | s | G-E | E [r] | 14d | 6 |
| Mettle 125ME | 3 | 3-5 fl. oz. | 3-5 fl. oz. | x | x | 3-5 fl. oz. | 12h/7d7 | 10 oz. |
| tetraconazole | | E | E | x | x | E [r] | 14d | varies |
| Microthiol Disperss | M | x | x | x | 3-10 lb. | 3-10 lb. | 24h | NA |
| sulfur | | x | x | x | F | E | 0d | NA |
| Miravis Prime | 7+12 | 9.2-13.4 fl. oz. | 9.2-13.4 fl. oz. | x | 9.2-13.4 fl. oz. | 9.2-13.4 fl. oz. | 12h | 37 fl. oz. |
| pydiflumetofen+fludioxonil | | u | E | x | F-G | G-E | 14d | 2 |
| OSO 5% SC | 19 | x | x | x | x | 6.5-13 fl. oz. | 4h | 5 oz. |
| polyoxin D | | x | x | x | x | G | 0d | 6 |
| Pristine | 11+7 | 8-12.5 oz. | 8-12.5 oz. | 8-12.5 oz. | 8-12.5 oz. | 8-12.5 oz. | 12h/5d6 | 69 oz. |
| pyraclostrobin + boscalid | | E | E | E [r] | F | E | 14d | varies |
| Procure 480SC | 3 | x | x | x | x | 4-8 fl. oz. | 24h | 32 fl. oz. |
| triflumizole | | x | x | x | x | E [r] | 7d | 4 |
| ProPhyt | P07 (33) | x | x | 2-4 pt. | 2-4 pt. | x | 4h | NA |
| potassium phosphite | | x | x | G-E | G-E | x | 0d | NA |
| Quadris Top (SC) | 3+11 | 12-14 fl. oz. | 12-14 fl. oz. | 12-14 fl. oz. | 12-14 fl. oz. | 12-14 fl. oz. | 12h | 56 fl. oz. |
| difenoconazole + azoxystrobin | | E | E | F | F | E | 14d | NA |
| Quintec (2.08F) | 13 | x | x | x | x | 4-6.6 fl. oz. | 12h | 33 fl. oz. |
| quinoxifen | | x | x | x | x | E | 21d | 5 |
| Rally 40WSP | 3 | 3-5 oz. | 3-5 oz. | x | x | 3-5 oz. | 24h | 2 lb. |
| myclobutanil | | E | E | x | x | E [r] | 14d | NA |

(Continued)

Table 6-3. Grape Diseases - Bug Break Through Prebloom¹ (continued)

| Product And Formulation Active Ingredient | FRAC Code ² | Anthraco- nose Foliar | Black Rot | Downy Mildew | Phomopsis Blight | Powdery Mildew | REI ³ PHI ⁴ | Max Amt ⁵ Max App ⁶ |
|--|---------------------------|-----------------------------|--------------|---------------------|---------------------|----------------------|--------------------------------------|--|
| Ranman 400SC | 21 | x | x | E | x | x | 12h | 17 fl. oz. |
| cyazofamid | | x | x | 2.1-2.75 fl. oz. | x | x | 30d | 6 |
| Reason 500SC | 11 | x | x | 2.7 fl. oz. | x | x | 12h | 9 fl. oz. |
| fenamidone | | x | x | G [r] | x | x | 30d | NA |
| Revus | 40 | x | x | 8 fl. oz. | x | x | 4h | 32 fl. oz. |
| mandipropamid | | x | x | E | x | x | 14d | NA |
| Revus Top | 3+40 | 7 fl. oz. | 7 fl. oz. | 7 fl. oz. | 7 fl. oz. | 7 fl. oz. | 12h | 28 fl. oz. |
| difenoconazole + mandiprop- amid | | E | E | E | E | E | 14d | NA |
| Rhyme | 3 | x | 4-5 fl. oz. | x | x | 4-5 fl. oz. | 14d | 30 fl. oz. |
| flutriafol | | x | E | x | x | E | 12h | 6 |
| Ridomil Gold Copper | 4+M | x | x | 5 lb. (1 pack) | x | x | 48h | 8 lb. |
| mefenoxam + copper hydroxide | | x | x | E | x | x | 42d | 4 |
| Ridomil Gold MZ WG | 4+M | x | x | 2.5 lb. | x | x | 48h | 10 lb. |
| mefenoxam + mancozeb | | x | x | E | x | x | 66d | 4 |
| Roper DF | M | x | 1.5-4 | 1.5-4 | 1.5-4 | x | 24h | 24 lb. |
| mancozeb | | x | E | E | E | x | 66d | 6 |
| Sovran (50WG) | 11 | x | 3.2-4.8 oz. | 4-6.4 oz. | 3.2-4.8 oz. | 3.2-4.8 oz. | 12h | 26 oz. |
| kresoxim-methyl | | x | E | F [r] | F | E [r] | 14d | 4 |
| Sulforix | M | x | x | x | 1-2 gal., 1 pt. | 1-2 gal., 1 pt. | 48h | NA |
| calcium polysulfide | | x | x | x | G-E | i | NA | 8 |
| Tanos | 11+27 | x | x | 8 oz. | x | x | 12h | 72 oz. |
| famoxadone + cymoxanil | | x | x | G [r] | x | x | 30d | 9 |
| Tebustar 45WSP | 3 | x | 4 oz. | x | x | 4 oz. | 12h | 2 lb. |
| tebuconazole | | x | E | x | x | E [r] | 0d | NA |
| Topguard EQ | 3+11 | x | 5-6 fl. oz. | 8 fl. oz. | 8 fl. oz. | 5-6 fl. oz. | 12h | 34 fl. oz. |
| azoxystrobin+flutriafol | | x | u | u | u | E | 14d | 6 |
| Topsin M WSB | 1 | x | 0.75-1.5 lb. | x | 0.75-1.5 lb. | 0.75-1.5 lb. | 2d | 6 lb. |
| thiophanate-methyl | | x | F | x | G | E | 7d | NA |
| Torino (SC) | U6 | x | x | x | x | 3.4-6.8 oz. | 4h | 7 oz. |
| cyflufenamid | | x | x | x | x | E | 3d | 1 or 2 |
| Vanguard WG | 9 | x | x | x | x | 10 oz. | 12h | 30 oz. |
| cyprodinil | | x | x | x | x | s | 7d | NA |
| Vivando 2.5F | U8 | x | x | x | x | 10.3-15.4 fl. oz. | 12h | 43 fl. oz. |
| metrafenone | | x | x | x | x | E | 14d | 3 |
| Zampro | 45+40 | x | x | 11-14 fl. oz. | x | x | 12h | 56 fl. oz. |
| ametoctradin + dimethomorph | | x | x | E | x | x | 14d | NA |
| Ziram 76DF | M3 | x | 3-4 lb. | 3-4 lb. | 3-4 lb. | x | 48h | 28 lb. |
| ziram | | x | E | G | G | x | 21d | NA |

Footnotes: See how to read the spray schedule tables section at the beginning of the chapter.

Grape Bud Break to Bloom - Insects

Notes on insect pest management

- **Grape phylloxera:** Admire Pro is soil-applied for systemic control. Use if there is a history of leaf-galling. Apply from bud swell until the first expanded leaf to be sure the chemical is available as soon as the roots begin to take up water, as it takes several weeks for the chemical to get to the leaves.
 - **Rose chafers:** May be present any time between 4- and 10-inch shoot growth and bloom.
 - **Redbanded leafroller and grape berry moth:** Pheromone traps for grape berry moth and red-banded leafroller indicate if they are present and help determine the need for control.
 - **Grape phylloxera (leaf form):** Control the root gall form of grape phylloxera by using rootstocks derived from American grapes. Native American grapes (Eastern U.S.) are highly resistant to this pest.
 - While all grapes do not benefit from insect pollination, protection of bees and other pollinators
- harvesting pollen from grape flowers is important. Avoid spraying insecticides that are toxic to pollinators during bloom.
 - **Scorpion 35SL:** Use low rate for foliar. Use high rate for soil application.
 - **Movento 2SC:** See label regarding adjuvants. Allow 30 days between applications. Movento is applied to leaves but moves to the roots. It is most effective if applied at this early stage of grape growth.
 - **Grape scale:** Not a common pest in most of the Midwest. In southern areas, flag scale-infested vines during dormant pruning. In early May begin weekly inspections of flagged vines for scale crawlers. Lift live adult scale covers and look for yellow moving crawlers (use a hand lens with 10x magnification). Protect canes by applying sprays every 10 days as long as you see moving crawlers (2-3 week crawler emergence period).
 - Other insecticide formulations may be available. See generic insecticides table for product with the same active ingredient, pages 279-280.

Table 6-4. Grape Insects - Bud Break Through Bloom¹

| Product And Formulation Active Ingredient | IRAC Code ² | Climbing Cutworm | Grape Cane Girdler/ Gallmaker | Grape Flea Beetle | Rose Chafer | REI ³ PHI ⁴ | Max Amt ⁵ Max App ⁶ |
|--|---------------------------|---------------------|-------------------------------------|----------------------|-------------------|--------------------------------------|--|
| Altacor eVo(35WDG) | 28 | 3-4.5 oz. | x | x | x | 4h | 4.6 oz. |
| chlorantraniliprole | | G | x | x | x | 1d | 3 |
| Assail 30SG | 4A | x | 2.5-5.3 oz. | x | 2.5-5.3 oz. | 12h | 10.6 oz. |
| acetamiprid | | x | u | x | E | 3d | 2 |
| Azera 0.21EC | 3A | 16-56 fl. oz. | 16-56 fl. oz. | 16-56 fl. oz. | 16-56 fl. oz. | 12h | NA |
| azadirachtin + pyrethrins | | u | u | u | u | 0d | 10 |
| Baythroid XL (1EC) (RUP) | 3A | 2.4 -3.2 fl. oz. | 2.4 -3.2 fl. oz. | 2.4 -3.2 fl. oz. | x | 12h | 12.8 fl. oz. |
| beta-cyfluthrin | | E | G | G | x | 3d | NA |
| Brigade 2EC (RUP) | 3A | 3.2-6.4 fl. oz. | x | x | x | 12h | 7 fl. oz. |
| bifenthrin | | G | x | x | x | 30d | NA |
| Brigade WSB (10WP) (RUP) | 3A | 8-16 oz. | x | x | x | 12h | 16 oz. |
| bifenthrin | | G | x | x | x | 30d | NA |
| Danitol 2.4EC (RUP) | 3A | 10.6-21.3 fl. oz. | 10.6-21.3 fl. oz. | 5.3 to 10.6 fl. oz. | 10.6-21.3 fl. oz. | 24h | 42.6 fl. oz. |
| fenpropathrin | | G | u | E | E | 21d | NA |
| Delegate WG (25WG) | 5 | 3-5 oz. | x | x | x | 4h | 19.5 oz. |
| spinetoram | | G | x | x | x | 3d | 5 |
| Entrust SC (2SC) | 5 | 4-8 fl. oz. | x | x | x | 4h | 23 fl. oz. |

(Continued)

Table 6-4. Grape Insects - Bud Break Through Bloom¹ (continued)

| Product And Formulation Active Ingredient | IRAC Code ² | Climbing Cutworm | Grape Cane Girdler/ Gallmaker | Grape Flea Beetle | Rose Chafer | REI ³ PHI ⁴ | Max Amt ⁵ Max App ⁶ |
|--|---------------------------|---------------------|-------------------------------------|----------------------|----------------|--------------------------------------|--|
| spinosad | | G | x | x | x | 3d | 5 |
| Imidan 70W | 1B | x | 1.3-2.1 lb. | 1.3-2.1 lb. | 1.3-2.1 lb. | 14d | 6.5 lb. |
| phosmet | | x | u | F | G | 7/14d | NA |
| Mustang Maxx (0.83EC) (RUP) | 3A | 2-4 fl. oz. | x | x | x | 12h | 24 fl. oz. |
| zeta-cypermethrin | | E | x | x | x | 1d | NA |
| Pyganic 5%EC | 3A | x | x | 4.5-15.6 fl. oz. | x | 12h | NA |
| pyrethrins | | x | x | F | x | 0d | 10 |
| Sevin XLR Plus (4F) | 1A | 2 qt. | x | 1-2 qt. | 1-2 qt. | 2d/6d | 10 qt. |
| carbaryl | | E | x | E | E | 7d | 5 |
| Surround WP (95WP) | UN | x | x | x | 25-50 lb. | 4h | NA |
| kaolin | | x | x | x | F | 0 | NA |
| Verdepryn 100SL (0.83SL) | 28 | 8.2-11 fl. oz. | x | x | x | 4h | 33 fl. oz. |
| cyclaniliprole | | u | x | x | x | 7d | 3 |

Footnotes: See how to read the spray schedule tables section at the beginning of the chapter.

Grape Prebloom To Shatter - Diseases

Notes on disease management

Prebloom through bloom

- Begin prebloom applications when shoots are 10- to 12- inches through the bloom period.
- Critical period for disease control: The period from immediate prebloom through four or five weeks after bloom (when grapes reach pea-size) is critical to control fruit infections by the black rot, powdery mildew, and the downy mildew pathogens.
- Although fruit of the most commonly planted varieties become resistant to powdery and downy mildews as they mature, the rachises (cluster stems) and leaves remain susceptible. Fungicide protection against powdery and downy mildews is therefore required throughout the growing season.

Bloom

- Begin bloom applications when the flower caps start to drop.
- If wet weather persists during bloom, or if the interval between the pre-bloom and shatter spray is greater than 10 days, a fungicide application during bloom should be applied.
- Downy mildew** is the most common disease in the Midwest. Initial infections can occur as early as bloom. Leaf infections may occur throughout the summer, so it may be necessary to protect susceptible varieties from bloom to post-harvest.

- Ripe rot** infections occur during this period, but symptoms don't appear until the grapes mature. FRAC 11 and 12 fungicides provide the best level of control. FRAC 7 fungicides can be used if disease pressure is low. A fungicide application at veraison may be required if disease pressure is high (wet and warm). See page 159 for additional information on ripe rot.
- Botrytis bunch rot:** A fungicide application during this period is critical on tight-clustered varieties (especially French hybrids or Vinifera) or in vineyards where Botrytis bunch rot has been a problem in the past. See Botrytis Bunch Rot, page 160.

Postharvest Interval (PHI) Reminder

- Pay close attention to the PHI on products that contain mancozeb (66 days) or mefenoxam (60 days).

Fungicide Resistance Alert

- The downy and powdery mildew pathogens are especially prone to fungicide resistance. Avoid back-to-back applications of any one systemic fungicide class (i.e., FRAC 3, 7, or 11).
- See Fungicide Resistance Management, page 180, for information about fungicide resistance development in powdery and downy mildews. See generic fungicides table for product with the same active ingredient, page 251.

Table 6-5. Grape Diseases – Prebloom Through Shatter¹

| Product And Formulation Active Ingredient | FRAC Code ² | Anthraco- nose Foliar | Black Rot | Botrytis Bunch Rot | Downy Mildew | Phomopsis Blight | Powdery Mildew | REI ³ PHI ⁴ | Max Amt ⁵ Max App ⁶ |
|--|---------------------------|-----------------------------|---------------------|--------------------------|--------------------|---------------------|----------------------|--------------------------------------|--|
| Abound (SC) | 11 | x | 10-15.5 fl. oz. | 10-15.5 fl. oz. | 10-15.5 fl. oz. | 10-15.5 fl. oz. | 10-15.5 fl. oz. | 4h | 90 fl. oz. |
| azoxystrobin | | x | E | s | E [r] | F | E [r] | 14d | varies |
| Aliette WDG | P07 (33) | x | x | x | 3-5 lb. | x | x | 12h | NA |
| aluminum tris | | x | x | x | E | x | x | 15d | 3 |
| Aprovia (EC) | 7 | 8.6-10.5 fl. oz. | 8.6-10.5 fl. oz. | x | x | 8.6-10.5 fl. oz. | 8.6-10.5 fl. oz. | 12h | 32 fl. oz. |
| benzovindiflupyr | | i | G-E | x | x | i | G-E | 21d | 3 |
| Captan 80 WDG | M | 1.25-2.5 lb. | 1.25-2.5 lb. | 2.5 lb | 1.25-2.5 lb. | 1.2-2.5 lb. | 1.2-2.5 lb. | 48h | 15 lb. |
| captan | | s(G) | s(F) | F | G | E | i | 0d | NA |
| Cevya | 3 | x | 4 fl. oz. | x | x | 3-4 fl. oz. | 3-4 fl. oz. | 12h | 8 fl. oz. |
| mefentrifluconazole | | x | E | x | x | G | G-E | 14d | 2 |
| Cuprofix Ultra 40D | M | x | 1.25-3 lb. | x | 1.2-3 lb. | 1.2-3 lb. | 1.2-3 lb. | 48h | 50 lb. |
| copper sulfate | | x | F | x | F | F | F-i | x | |
| Elevate 50 WDG | 17 | x | x | 1 lb. | x | x | 1 lb. | 12h | 3 lb. |
| fenhexamid | | x | x | E | x | x | s | 0d | 3 |
| Endura (70WG) | 7 | x | x | 8 oz. | x | x | 4.5 oz. | 12h | 24 oz. |
| boscalid | | x | x | G | x | x | E | 14d | varies |
| Fervent 475SC | 3+7 | x | 8.5 fl. oz. | 8.5 fl. oz. | x | 8.5 fl. oz. | 8.5 fl. oz. | 12h | 26 fl. oz. |
| isofetamid+tebuconazole | | x | u | s | x | u | E | 14d | NA |
| Flint Extra | 11 | x | 3.5-3.8 fl. oz. | 3.8 fl. oz. | 3.8 fl. oz. | 3.5-3.8 fl. oz. | 3-3.5 fl. oz. | 12h | 23 fl. oz. |
| trifloxystrobin | | x | E | G | s | F | E | 14d | 6 |
| Forum | 40 | x | x | x | 6 oz. | x | x | 12h | 24 oz. |
| dimethomorph | | x | x | x | E | x | x | 14d | 4 |
| Fracture | M | x | x | 24.4-36.6 fl. oz. | x | x | 20.5-24.4 fl. oz. | 4h | NA |
| Banda de Lupinus albus doce (BLAD) | | x | x | E | x | x | E | 1d | 5 |
| Gatten | U13 | x | x | x | x | x | 6.4 fl. oz. | 12h | 1 lb. |
| flutianil | | x | x | x | x | x | G-E | 14d | 4 |
| Inspire Super (EW) | 3+9 | 16-20 fl. oz. | 16-20 fl. oz. | 16-20 fl. oz. | x | x | 16-20 fl. oz. | 12h | 80 fl. oz. |
| difenoconazole + cyprodinil | | E | E | E | x | x | G | 14d | NA |
| Intuity (SC) | 11 | x | x | 6 fl. oz. | x | x | 6 fl. oz. | 12h | 18 fl. oz. |
| mandestrobin | | x | x | G-E | x | x | s | 10d | 3 |
| Kenja 400SC | 7 | 20-22 fl. oz. | x | 20-22 fl. oz. | x | x | 20-22 fl. oz. | 12h | 66 fl. oz. |
| isofetamid | | G | x | F | x | x | F | 14d | NA |
| Lifegard WG | M | x | 4.5 oz. | x | 4.5 oz. | 4.5 oz. | 4.5 oz. | 4h | NA |
| <i>Bacillus mycooides</i> isolate J | | x | x | x | F | x | F | 0d | NA |

(Continued)

Table 6-5. Grape Diseases – Prebloom Through Shatter¹ (continued)

| Product And Formulation Active Ingredient | FRAC Code ² | Anthraco- nose Foliar | Black Rot | Botrytis Bunch Rot | Downy Mildew | Phomopsis Blight | Powdery Mildew | REI ³ PHI ⁴ | Max Amt ⁵ Max App ⁶ |
|--|---------------------------|-----------------------------|---------------------|--------------------------|------------------|---------------------|---------------------|--------------------------------------|--|
| Luna Experience (SC) | 7+3 | x | 8-8.6 fl. oz. | 8-8.6 fl. oz. | x | 8.6 fl. oz. | 6-8.6 fl. oz. | 12h/ 5d6 | 34 fl. oz. |
| fluopyram + tebuconazol | | x | G | E | x | s | E | 14d | NA |
| Luna Privilege (SC) | 7 | x | 6-6.8 fl. oz. | 6-6.8 fl. oz. | x | x | 3.2-6.8 fl. oz. | 12h | 14 fl. oz. |
| fluopyram | | x | G | E | x | x | G | 7d | 2 |
| Luna Sensation (SC) | 7+11 | x | 5-7.6 fl. oz. | 5-7.6 fl. oz. | 7.6 fl. oz. | 5-7.6 fl. oz. | 4-7.6 fl. oz. | 12h | 28 fl. oz. |
| fluopyram + trifloxystrobin | | x | G | G-E | s | F-G | E | 14d | 6 |
| Merivon (2.09SC) | 7+11 | 4-5.5 fl. oz. | 4-5.5 fl. oz. | 4-5.5 fl. oz. | 4-5.5 fl. oz. | 4-5.5 fl. oz. | 4-5.5 fl. oz. | 12h | 33 fl. oz. |
| fluxapyroxad + pyraclos- trobin | | E | G-E | s | s | G-E | E [r] | 14d | 6 |
| Meteor | 2 | x | x | 1-2 pts. | x | x | x | 48h | x |
| lprodione | | x | x | G-E | x | x | x | 7d | 1 or 4 |
| Mettle 125ME | 3 | 3-5 fl. oz. | 3-5 fl. oz. | x | x | x | 3-5 fl. oz. | 12h/7d7 | 10 oz. |
| tetraconazole | | E | E | x | x | x | E [r] | 14d | varies |
| Microthiol Disperss | M | x | x | x | x | 3-10 lb. | 3-10 lb. | 24h | NA |
| sulfur | | x | x | x | x | F | E | 0d | NA |
| Miravis Prime | 7+12 | 9.2-13.4 fl. oz. | 9.2-13.4 fl. oz. | 10.3-13.4 fl. oz. | x | 9.2-13.4 fl. oz. | 9.2-13.4 fl. oz. | 12h | 37 fl. oz. |
| pydiflumetofen+fludioxonil | | u | E | G-E | x | F-G | G-E | 14d | 2 |
| Orius 20AQ | 3 | x | 8.6 oz. | x | x | x | 8.6 oz. | 12h | 68.8 fl. oz. |
| tebuconazole | | x | E | x | x | x | E | 14d | NA |
| OSO 5% SC | 19 | x | x | 6.5-13 fl. oz. | x | x | 6.5-13 fl. oz. | 4h | 5 oz. |
| polyoxin D | | x | x | G | x | x | G | 0d | 6 |
| Pristine | 11+7 | 8-12.5 oz. | 8-12.5 oz. | 8-23 oz. | 8-12.5 oz. | 8-12.5 oz. | 8-12.5 oz. | 12h/ 5d | 69 oz. |
| pyraclostrobin + boscalid | | E | E | G | E [r] | F | E | 14d | varies |
| Procure 480SC | 3 | x | x | x | x | x | 4-8 fl. oz. | 24h | 32 fl. oz. |
| triflumizole | | x | x | s | x | x | E [r] | 7d | 4 |
| ProPhyt | P07 (33) | x | x | x | 2-4 pt | 2-4 pt | x | 4h | NA |
| potassium phosphite | | x | x | x | G-E | G-E | x | 0d | NA |
| Quadris Top (SC) | 3+11 | 12-14 fl. oz. | 12-14 fl. oz. | 12-14 fl. oz. | 12-14 fl. oz. | 12-14 fl. oz. | 12-14 fl. oz. | 12h | 56 fl. oz. |
| difenoconazole + azox- ystrobin | | E | E | s | F | F | E | 14d | NA |
| Quintec (2.08F) | 13 | x | x | x | x | x | 4-6.6 fl. oz. | 12h | 33 fl. oz. |
| quinoxifen | | x | x | x | x | x | E | 21d | 5 |
| Rally 40WSP | 3 | 3-5 oz. | 3-5 oz. | x | x | x | 3-5 oz. | 24h | 2 lb. |
| myclobutanil | | E | E | x | x | x | E[r] | 14d | NA |

(Continued)

Table 6-5. Grape Diseases – Prebloom Through Shatter¹ (continued)

| Product And Formulation Active Ingredient | FRAC Code ² | Anthraco- nose Foliar | Black Rot | Botrytis Bunch Rot | Downy Mildew | Phomopsis Blight | Powdery Mildew | REI ³ PHI ⁴ | Max Amt ⁵ Max App ⁶ |
|--|---------------------------|-----------------------------|-----------------|--------------------------|---------------------|---------------------|--------------------|--------------------------------------|--|
| Ranman 400SC | 21 | x | x | x | 2.1-2.75 fl. oz. | x | x | 12h | 17 fl. oz. |
| cyazofamid | | x | x | x | E | x | x | 30d | 6 |
| Reason 500SC | 11 | x | x | x | 2.7 fl. oz. | x | x | 12h | 9 fl. oz. |
| fenamidone | | x | x | x | G[r] | x | x | 30d | NA |
| Revus | 40 | x | x | x | 8 fl. oz. | x | x | 4h | 32 fl. oz. |
| mandipropamid | | x | x | x | E | x | x | 14d | NA |
| Revus Top | 3+40 | 7 fl. oz. | 7 fl. oz. | x | 7 fl. oz. | 7 fl. oz. | 7 fl. oz. | 12h | 28 fl. oz. |
| difenoconazole + mandipropamid | | E | E | x | E | E | E | 14d | NA |
| Ridomil Gold Copper | 4+M | x | x | x | 5 lb. (1 pack) | x | x | 48h | 8 lb. |
| mefenoxam + copper hydroxide | | x | x | x | E | x | x | 42d | 4 |
| Rhyme | 3 | x | 4-5 fl. oz. | x | x | x | 4-5 fl. oz. | 14d | 30 fl. oz. |
| flutriafol | | x | E | x | x | x | E | 12h | 6 |
| Ridomil Gold MZ WG | 4+M | x | x | x | 2.5 lb. | x | x | 48h | 10 lb. |
| mefenoxam + mancozeb | | x | x | x | E | x | x | 66d | 4 |
| Roper DF | M | x | 1.5-4 | 1.5-4 | 1.5-4 | 1.5-4 | x | 24h | 24 lb. |
| mancozeb | | x | E | i | E | E | x | 66d | 6 |
| Rovral 4 F | 2 | x | x | 1-2 pt. | x | x | x | 48h | 2 or 8 pt. |
| iprodione | | x | x | G | x | x | x | 7d | 1 or 4 |
| Scala SC | 9 | x | x | 18 fl. oz. | x | x | x | 12h | 36 fl. oz. |
| pyrimethanil | | x | x | G | x | x | x | 7d | NA |
| Sovran (50WG) | 11 | x | 3.2-4.8 oz. | 3.2-6.4 oz. | 4-6.4 oz. | 3.2-4.8 oz. | 3.2-4.8 oz. | 12h | 26 oz. |
| kresoxim-methyl | | x | E | s | F [r] | F | E [r] | 14d | 4 |
| Sulforix | M | x | x | x | x | 1-2 gal., 1 pt. | 1-2 gal., 1 pt. | 48h | NA |
| calcium polysulfide | | x | x | x | x | G-E | i | NA | 8 |
| Switch 62.5 WG | 9+12 | x | x | 11-14 oz. | x | x | x | 12h | 56 oz. |
| cyprodinil + fludioxonil | | x | x | G | x | x | x | 7d | NA |
| Tanos | 11+27 | x | x | x | 8 oz. | x | x | 12h | 72 oz. |
| famoxadone + cymoxanil | | x | x | x | G [r] | x | x | 30d | 9 |
| Tebustar 45WSP | 3 | x | 4 oz. | x | x | x | 4 oz. | 12h | 2 lb. |
| tebuconazole | | x | E | x | x | x | E [r] | 0d | NA |
| Topguard EQ | 3+11 | x | 5-6 fl. oz. | 8 fl. oz. | 8 fl. oz. | 8 fl. oz. | 5-6 fl. oz. | 12h | 34 fl. oz. |
| azoxystrobin + flutriafol | | x | u | s | u | u | E | 14d | 6 |
| Topsin M WSB | 1 | x | 0.75-1.5 lb. | x | x | 0.75-1.5 lb. | 0.75-1.5 lb. | 2d | 6 lb. |
| thiophanate-methyl | | x | F | x | x | G | E | 7d | NA |

(Continued)

Table 6-5. Grape Diseases – Prebloom Through Shatter¹ (continued)

| Product And Formulation Active Ingredient | FRAC Code ² | Anthraco- nose Foliar | Black Rot | Botrytis Bunch Rot | Downy Mildew | Phomopsis Blight | Powdery Mildew | REI ³ PHI ⁴ | Max Amt ⁵ Max App ⁶ |
|--|---------------------------|-----------------------------|--------------|--------------------------|------------------|---------------------|----------------------|--------------------------------------|--|
| Torino (SC) | U6 | x | x | x | x | x | 3.4-6.8 oz. | 4h | 7 oz. |
| cyflufenamid | | x | x | x | x | x | E | 3d | 1 or 2 |
| Vanguard WG | 9 | x | x | 10 oz. | x | x | 10 oz. | 12h | 30 oz. |
| cyprodinil | | x | x | G-E | x | x | s | 7d | NA |
| Vivando 2.5F | U8 | x | x | x | x | x | 10.3-15.4 fl. oz. | 12h | 43 fl. oz. |
| metrafenone | | x | x | x | x | x | E | 14d | 3 |
| Zampro | 45+40 | x | x | x | 11-14 fl. oz. | x | x | 12h | 56 fl. oz. |
| ametoctradin + dimetho- morph | | x | x | x | E | x | x | 14d | NA |
| Ziram 76DF | M3 | x | 3-4 lb. | 3-4 lb. | 3-4 lb. | 3-4 lb. | x | 48h | 28 lb. |
| ziram | | x | E | s | G | G | x | 21d | NA |

Footnotes: See how to read the spray schedule tables section at the beginning of the chapter.

Grape Shatter To Veraison (Berry Coloring) - Insects

Apply when unfertilized berries fall from clusters, about 7-10 days after bloom or 7-10 days after last spray.

Notes on insect pest management

- **Grape rootworm:** Occasional problems from grape rootworm (adult beetles) are also controlled by Sevin, Imidan, Danitol, Baythroid, or Brigade
- **Redbanded leafroller:** Although adult moths are commonly caught in traps, the larvae of this pest are not common in grapes in the Midwest.
- **Leafhoppers (including sharpshooters):** Examining the undersides of grape leaves indicates if leafhoppers are present.

applied for grape berry moth control. When found, grape rootworm is typically a perimeter problem, low in the canopy.

Table 6-6. Grape Insects – Shatter Through Veraison¹

| Product And Formulation Active Ingredient | IRAC Code ² | Grape berry- Moth | Grape Phylloxera | Japanese Beetle | Leafhopper | Red- banded Leaf- roller | Rose Chafer | Spider Mite | REI ³ PHI ⁴ | Max Amt ⁵ Max App ⁶ |
|--|---------------------------|-------------------------|------------------------|--------------------|---|-----------------------------------|----------------|---------------------|--------------------------------------|--|
| Actara (25WDG) | 4A | x | x | 1.5-3.5 oz. | 1.5-3.5 oz. | x | x | x | 12h | 7 oz. |
| thiamethoxam | | x | x | G | G | x | x | x | 5d | NA |
| Admire Pro (4.6F) | 4A | x | 7-14 fl. oz. (soil) | x | 7-14 fl. oz. (soil); 1-1.4 fl. oz. (foliar) | x | x | x | 12h | 2.8/14 fl. oz. |
| imidacloprid | | x | G | x | E | x | x | x | 0/30d | NA |
| Agri-Mek SC (0.7SC) (RUP) | 6 | x | x | x | 1.75-3.5 fl. oz. | x | x | 1.75-3.5 fl. oz. | 12h/ 4d | 7 fl. oz. |
| abamectin | | x | x | x | F | x | x | G | 28d | 2 |
| Altacor eVo (35WDG) | 28 | 2-4.5 oz. | x | 3-4.5 oz. | x | x | x | x | 4h | 4.6 oz. |
| chlorantraniliprole | | E | x | G | x | x | x | x | 1d | 3 |

(Continued)

Table 6-6. Grape Insects - Shatter Through Veraison¹ (continued)

| Product And Formulation Active Ingredient | IRAC Code ² | Grape berry-Moth | Grape Phylloxera | Japanese Beetle | Leafhopper | Red-banded Leaf-roller | Rose Chafer | Spider Mite | REI ³ PHI ⁴ | Max Amt ⁵ Max App ⁶ |
|---|------------------------|--------------------|---------------------|----------------------|---|------------------------|-------------------|-------------------|-----------------------------------|---|
| Apollo SC (ISC) | 10A | x | x | x | x | x | x | 4-8 oz. | 12h/14d | NA |
| clofentezine | | x | x | x | x | x | x | E | 21d | 1 |
| Assail 30SG | 4A | 2.5-5.3 oz. | 2.5-5.3 oz. | 2.5-5.3 oz. | 2.5-5.3 oz. | x | 2.5-5.3 oz. | x | 12h | 10.6 oz. |
| acetamiprid | | u | G | G | E | x | E | x | 3d | 2 |
| Avaunt eVo (30WDG) | 22 | 5-6 oz. | x | 3.5-6 oz. | 5-6 oz. | x | x | x | 12h | 12 oz. |
| indoxacarb | | G | x | G | s | x | x | x | 7d | 2 |
| Azera 0.21EC | 3A | 16-56 fl. oz. | 16-56 fl. oz. | 16-56 fl. oz. | 16-56 fl. oz. | 16-56 fl. oz. | 16-56 fl. oz. | 16-56 fl. oz. | 12h | NA |
| azadirachtin + pyrethrins | | u | u | u | u | u | u | u | 0d | 10 |
| Baythroid XL (1EC) (RUP) | 3A | 2.4-3.2 fl. oz. | x | x | 1.6-3.2 fl. oz. | x | x | x | 12h | 12.8 fl. oz. |
| beta-cyfluthrin | | E | x | x | G | x | x | x | 3d | NA |
| BeetleGONE! ag | 11 | x | x | 1-175 lb. | x | x | x | x | 4h | NA |
| <i>B.t. galleriae</i> | | x | x | G | x | x | x | x | 0d | NA |
| Belay (2.13SC) | 4A | 6 fl. oz. (foliar) | 6-12 fl. oz. (soil) | 2-4 fl. oz. (foliar) | 6-12 fl. oz. (soil); 2-4 fl. oz. (foliar) | x | x | x | 12h | 12 fl. oz. |
| clothianidin | | F | s | F | E | x | x | x | 0/30d | 1 |
| Brigade 2EC (RUP) | 3A | 3.2-6.4 fl. oz. | x | 3.2-6.4 fl. oz. | 3.2-6.4 fl. oz. | x | x | 6.4 fl. oz. | 12h | 7 fl. oz. |
| bifenthrin | | G | x | G | G | x | x | u | 30d | NA |
| Brigade WSB (10WP) (RUP) | 3A | 8-16 oz. | x | 8-16 oz. | 8-16 oz. | x | x | 16 oz. | 12h | 16 oz. |
| bifenthrin | | G | x | G | G | x | x | u | 30d | NA |
| <i>B.t.</i> (Agree, Dipel, etc.) | 11A | 1-2 lb. | x | x | x | 1-2 lb. | x | x | 4h | NA |
| <i>Bacillus thuringiensis</i> | | F | x | x | x | F | x | x | 0d | NA |
| Closer SC (2SC) | 4C | x | x | x | 2.75-5.75 fl. oz. | x | x | x | 12h | 17 fl. oz. |
| sulfoxaflor | | x | x | x | E | x | x | x | 7d | 4 |
| Danitol 2.4EC (RUP) | 3A | 10.6-21.3 fl. oz. | 10.6-21.3 fl. oz. | 10.6-21.3 fl. oz. | 5.3-10.6 fl. oz. | 10.6-21.3 fl. oz. | 10.6-21.3 fl. oz. | 10.6-21.3 fl. oz. | 24h | 42.6 fl. oz. |
| fenpropathrin | | E | E | E | G | u | E | G | 21d | NA |
| Delegate WG (25WG) | 5 | 3-5 oz. | x | x | x | 3-5 oz. | x | x | 4h | 19.5 oz. |
| spinetoram | | E | x | x | x | E | x | x | 3d | 5 |
| Dibrom 8E (RUP) | 1B | x | x | x | 0.5-0.6 pt. | x | x | x | 48h | 6 pt. |
| naled | | x | x | x | u | x | x | x | 10d | NA |
| Entrust SC (2SC) | 5 | 4-8 fl. oz. | x | x | x | 4-8 fl. oz. | x | x | 4h | 23 fl. oz. |
| spinosad | | G | x | x | x | G | x | x | 3d | 5 |

(Continued)

Table 6-6. Grape Insects - Shatter Through Veraison¹ (continued)

| Product And Formulation Active Ingredient | IRAC Code ² | Grape berry-Moth | Grape Phylloxera | Japanese Beetle | Leafhopper | Red-banded Leaf-roller | Rose Chafer | Spider Mite | REI ³ PHI ⁴ | Max Amt ⁵ Max App ⁶ |
|---|------------------------|-------------------|---------------------|-------------------|-------------------|------------------------|-------------|------------------|-----------------------------------|---|
| Envidor 2SC | 23 | x | x | x | x | x | x | 16-34 fl. oz. | 12h | 34 fl. oz. |
| spirodiclofen | | x | x | x | x | x | x | E | 14d | 1 |
| Imidan 70W | 1B | 1.3-2.1 lb. | x | 1.3-2.1 lb. | 1.3-2.1 lb. | 1.3-2.1 lb. | 1.3-2.1 lb. | x | 14d | 6.5 lb. |
| phosmet | | G | x | G | G | G | G | x | 7/14d | NA |
| Intrepid 2F | 18 | 8-16 fl. oz. | x | x | x | 8-16 fl. oz. | x | x | 4h | 48 fl. oz. |
| methoxyfenozide | | E | x | x | x | G | x | x | 30d | NA |
| Kanemite 15SC | 20B | x | x | x | x | x | x | 21-31 fl. oz. | 12h | 62 fl. oz. |
| acequinocyl | | x | x | x | x | x | x | F | 7d | 2 |
| Magister SC (1.7SC) | 21A | x | x | x | 32-36 fl. oz. | x | x | 32-36 fl. oz. | 12h | 36 fl. oz. |
| fenazaquin | | x | x | x | u | x | x | u | 7d | 1 |
| Malathion 5EC | 1B | x | x | 3 pt. | 3 pt. | x | x | 3 pt. | 72h | NA |
| malathion | | x | x | G | G | x | x | u | 3d | 2 |
| Movento (2SC) | 23 | x | 6-8 fl. oz. | x | x | x | x | 6-8 fl. oz. | 24h | 12.5 fl. oz. |
| spirotetramat | | x | E | x | x | x | x | s | 7d | NA |
| Mustang Maxx (0.83EC) (RUP) | 3A | 4 fl. oz. | x | 4 fl. oz. | 4 fl. oz. | x | x | x | 12h | 24 fl. oz. |
| zeta-cypermethrin | | E | x | E | E | x | x | x | 1d | NA |
| Nealta (1.67SC) | 25 | x | x | x | x | x | x | 13.7 fl. oz. | 12h | 28 fl. oz. |
| cyflumetofen | | x | x | x | x | x | x | G | 14d | 2 |
| Nexter SC (3.75SC) | 21A | x | x | x | 7.5-17 fl. oz. | x | x | 7.5-17 fl. oz. | 12h | NA |
| pyridaben | | x | x | x | G | x | x | G | 7d | 2 |
| Onager (1EC) | 10A | x | x | x | x | x | x | 12-24 oz. | 12h | 24 oz. |
| hexythiazox | | x | x | x | x | x | x | E | 7d | 1 |
| Platinum (755G) | 4A | x | 2.67-5.67 oz. | 2.67-5.67 oz. | 2.67-5.67 oz. | x | x | x | 12h | 5.67 oz. |
| thiamethoxam | | x | G | F | G | x | x | x | 60d | NA |
| Portal XLO (0.4EC) | 21A | x | x | x | 1-2 pt. | x | x | 2 pt. | 12h | 2 pt. |
| fenpyroximate | | x | x | x | F | x | x | E | 14d | 2 |
| PQZ (1.87SC) | 9B | x | x | x | 3.2 fl. oz. | x | x | x | 12h | 4.8 fl. oz. |
| pyrifluquinazon | | x | x | x | G | x | x | x | 3d | 2 |
| Pyganic 5%EC | 3A | x | x | 4.5-15.6 fl. oz. | 4.5-15.6 fl. oz. | 4.5-15.6 fl. oz. | x | 4.5-15.6 fl. oz. | 12h | NA |
| pyrethrins | | x | x | F | u | u | x | u | 0d | 10 |
| Scorpion 355L | 4A | 1.75-5.25 fl. oz. | 9.0-13.25 fl. oz.** | 1.75-5.25 fl. oz. | 1.75-5.25 fl. oz. | x | x | x | 12h | 21.25 fl. oz. |
| dinotefuran | | F | s | u | G | x | x | x | 1d | NA |

(Continued)

Table 6-6. Grape Insects - Shatter Through Veraison¹ (continued)

| Product And Formulation Active Ingredient | IRAC Code ² | Grape berry-Moth | Grape Phylloxera | Japanese Beetle | Leafhopper | Red-banded Leaf-roller | Rose Chafer | Spider Mite | REI ³ PHI ⁴ | Max Amt ⁵ Max App ⁶ |
|---|------------------------|------------------|------------------|-----------------|------------------|------------------------|-------------|-------------|-----------------------------------|---|
| Sevin XLR Plus (4F) | 1A | 2 qt. | x | 1-2 qt. | 1-2 qt. | 2 qt. | 1-2 qt. | x | 2d/6d | 10 qt. |
| carbaryl | | G | x | E | G | G | E | x | 7d | 5 |
| Sivanto Prime (1.67SC) | 4D | x | x | x | 7-14 fl. oz. | x | x | x | 4h | 28 fl. oz. |
| flupyradifurone | | x | x | x | G | x | x | x | 0/30d | NA |
| Surround WP (95WP) | UN | x | x | 25-50 lb. | 25-50 lb. | x | 25-50 lb. | x | 4h | NA |
| kaolin | | x | x | F | F | x | F | x | 0 | NA |
| Transform WG | 4C | x | x | x | 1.5-2.75 oz. | x | x | x | 24h | 8.5 oz. |
| sulfoxaflor | | x | x | x | E | x | x | x | 7d | 4 |
| Vendex 50WP (RUP) | 12B | x | x | x | x | x | x | 1-2.5 lb. | 48h | 4 lb. |
| fenbutatin-oxide | | x | x | x | x | x | x | F | 28d | 2 |
| Venom (70SG) | 4A | 1-3 oz. (foliar) | 5-7.5 oz. (soil) | x | 1-3 oz. (foliar) | x | x | x | 12h | 12 oz. |
| dinotefuran | | F | s | x | G | x | x | x | 1/28d | NA |
| Verdepryn 100SL (0.83SL) | 28 | 8.2-11 fl. oz. | x | 8.2-11 fl. oz. | x | 8.2-11 fl. oz. | x | x | 4h | 33 fl. oz. |
| cyclanilprole | | E | x | u | x | u | x | x | 7d | 3 |
| Zeal (72WP) | 10B | x | x | x | x | x | x | 2-3 oz. | 12h | 3 oz. |
| etoxazole | | x | x | x | x | x | x | E | 14d | 1 |

**soil applications use higher rate

Footnotes: See how to read the spray schedule tables section at the beginning of the chapter.

Grape Shatter To Veraison (Berry Coloring) - Diseases

Notes on disease management

- Begin shatter applications when unfertilized berries fall from clusters, about 7-10 days after bloom or 7-10 days after the last spray.
- First cover applications should follow shatter by 7-10 days. Thereafter, sprays for disease control can be applied every 10-14 days until veraison. If heavy rainfall occurs, shorten the interval between sprays. Refer to labels for application timing and harvest restrictions. After bloom the threat of Phomopsis infection is greatly reduced. Fruit remain susceptible to black rot, powdery mildew, and downy mildew until about 4-5 weeks after bloom. It is critical to maintain a fungicide program that controls all three of these diseases until about 4-5 weeks after bloom. At 4-5 weeks after bloom, the fruit should be resistant to black rot, powdery mildew, and downy mildew; however, the leaves and rachises (cluster stems) remain susceptible to both

powdery and downy mildew for the rest of the season. Therefore, fungicide protection against both diseases may be required throughout the growing season.

Postharvest interval (PHI) reminder

- Pay close attention to the PHI on products that contain mancozeb (66 days) or mefenoxam (60 days).

Fungicide resistance alert

- The downy and powdery mildew pathogens are especially prone to fungicide resistance. Avoid back-to-back applications of any one systemic fungicide class (i.e., FRAC 3, 7, 11, 21, 40).
- See Fungicide Resistance Management, page 160, for information about fungicide resistance development in powdery and downy mildews. See Generic Fungicides table for product(s) with the same active ingredient, page 251.

Table 6-7. Grape Diseases – Shatter To Veraison¹

| Product And Formulation Active Ingredient | FRAC Code ² | Anthraco- nose Foliar | Black Rot | Downy Mildew | Phomopsis Blight | Powdery Mildew | REI ³ PHI ⁴ | Max Amt ⁵ Max App ⁶ |
|--|---------------------------|-----------------------------|---------------------|--------------------|---------------------|----------------------|--------------------------------------|--|
| Abound (SC) | 11 | x | 10-15.5 fl. oz. | 10-15.5 fl. oz. | 10-15.5 fl. oz. | 10-15.5 fl. oz. | 4h | 90 fl. oz. |
| azoxystrobin | | x | E | E [r] | F | E [r] | 14d | varies |
| Aliette WDG | 33 | x | x | 3-5 lb. | x | x | 12h | NA |
| aluminum tris | | x | x | E | x | x | 15d | 3 |
| Aprovia (EC) | 7 | 8.6-10.5 fl. oz. | 8.6-10.5 fl. oz. | x | 8.6-10.5 fl. oz. | 8.6-10.5 fl. oz. | 12h | 32 fl. oz. |
| benzovindiflupyr | | i | G-E | x | i | G-E | 21d | 3 |
| Captan 80 WDG | M | 1.25-2.5 lb. | 1.25-2.5 lb. | 1.25-2.5 lb. | 1.2-2.5 lb. | 1.2-2.5 lb. | 48h | 15 lb |
| captan | | s(G) | s(F) | G | E | i | 0d | NA |
| Cevya | 3 | x | 4 fl. oz. | x | 3-4 fl. oz. | 3-4 fl. oz. | 12h | 8 fl. oz. |
| mefentrifluconazole | | x | E | x | G | G-E | 14d | 2 |
| Cuprofix Ultra 40D | M | x | 1.25-3 lb. | 1.2-3 lb. | 1.2-3 lb. | 1.2-3 lb. | 48 h | 50 lb. |
| copper sulfate | | x | F | F | F | F-i | x | NA |
| Endura (70WG) | 7 | x | x | x | x | 4.5 oz. | 12h | 24 oz. |
| boscalid | | x | x | x | x | E | 14d | varies |
| Fervent 475SC | 3+7 | x | 8.5 fl. oz. | x | 8.5 fl. oz. | 8.5 fl. oz. | 12h | 26 fl. oz. |
| isofetamid+tebuconazole | | x | u | x | u | E | 14d | NA |
| Flint Extra | 11 | x | 3.5-3.8 fl. oz. | 3.8 fl. oz. | 3.5-3.8 fl. oz. | 3-3.5 fl. oz. | 12h | 23 fl. oz. |
| trifloxystrobin | | x | E | s | F | E | 14d | 6 |
| Forum | 40 | x | x | 6 oz. | x | x | 12h | 24 oz. |
| dimethomorph | | x | x | E | x | x | 14d | 4 |
| Fracture | M | x | x | x | x | 20.5-24.4 fl. oz. | 4h | NA |
| Banda de Lupinus albus doce (BLAD) | | x | x | x | x | F | 1d | 5 |
| Gatten | U13 | x | x | x | x | 6.4 fl. oz. | 12h | 1 lb. |
| flutianil | | x | x | x | x | G-E | 14d | 4 |
| Inspire Super (EW) | 3+9 | 16-20 fl. oz. | 16-20 fl. oz. | x | x | 16-20 fl. oz. | 12h | 80 fl. oz. |
| difenoconazole + cyprodinil | | E | E | x | x | G | 14d | NA |
| Kenja 400SC | 7 | 20-22 fl. oz. | x | x | x | 20-22 fl. oz. | 12h | 66 fl. oz. |
| isofetamid | | G | x | x | x | F | 14d | NA |
| Lifegard WG | M | x | 4.5 oz. | 4.5 oz. | 4.5 oz. | 4.5 oz. | 4h | NA |
| <i>Bacillus mycooides</i> isolate J | | x | x | F | x | F | 0d | NA |
| Luna Experience (SC) | 7+3 | x | 8-8.6 fl. oz. | x | 8.6 fl. oz. | 6-8.6 fl. oz. | 12h/5d6 | 34 fl. oz. |
| fluopyram + tebuconazol | | x | G | x | s | E | 14d | NA |
| Luna Privilege (SC) | 7 | x | 6-6.8 fl. oz. | x | x | 3.2-6.8 fl. oz. | 12h | 14 fl. oz. |
| fluopyram | | x | G | x | x | G | 7d | 2 |
| Luna Sensation (SC) | 7+11 | x | 5-7.6 fl. oz. | 7.6 fl. oz. | 5-7.6 fl. oz. | 4-7.6 fl. oz. | 12h | 28 fl. oz. |
| fluopyram + trifloxystrobin | | x | G | s | F-G | E | 14d | 6 |

(Continued)

Table 6-7. Grape Diseases – Shatter To Veraison¹ (continued)

| Product And Formulation Active Ingredient | FRAC Code ² | Anthraco- nose Foliar | Black Rot | Downy Mildew | Phomopsis Blight | Powdery Mildew | REI ³ PHI ⁴ | Max Amt ⁵ Max App ⁶ |
|--|---------------------------|-----------------------------|------------------|------------------|---------------------|-------------------|--------------------------------------|--|
| Merivon (2.09SC) | 7+11 | 4-5.5 fl. oz. | 4-5.5 fl. oz. | 4-5.5 fl. oz. | 4-5.5 fl. oz. | 4-5.5 fl. oz. | 12h | 33 fl. oz. |
| fluxapyroxad + pyraclostrobin | | E | G-E | s | G-E | E [r] | 14d | 6 |
| Mettle 125ME | 3 | 3-5 fl. oz. | 3-5 fl. oz. | x | x | 3-5 fl. oz. | 12h/7d7 | 10 oz. |
| tetraconazole | | E | E | x | x | E [r] | 14d | varies |
| Microthiol Disperss | M | x | x | x | 3-10 lb. | 3-10 lb. | 24h | NA |
| sulfur | | x | x | x | F | E | 0d | NA |
| Miravis Prime | 7+12 | 9.2-13.4 fl. oz. | 9.2-13.4 fl. oz. | x | 9.2-13.4 fl. oz. | 9.2-13.4 fl. oz. | 12h | 37 fl. oz. |
| pydiflumetofen+fludioxonil | | u | E | x | F-G | G-E | 14d | 2 |
| OSO 5% SC | 19 | x | x | x | x | 6.5-13 fl. oz. | 4h | 5 oz. |
| polyoxin D | | x | x | x | x | G | 0d | 6 |
| Pristine | 11+7 | 8-12.5 oz. | 8-12.5 oz. | 8-12.5 oz. | 8-12.5 oz. | 8-12.5 oz. | 12h/5d6 | 69 oz. |
| pyraclostrobin + boscalid | | E | E | E [r] | F | E | 14d | varies |
| Procure 480SC | 3 | x | x | x | x | 4-8 fl. oz. | 24h | 32 fl. oz. |
| triflumizole | | x | x | x | x | E [r] | 7d | 4 |
| ProPhyt | 33 | x | x | 2-4 pt. | 2-4 pt. | x | 4h | NA |
| potassium phosphite | | x | x | G-E | G-E | x | 0d | NA |
| Quadris Top (SC) | 3+11 | 12-14 fl. oz. | 12-14 fl. oz. | 12-14 fl. oz. | 12-14 fl. oz. | 12-14 fl. oz. | 12h | 56 fl. oz. |
| difenoconazole + azoxystrobin | | E | E | F | F | E | 14d | NA |
| Quintec (2.08F) | 13 | x | x | x | x | 4-6.6 fl. oz. | 12h | 33 fl. oz. |
| quinoxifen | | x | x | x | x | E | 21d | 5 |
| Rally 40WSP | 3 | 3-5 oz. | 3-5 oz. | x | x | 3-5 oz. | 24h | 2 lb. |
| myclobutanil | | E | E | x | x | E[r] | 14d | NA |
| Ranman 400SC | 21 | x | x | 2.1-2.75 fl. oz. | x | x | 12h | 17 fl. oz. |
| cyazofamid | | x | x | E | x | x | 30d | 6 |
| Reason 500SC | 11 | x | x | 2.7 fl. oz. | x | x | 12h | 9 fl. oz. |
| fenamidone | | x | x | G[r] | x | x | 30d | NA |
| Revus | 40 | x | x | 8 fl. oz. | x | x | 4h | 32 fl. oz. |
| mandipropamid | | x | x | E | x | x | 14d | NA |
| Revus Top | 3+40 | 7 fl. oz. | 7 fl. oz. | 7 fl. oz. | 7 fl. oz. | 7 fl. oz. | 12h | 28 fl. oz. |
| difenoconazole + mandipropamid | | E | E | E | E | E | 14d | NA |
| Rhyme | 3 | x | 4-5 fl. oz. | x | x | 4-5 fl. oz. | 14d | 30 fl. oz. |
| flutriafol | | x | E | x | x | | 12h | 6 |
| Ridomil Gold Copper | 4+M | x | x | 5 lb. (1 pack) | x | x | 48h | 8 lb. |
| mefenoxam + copper hydroxide | | x | x | E | x | x | 42d | 4 |
| Ridomil Gold MZ WG | 4+M | x | x | 2.5 lb. | x | x | 48h | 10 lb. |
| mefenoxam + mancozeb | | x | x | E | x | x | 66d | 4 |
| Sovran (50WG) | 11 | x | 3.2-4.8 oz. | 4-6.4 oz. | 3.2-4.8 oz. | 3.2-4.8 oz. | 12h | 26 oz. |
| kresoxim-methyl | | x | E | F [r] | F | E [r] | 14d | 4 |

(Continued)

Table 6-7. Grape Diseases – Shatter To Veraison¹ (continued)

| Product And Formulation Active Ingredient | FRAC Code ² | Anthrachnose Foliar | Black Rot | Downy Mildew | Phomopsis Blight | Powdery Mildew | REI ³ PHI ⁴ | Max Amt ⁵ Max App ⁶ |
|--|---------------------------|------------------------|--------------|-----------------|---------------------|----------------------|--------------------------------------|--|
| Sulforix | M | x | x | x | 1-2 gal., 1 pt. | 1-2 gal., 1 pt. | 48h | NA |
| calcium polysulfide | | x | x | x | G-E | i | NA | 8 |
| Switch 62.5 WG | 9+12 | x | x | x | x | x | 12h | 56 oz. |
| cyprodinil + fludioxonil | | x | x | x | x | x | 7d | NA |
| Tanos | 11+27 | x | x | 8 oz. | x | x | 12h | 72 oz |
| famoxadone + cymoxanil | | x | x | G [r] | x | x | 30d | 9 |
| Tebustar 45WSP | 3 | x | 4 oz. | x | x | 4 oz | 12h | 2 lb. |
| tebuconazole | | x | E | x | x | E [r] | 0d | NA |
| Topguard EQ | 3+11 | x | 5-6 fl. oz. | 8 fl. oz. | 8 fl. oz. | 5-6 fl. oz. | 12h | 34 fl. oz. |
| azoxystrobin+flutriafol | | x | u | u | u | E | 14d | 6 |
| Topsin M WSB | 1 | x | 0.75-1.5 lb. | x | 0.75-1.5 lb. | 0.75-1.5 lb. | 2d | 6 lb. |
| thiophanate-methyl | | x | F | x | G | E | 7d | NA |
| Torino (SC) | U6 | x | x | x | x | 3.4-6.8 oz. | 4h | 7 oz. |
| cyflufenamid | | x | x | x | x | E | 3d | 1 or 2 |
| Vanguard WG | 9 | x | x | x | x | 10 oz. | 12h | 30 oz. |
| cyprodinil | | x | x | x | x | s | 7d | NA |
| Vivando 2.5F | U8 | x | x | x | x | 10.3-15.4 fl. oz. | 12h | 43 fl. oz. |
| metrafenone | | x | x | x | x | E | 14d | 3 |
| Zampro | 45+40 | x | x | 11-14 fl. oz. | x | x | 12h | 56 fl. oz. |
| ametoctradin + dimethomorph | | x | x | E | x | x | 14d | NA |
| Ziram 76DF | M3 | x | 3-4 lb. | 3-4 lb. | 3-4 lb. | x | 48h | 28 lb. |
| ziram | | x | E | G | G | x | 21d | NA |

Footnotes: See how to read the spray schedule tables section at the beginning of the chapter.

Grape Veraison to Harvest – Insects

Notes on insect pest management

- Continue to monitor for insect and mite pests and apply insecticide as needed. Refer to product labels for specific insects, rates, and harvest restrictions.
- **Multi-colored Asian lady beetle:** Scout vineyards several days before harvest to determine the abundance of multi-colored Asian lady beetle.
- **RESIDUE REMINDER:** Wettable powder formulations may leave visible residues on fruit at harvest.
- Pay close attention to the preharvest intervals on products.

Table 6-8. Grape Insect - Veraison Through Harvest¹

| Product And Formulation Active Ingredient | IRAC Code ² | Grape Berry Moth | Green June Beetle | Japanese Beetle | Leafhopper | Multi-colored Asian Lady Beetle | Spider Mite | Spotted-Wing Drosophila | Stink Bug | REI ³ PHI ⁴ | Max Amt ⁵ Max App ⁶ |
|---|------------------------|--------------------|-------------------|----------------------|---|---------------------------------|------------------|-------------------------|---------------|-----------------------------------|---|
| Actara (25WDG) | 4A | x | x | 1.5-3.5 oz. | 1.5-3.5 oz. | x | x | x | 4 oz. | 12h | 7 oz. |
| thiamethoxam | | x | x | G | G | x | x | x | G | 5d | NA |
| Admire Pro (4.6F) | 4A | x | x | x | 7-14 fl. oz. (soil); 1-1.4 fl. oz. (foliar) | x | x | x | x | 12h | 2.8/14 fl. oz. |
| imidacloprid | | x | x | x | E | x | x | x | x | 0/30d | NA |
| Agri-Mek SC (0.7SC) (RUP) | 6 | x | x | x | 1.75-3.5 fl. oz. | x | 1.75-3.5 fl. oz. | x | x | 12h/4d | 7 fl. oz. |
| abamectin | | x | x | x | F | x | G | x | x | 28d | 2 |
| Altacor eVo (35WDG) | 28 | 2-4.5 oz. | x | 3-4.5 oz. | x | x | x | x | x | 4h | 4.6 oz. |
| chlorantraniliprole | | E | x | G | x | x | x | x | x | 1d | 3 |
| Apollo SC (1SC) | 10A | x | x | x | x | x | 4-8 oz. | x | x | 12h/14d | NA |
| clofentezine | | x | x | x | x | x | E | x | x | 21d | 1 |
| Assail 30SG | 4A | 2.5-5.3 oz. | x | 2.5-5.3 oz. | 2.5-5.3 oz. | x | x | x | x | 12h | 10.6 oz. |
| acetamiprid | | u | x | G | E | x | x | x | x | 3d | 2 |
| Avaunt eVo (30WDG) | 22 | 5-6 oz. | x | 3.5-6 oz. | 5-6 oz. | x | x | x | x | 12h | 12 oz. |
| indoxacarb | | G | G | G | s | x | x | x | x | 7d | 2 |
| Azera 0.21EC | 3A | 16-56 fl. oz. | 16-56 fl. oz. | 16-56 fl. oz. | 16-56 fl. oz. | 16-56 fl. oz. | 16-56 fl. oz. | 16-56 fl. oz. | 16-56 fl. oz. | 12h | NA |
| azadirachtin + pyrethrins | | u | u | u | u | u | u | u | u | 0d | 10 |
| Baythroid XL (1EC) (RUP) | 3A | 2.4-3.2 fl. oz. | x | x | 1.6 -3.2 fl. oz. | x | x | 2.4-3.2 fl. oz. | x | 12h | 12.8 fl. oz. |
| beta-cyfluthrin | | E | x | x | G | x | x | E | x | 3d | NA |
| BeetleGONE! ag | 11 | x | x | 1-175 lb. | x | x | x | x | x | 4h | NA |
| <i>B.t. galleriae</i> | | x | x | G | x | x | x | x | x | 0d | NA |
| Belay (2.13SC) | 4A | 6 fl. oz. (foliar) | x | 2-4 fl. oz. (foliar) | 6-12 fl. oz. (soil); 2-4 fl. oz. (foliar) | 2-4 fl. oz. (foliar) | x | x | x | 12h | 12 fl. oz. |
| clothianidin | | F | x | F | E | E | x | x | x | 0/30d | 1 |
| Brigade 2EC (RUP) | 3A | 3.2-6.4 fl. oz. | x | 3.2-6.4 fl. oz. | 3.2-6.4 fl. oz. | x | 6.4 fl. oz. | x | x | 12h | 7 fl. oz. |
| bifenthrin | | G | x | G | G | x | u | x | x | 30d | NA |
| Brigade WSB (10WP) (RUP) | 3A | 8-16 oz. | x | 8-16 oz. | 8-16 oz. | x | 16 oz. | x | x | 12h | 16 oz. |
| bifenthrin | | G | x | G | G | x | u | x | x | 30d | NA |

(Continued)

Table 6-8. Grape Insect - Veraison Through Harvest¹ (continued)

| Product And Formulation Active Ingredient | IRAC Code ² | Grape Berry Moth | Green June Beetle | Japanese Beetle | Leafhopper | Multi-colored Asian Lady Beetle | Spider Mite | Spotted-Wing Drosophila | Stink Bug | REI ³ PHI ⁴ | Max Amt ⁵ Max App ⁶ |
|--|------------------------|-------------------|-------------------|-------------------|-------------------|---------------------------------|-------------------|-------------------------|-------------------|-----------------------------------|--|
| <i>B.t.</i> (Agree, Dipel, etc.) | 11A | 1-2 lb. | x | x | x | x | x | x | x | 4h | NA |
| <i>Bacillus thuringiensis</i> | | u | x | x | x | x | x | x | x | 0d | NA |
| Closer SC (2SC) | 4C | x | x | x | 2.75-5.75 fl. oz. | x | x | x | x | 12h | 17 fl. oz. |
| sulfoxaflor | | x | x | x | E | x | x | x | x | 7d | 4 |
| Danitol 2.4EC (RUP) | 3A | 10.6-21.3 fl. oz. | x | 10.6-21.3 fl. oz. | 5.3-10.6 fl. oz. | x | 10.6-21.3 fl. oz. | 10.6-21.3 fl. oz. | 10.6-21.3 fl. oz. | 24h | 42.6 fl. oz. |
| fenpropathrin | | E | x | E | G | x | G | E | G | 21d | NA |
| Delegate WG (25WG) | 5 | 3-5 oz. | x | x | x | x | x | x | x | 4h | 19.5 oz. |
| spinetoram | | E | x | x | x | x | x | E | x | 3d | 5 |
| Dibrom 8E (RUP) | 1B | x | x | x | 0.5-0.6 pt. | x | x | 0.5-0.6 pt. | x | 48h | 6 pt. |
| naled | | x | x | x | u | x | x | u | x | 10d | NA |
| Entrust SC (2SC) | 5 | 4-8 fl. oz. | x | x | x | x | x | x | x | 4h | 23 fl. oz. |
| spinosad | | G | x | x | x | x | x | G | x | 3d | 5 |
| Envidor 2SC | 23 | x | x | x | x | x | 16-34 fl. oz. | x | x | 12h | 34 fl. oz. |
| spiroticlofen | | x | x | x | x | x | E | x | x | 14d | 1 |
| Imidan 70W | 1B | 1.3-2.1 lb. | x | 1.3-2.1 lb. | 1.3-2.1 lb. | x | x | 1.3-2.1 lb. | x | 14d | 6.5 lb. |
| phosmet | | G | x | G | G | x | x | G | x | 7/ 14d | NA |
| Intrepid 2F | 18 | 8-16 fl. oz. | x | x | x | x | x | x | x | 4h | 48 fl. oz. |
| methoxyfenozide | | E | x | x | x | x | x | x | x | 30d | NA |
| Kanemite 15SC | 20B | x | x | x | x | x | 21-31 fl. oz. | x | x | 12h | 62 fl. oz. |
| acequinocyl | | x | x | x | x | x | F | x | x | 7d | 2 |
| Magister SC (1.7SC) | 21A | x | x | x | 32-36 fl. oz. | x | 32-36 fl. oz. | x | x | 12h | 36 fl. oz. |
| fenazaquin | | x | x | x | u | x | u | x | x | 7d | 1 |
| Malathion 5EC | 1B | x | x | 3 pt. | 3 pt. | x | 3 pt. | 3 pt. | x | 72h | NA |
| malathion | | x | x | G | G | x | u | G | x | 3d | 2 |
| Movento (2SC) | 23 | x | x | x | x | x | 6-8 fl. oz. | x | x | 24h | 12.5 fl. oz. |
| spirotetramat | | x | x | x | x | x | s | x | x | 7d | NA |
| Mustang Maxx (0.83EC) (RUP) | 3A | 4 fl. oz. | x | 4 fl. oz. | 4 fl. oz. | 2-4 fl. oz. | x | x | x | 12h | 24 fl. oz. |
| zeta-cypermethrin | | E | x | E | E | G | x | E | x | 1d | NA |
| Nealta (1.67SC) | 25 | x | x | x | x | x | 13.7 fl. oz. | x | x | 12h | 28 fl. oz. |
| cyflumetofen | | x | x | x | x | x | G | x | x | 14d | 2 |

(Continued)

Table 6-8. Grape Insect - Veraison Through Harvest¹ (continued)

| Product And Formulation Active Ingredient | IRAC Code ² | Grape Berry Moth | Green June Beetle | Japanese Beetle | Leafhopper | Multi-colored Asian Lady Beetle | Spider Mite | Spotted-Wing Drosophila | Stink Bug | REI ³ PHI ⁴ | Max Amt ⁵ Max App ⁶ |
|--|------------------------|-------------------|-------------------|-------------------|-------------------|---------------------------------|------------------|-------------------------|------------------|--------------------------------------|--|
| Nexter SC (3.75SC) | 21A | x | x | x | 7.5-17 fl. oz. | x | 7.5-17 fl. oz. | x | x | 12h | NA |
| pyridaben | | x | x | x | G | x | G | x | x | 7d | 2 |
| Onager (1EC) | 10A | x | x | x | x | x | 12-24 oz. | x | x | 12h | 24 oz. |
| hexythiazox | | x | x | x | x | x | E | x | x | 7d | 1 |
| Portal XLO (0.4EC) | 21A | x | x | x | 1-2 pt. | x | 2 pt. | x | x | 12h | 2 pt. |
| fenpyroximate | | x | x | x | F | x | E | x | x | 14d | 2 |
| PQZ (1.87SC) | 9B | x | x | x | 3.2 fl. oz. | x | x | x | x | 12h | 4.8 fl. oz. |
| pyrifluquinazon | | x | x | x | G | x | x | x | x | 3d | 2 |
| Pyganic 5%EC | 3A | x | x | 4.5-15.6 fl. oz. | 4.5-15.6 fl. oz. | x | 4.5-15.6 fl. oz. | 4.5-15.6 fl. oz. | 4.5-15.6 fl. oz. | 12h | NA |
| pyrethrins | | x | x | F | u | G | u | F | u | 0d | 10 |
| Scorpion 35SL | 4A | 1.75-5.25 fl. oz. | x | 1.75-5.25 fl. oz. | 1.75-5.25 fl. oz. | 1.75-5.25 fl. oz. | x | x | x | 12h | 21.25 fl. oz. |
| dinotfuran | | F | x | u | G | G | x | x | x | 1d | NA |
| Sevin XLR Plus (4F) | 1A | 2 qt. | 2 qt. | 1-2 qt. | 1-2 qt. | x | x | x | x | 2d/ 6d | 10 qt. |
| carbaryl | | G | E | E | G | x | x | x | x | 7d | 5 |
| Sivanto Prime (1.67SC) | 4D | x | x | x | 7-14 fl. oz. | x | x | x | x | 4h | 28 fl. oz. |
| flupyradifurone | | x | x | x | G | x | x | x | x | 0/ 30d | NA |
| Surround WP (95WP) | UN | x | 25-50 lb. | 25-50 lb. | 25-50 lb. | x | x | x | x | 4h | NA |
| kaolin | | x | u | F | F | x | x | x | x | 0 | NA |
| Transform WG | 4C | x | x | x | 1.5-2.75 oz. | x | x | x | x | 24h | 8.5 oz. |
| sulfoxaflor | | x | x | x | E | x | x | x | x | 7d | 4 |
| Vendex 50WP (RUP) | 12B | x | x | x | x | x | 1-2.5 lb. | x | x | 48h | 4 lb. |
| fenbutatin-oxide | | x | x | x | x | x | F | x | x | 28d | 2 |
| Venom (70SG) | 4A | 1-3 oz. (foliar) | x | x | 1-3 oz. (foliar) | 1-3 oz. (foliar) | x | x | x | 12h | 6 oz.; 7.5 oz.** |
| dinotefuran | | F | x | x | G | G | x | x | x | 1/ 28d | 1** |
| Verdepryn 100SL (0.83SL) | 28 | 8.2-11 fl. oz. | x | 8.2-11 fl. oz. | x | x | x | 8.2-11 fl. oz. | 8.2-11 fl. oz. | 4h | 33 fl. oz. |
| cyclaniliprole | | E | x | u | x | x | x | E | s | 7d | 3 |
| Zeal (72WP) | 10B | x | x | x | x | x | 2-3 oz. | x | x | 12h | 3 oz. |
| etoxazole | | x | x | x | x | x | E | x | x | 14d | 1 |

** denotes soil application

Footnotes: See how to read the spray schedule tables section at the beginning of the chapter.

Grape Veraison to Harvest - Diseases

Notes on disease management

- **Botrytis bunch rot:** See comments under Grape Bloom for Topsin M, Rovral, Vanguard, and Elevate. See Botrytis Bunch Rot, page 1160. Same as for Grape Bloom, pages 163-166.
- **Sour rot complex:** Mix Oxidate or Blight Ban 506 with an insecticide (for Drosophila control). See discussion on page 160.
- **Black rot:** Sprays for black rot should not be needed at this time.
- **Ripe rot:** A fungicide application at veraison may be required if disease pressure high (wet and warm). See page 159 for additional information on ripe rot.

Table 6-9. Grape Diseases - Veraison Through Harvest¹

| Product And Formulation Active Ingredient | FRAC Code ² | Anthraxnose Foliar | Anthraxnose Fruit rot | Black Rot | Botrytis Bunch Rot | Downy Mildew | Phomopsis Blight | Powdery Mildew | Bitter Rot | Ripe Rot | REI ³ PHI ⁴ | Max Amt ⁵ Max App ⁶ |
|--|------------------------|--------------------|-----------------------|-----------------|--------------------|-----------------|------------------|-------------------|------------|----------|--------------------------------------|--|
| Abound (SC) | 11 | x | x | 10-15.5 fl. oz. | 10-15.5 fl. oz. | 10-15.5 fl. oz. | 10-15.5 fl. oz. | 10-15.5 fl. oz. | x | x | 4h | 90 fl. oz. |
| azoxystrobin | | x | x | E | s | E [r] | F | E [r] | x | x | 14d | varies |
| Aliette WDG | 33 | x | x | x | x | 3-5 lb. | x | x | x | x | 12h | NA |
| aluminum tris | | x | x | x | E | x | x | x | x | x | 15d | 3 |
| Captan 80 WDG | M | 1.25-2.5 lb. | 1.25-2.5 lb. | 1.25-2.5 lb. | 2.5 lb. | 1.25-2.5 lb. | 1.2-2.5 lb. | 1.2-2.5 lb. | x | x | 48h | 15 lb. |
| captan | | s(G) | s(F) | F | G | E | i | s(G) | x | x | 0d | NA |
| Cevya | 3 | x | x | 4 fl. oz. | x | x | 3-4 fl. oz. | 3-4 fl. oz. | x | x | 12h | 8 fl. oz. |
| mefentrifluconazole | | x | x | E | x | x | G | G-E | x | x | 14d | 2 |
| Cuprofix Ultra 40D | M | x | x | 1.25-3 lb. | x | 1.2-3 lb. | 1.2-3 lb. | 1.2-3 lb. | x | x | 48 h | 50 lb. |
| copper sulfate | | x | x | F | x | F | F | F-i | x | x | x | NA |
| Elevate 50 WDG | 17 | x | x | x | 1 lb. | x | x | 1 lb. | x | x | 12h | 3 lb. |
| fenhexamid | | x | x | x | E | x | x | s | x | x | 0d | 3 |
| Endura (70WG) | 7 | x | x | x | 8 oz. | x | x | 4.5 oz. | x | x | 12h | 24 oz. |
| boscalid | | x | x | x | G | x | x | E | x | x | 14d | varies |
| Fervent 475SC | 3+7 | x | x | 8.5 fl. oz. | 8.5 fl. oz. | x | 8.5 fl. oz. | 8.5 fl. oz. | x | x | 12h | 26 fl. oz. |
| isofetamid+tebuconazole | | x | x | u | s | x | u | E | x | x | 14d | NA |
| Flint Extra | 11 | x | x | 3.5-3.8 fl. oz. | 3.8 fl. oz. | 3.8 fl. oz. | 3.5-3.8 fl. oz. | 3-3.5 fl. oz. | x | x | 12h | 23 fl. oz. |
| trifloxystrobin | | x | x | E | G | s | F | E | x | x | 14d | 6 |
| Forum | 40 | x | x | x | x | 6 oz. | x | x | x | x | 12h | 24 oz. |
| dimethomorph | | x | x | x | x | E | x | x | x | x | 14d | 4 |
| Fracture | M | x | x | x | 24.4-36.6 fl. oz. | x | x | 20.5-24.4 fl. oz. | x | x | 4h | NA |
| Banda de Lupinus albus doce (BLAD) | | x | x | x | E | x | x | E | x | x | 1d | 5 |

(Continued)

Table 6-9. Grape Diseases - Veraison Through Harvest¹ (continued)

| Product And Formulation Active Ingredient | FRAC Code ² | Anthraco- se Foliar | Anthraco- se Fruit rot | Black Rot | Botrytis Bunch Rot | Downy Mildew | Phomopsis Blight | Powdery Mildew | Bitter Rot | Ripe Rot | REI ³ PHI ⁴ | Max Amt ⁵ Max App ⁶ |
|--|------------------------|------------------------|---------------------------|------------------|-----------------------|------------------|---------------------|-------------------|------------|---------------|--------------------------------------|--|
| Gatten | U13 | x | x | x | x | x | x | 6.4 fl. oz. | x | x | 12h | 1 lb. |
| flutianil | | x | x | x | x | x | x | G-E | x | x | 14d | 4 |
| Inspire Super (EW) | 3+9 | 16-20 fl. oz. | 16-20 fl. oz. | 16-20 fl. oz. | 16-20 fl. oz. | x | x | 16-20 fl. oz. | x | x | 12h | 80 fl. oz. |
| difenoconazole + cyprodinil | | E | E | E | x | x | x | G | x | x | 14d | NA |
| Intuity (SC) | 11 | x | x | x | 6 fl. oz. | x | x | 6 fl. oz. | x | x | 12h | 18 fl. oz. |
| mandestrobin | | x | x | x | G-E | x | x | s | x | x | 10d | 3 |
| Kenja 400SC | 7 | 20-22 fl. oz. | 20-22 fl. oz. | x | 20-22 fl. oz. | x | x | 20-22 fl. oz. | x | 20-22 fl. oz. | 12h | 66 fl. oz. |
| isofetamid | | G | G | x | F | x | x | F | x | u | 14d | NA |
| Lifegard WG | M | x | x | x | x | 4.5 oz. | x | 4.5 oz. | x | x | 4h | NA |
| <i>Bacillus mycooides</i> isolate J | | x | x | x | x | F | x | F | x | x | 0d | NA |
| Luna Experience (SC) | 7+3 | x | x | 8-8.6 fl. oz. | 8-8.6 fl. oz. | x | 8.6 fl. oz. | 6-8.6 fl. oz. | x | x | 12h/ 5d6 | 34 fl. oz. |
| fluopyram + tebuconazol | | x | x | G | E | x | s | E | x | x | 14d | NA |
| Luna Privilege (SC) | 7 | x | x | 6-6.8 fl. oz. | 6-6.8 fl. oz. | x | x | 3.2-6.8 fl. oz. | x | x | 12h | 14 fl. oz. |
| fluopyram | | x | x | G | E | x | x | G | x | x | 7d | 2 |
| Luna Sensation (SC) | 7+11 | x | x | 5-7.6 fl. oz. | 5-7.6 fl. oz. | 7.6 fl. oz. | 5-7.6 fl. oz. | 4-7.6 fl. oz. | x | x | 12h | 28 fl. oz. |
| fluopyram + trifloxystrobin | | x | x | G | G-E | s | F-G | E | x | x | 14d | 6 |
| Merivon (2.09SC) | 7+11 | 4 to 5.5 fl. oz. | 4 to 5.5 fl. oz. | 4 to 5.5 fl. oz. | 4 to 5.5 fl. oz. | 4 to 5.5 fl. oz. | 4 to 5.5 fl. oz. | 4-5.5 fl. oz. | x | 4-5.5 fl. oz. | 12h | 33 fl. oz. |
| fluxapyroxad + pyraclostrobin | | E | G-E | E | s | s[r] | G-E | E [r] | x | s(G) | 14d | 6 |
| Meteor | 2 | x | x | x | 1.5-2 pts. | x | x | x | x | x | 48h | x |
| lprodione | | x | x | x | G-E | x | x | x | x | x | 7d | 1 or 4 |
| Mettle 125ME | 3 | 3-5 fl. oz. | 3-5 fl. oz. | 3-5 | x | x | x | 3-5 fl. oz. | x | x | 12h/ 7d7 | 10 oz. |
| tetraconazole | | E | E | u | x | x | x | E [r] | x | x | 14d | varies |
| Microthiol Disperss | M | x | x | x | x | x | 3-10 lb. | 3-10 lb. | x | x | 24h | NA |
| sulfur | | x | x | x | x | F | E | x | x | x | 0d | NA |
| Miravis Prime | 7+12 | 9.2-13.4 fl. oz. | 9.2-13.4 fl. oz. | 9.2-13.4 fl. oz. | 10.3-13.4 fl. oz. | x | 9.2-13.4 fl. oz. | 9.2-13.4 fl. oz. | x | x | 12h | 37 fl. oz. |
| pydiflumetofen+fludioxonil | | u | E | G-E | F-G | x | G-E | F-G | x | x | 14d | 2 |

(Continued)

Table 6-9. Grape Diseases - Veraison Through Harvest¹ (continued)

| Product And Formulation Active Ingredient | FRAC Code ² | Anthraxnose Foliar | Anthraxnose Fruit rot | Black Rot | Botrytis Bunch Rot | Downy Mildew | Phomopsis Blight | Powdery Mildew | Bitter Rot | Ripe Rot | REI ³ PHI ⁴ | Max Amt ⁵ Max App ⁶ |
|--|------------------------|--------------------|-----------------------|---------------|--------------------|-----------------|------------------|----------------|------------|----------|-----------------------------------|---|
| OSO 5% SC | 19 | x | x | x | 6.5-13 fl. oz. | x | x | 6.5-13 fl. oz. | x | x | 4h | 5 oz. |
| polyoxin D | | x | x | x | G | x | x | G | x | x | 0d | 6 |
| Pristine | 11+7 | 8-12.5 oz. | 8-12.5 oz. | 8-12.5 oz. | 8-23 oz. | 8-12.5 oz. | 8-12.5 oz. | 8-12.5 oz. | x | x | 12h/5d6 | 69 oz |
| pyraclostrobin + boscalid | | E | E | G | E [r] | F | E | u | x | x | 14d | varies |
| Procure 480SC | 3 | x | x | x | x | x | x | 4-8 fl. oz. | x | x | 24h | 32 fl. oz. |
| triflumizole | | x | x | s | x | x | E [r] | x | x | x | 7d | 4 |
| ProPhyt | 33 | x | x | x | x | 2-4 pt. | 2-4 pt. | x | x | x | 4h | NA |
| potassium phosphite | | x | x | x | G-E | G-E | x | x | x | x | 0d | NA |
| Quadris Top (SC) | 3+11 | 12-14 fl. oz. | 12-14 fl. oz. | 12-14 fl. oz. | 12-14 fl. oz. | 12-14 fl. oz. | 12-14 fl. oz. | 12-14 fl. oz. | x | x | 12h | 56 fl. oz. |
| difenoconazole + azoxystrobin | | E | E | s | F | s[r] | E | G | x | x | 14d | NA |
| Rally 40WSP | 3 | 3-5 oz. | 3-5 oz. | 3-5 oz. | x | x | x | 3-5 oz. | x | x | 24h | 2 lb. |
| myclobutanil | | E | E | x | x | x | E[r] | x | x | x | 14d | NA |
| Revus | 40 | x | x | x | x | 8 fl. oz. | x | x | x | x | 4h | 32 fl. oz. |
| mandipropamid | | x | x | x | E | x | x | x | x | x | 14d | NA |
| Revus Top | 3+40 | 7 fl. oz. | 7 fl. oz. | 7 fl. oz. | x | 7 fl. oz. | 7 fl. oz. | 7 fl. oz. | x | x | 12h | 28 fl. oz. |
| difenoconazole + mandipropamid | | E | E | E | x | E | E | F | x | x | 14d | NA |
| Rhyme | 3 | x | x | 4-5 fl. oz. | x | x | x | 4-5 fl. oz. | x | x | 14d | 30 fl. oz. |
| flutriafol | | x | x | E | x | x | x | E | x | x | 12h | 6 |
| Scala SC | 9 | x | x | x | 18 fl. oz. | x | x | x | x | x | 12h | 36 fl. oz. |
| pyrimethanil | | x | x | x | G | x | x | x | x | x | 7d | NA |
| Sovran (50WG) | 11 | x | x | 3.2-4.8 oz. | 3.2-6.4 oz. | 4-6.4 oz. | 3.2-4.8 oz. | 3.2-4.8 oz. | x | x | 12h | 26 oz. |
| kresoxim-methyl | | x | x | s | E-F [r] | E-F [r] | E | E-F [r] | x | x | 14d | 4 |
| Sulforix | M | x | x | x | x | 1-2 gal., 1 pt. | 1-2 gal., 1 pt. | 1 pt./100 g. | x | x | 48h | NA |
| calcium polysulfide | | x | x | x | x | G-E | i | F-G | x | x | 48h | 8 |
| Switch 62.5 WG | 9+12 | x | x | x | 11-14 oz. | x | x | x | x | x | 12h | 56 oz. |
| cyprodinil + fludioxonil | | x | x | x | G | x | x | x | x | x | 7d | NA |
| Tebustar 45WSP | 3 | x | x | 4 oz. | x | x | x | 4 oz. | x | x | 12h | 2 lb. |
| tebuconazole | | x | x | E | x | x | x | E [r] | x | x | 0d | NA |

(Continued)

Table 6-9. Grape Diseases - Veraison Through Harvest¹ (continued)

| Product And Formulation Active Ingredient | FRAC Code ² | Anthraco- se Foliar | Anthraco- se Fruit rot | Black Rot | Botrytis Bunch Rot | Downy Mildew | Phomopsis Blight | Powdery Mildew | Bitter Rot | Ripe Rot | REI ³ PHI ⁴ | Max Amt ⁵ Max App ⁶ |
|--|------------------------|---------------------------|------------------------------|-----------------|-----------------------|---------------|---------------------|--------------------------|------------------|----------|--------------------------------------|--|
| Topguard EQ | 3+11 | x | x | 5-6 fl. oz. | 8 fl. oz. | 8 fl. oz. | 8 fl. oz. | 5-6 fl. oz. | x | x | 12h | 34 fl. oz. |
| azoxystrobin+flutriafol | | x | x | u | s | u | u | E | x | x | 14d | 6 |
| Topsin M WSB | 1 | x | x | 0.75-1.5 lb. | x | x | 0.75-1.5 lb. | 0.75-1.5 lb. | 0.75- 1.5 lb. | x | 2d | 6 lb. |
| thiophanate-methyl | | x | x | F | x | x | G | E | G | x | 7d | NA |
| Torino (SC) | U6 | x | x | x | x | x | x | 3.4-6.8 oz. | x | x | 4h | 7 oz. |
| cyflufenamid | | x | x | x | x | x | x | E | x | x | 3d | 1 or 2 |
| Vanguard WG | 9 | x | x | x | 10 oz. | x | x | 10 oz. | x | x | 12h | 30 oz. |
| cyprodinil | | x | x | x | G-E | x | x | s | x | x | 7d | NA |
| Vivando 2.5F | U8 | x | x | x | x | x | x | 10.3- 15.4 fl. oz. | x | x | 12h | 43 fl. oz. |
| metrafenone | | x | x | x | x | x | x | E | x | x | 14d | 3 |
| Zampro | 45+40 | x | x | x | x | 11-14 fl. oz. | x | x | x | x | 12h | 56 fl. oz. |
| ametoctradin + dimethomorph | | x | x | x | x | E | x | x | x | x | 14d | NA |

Footnotes: See how to read the spray schedule tables section at the beginning of the chapter.

Grape Postharvest - Foliar Diseases

Downy mildew, powdery mildew: In some years, these diseases may cause defoliation well before the onset of cool weather in the fall. Postharvest early defoliation predisposes the vines to winter injury and reduces productivity for the following season. It is important to maintain at least some protection against foliar infections by these fungi. Postharvest rates for fungicides should be the same as pre-harvest rates. Check labels for season limits on quantity of products.

Special Comments on Grape Schedule

Disease Management

Anthracnose

Early-season applications are important to keep anthracnose from spreading to new tissues. As leaves and canes mature (fully expanded), they become resistant to infection; however, new leaves and succulent cane tips are susceptible throughout the season, and berries remain susceptible until veraison.

Foliar fungicides probably do not provide satisfactory anthracnose control unless you use them in conjunction with a delayed-dormant lime-sulfur or Sulforix application.

Grape bitter rot

Unlike black rot, which does not infect berries late in the season, bitter rot attacks only mature berries. Both diseases result in black, shriveled (mummified) fruit, and some growers have mistaken bitter rot for black rot. A rule of thumb is that if a rot develops on mature berries (8 percent or greater sugar), it more likely to be bitter rot than black rot.

If bitter rot is a problem, pre-harvest applications of captan may be beneficial. Observe all pre-harvest restrictions.

Grape ripe rot

Ripe rot symptoms appear on mature berries, resulting in shriveled berries with a vinegar odor or bitter taste. Infected berries become covered in blister-like lesions (acervuli) covered in salmon-colored spore masses.

Minnesota varieties Frontenac and Marquette are extremely susceptible to ripe rot.

Ripe rot is very difficult to manage. The systemic FRAC 11 fungicides (Abound, Sovran, Flint, and products that contain a FRAC 11 fungicide such as Pristine, Merivon, Luna Sensation and Quadris Top) or FRAC 12 fungicides (Miravis Prime, Switch) are moderately effective against ripe rot but must be applied during the critical period for berry infections. Captan may also be effective. Observe all pre-harvest restrictions.

Botrytis bunch rot

Botrytis bunch rot is most commonly a problem on tight-clustered French hybrid and *Vitis vinifera* cultivars. Infections can occur near bloom, but the disease does not appear until veraison or during harvest. Proper timing and thorough spray coverage are essential for good control.

Note: Growers in Europe and Canada have reported fungicide resistance due to overuse of Rovral over three to five years. Vanguard and Elevate are also at risk for fungicide resistance development. We therefore recommend limiting Rovral, Elevate, and Vanguard applications to two per year to reduce the probability of developing isolates of *Botrytis* resistant to these materials. In addition, consider alternating applications of Rovral, Elevate, and Vanguard during the growing season. Note: Removing leaves around clusters on mid- or low- wire cordon-trained vines before bunch closing has been shown to reduce *Botrytis*-caused losses.

Grape sour rot

Sour rot occurs late in the season near harvest (15 percent or greater sugar; 15 Brix) on berries damaged from bird pecks, rain cracking, and insects. The most obvious sour rot symptom is a pre-harvest decay accompanied by a strong vinegar smell; that is, acetic acid, or what winemakers call volatile acidity. The berries usually turn a tan color, soften, and eventually break down and disintegrate. The decayed berries seldom have any noticeable fungal growth or fruiting bodies on the surface like you would see with *Phomopsis*, *Botrytis* bunch rot, or black rot.

A combination of yeasts and bacteria in a step-wise process cause sour rot. Yeasts convert the fruit sugar to ethanol, and then the bacteria convert the ethanol to acetic acid. Both yeasts and bacteria require some type of physical injury or wound to infect the berries, so bird pecks, fruit fly injury (genus *Drosophila*) rain cracking, compression in tight clusters, and so on are all involved in the process. While filamentous fungi (such as *Botrytis*) may be associated with the rotting berries, they don't appear to be the cause. Thus, traditional fungicides do not control sour rot. For images

and more information on sour rot see <https://ohioline.osu.edu/factsheet/plpath-fru-50>.

The most effective sour rot control is to minimize the populations of yeasts, bacteria, and fruit flies. Starting at 15 Brix, apply an insecticide (Mustang Maxx, Delegate, and malathion) to control fruit flies and an antimicrobial (Oxidate, Fracture) to reduce yeasts and bacteria-associated with sour rot.

Fungicide resistance management

A spray program should be thoughtfully developed to prevent and slow the development of fungicide-resistant pathogens in the vineyard. Fungicides that have a site-specific mode of action are classified as medium to high risk for fungicide resistance development.

Fungicides with Fungicide Resistance Action Committee (FRAC) codes or numbers 1, 2, 3, 4, 7, 9, 11, and 49, are medium to high-risk fungicides. No more than one sequential applications of a high-risk fungicide (FRAC 11, 3, 7) should be applied before alternating to a fungicide with a different mode of action. High-risk fungicides have restrictions on how much and how often they can be applied. Do not overuse fungicides. It is unlawful to apply fungicides in a manner inconsistent with the product label.

Copper fungicides for grape disease control

When different formulations of copper are dissolved in water, copper ions are released into solution. These copper ions are toxic to fungi and bacteria because of their ability to denature proteins. However, using copper fungicides carries the risk of injuring foliage and fruit of most crops.

Factors promoting copper injury include:

1. The amount of actual copper applied.
2. Cold, wet weather (slow drying conditions) that apparently increases the availability of copper ions and, thus, increases the risk of plant injury.

Because of the potential to injure plants and to accumulate in soil, copper fungicides in conventional production systems have largely been replaced with other fungicides that are generally safer to plant tissues and often more effective.

Several terms are used when discussing copper as a fungicide. The original material used was called copper sulfate (also known as blue vitriol or bluestone). When this material was combined with lime in French vineyards, the combination became known as Bordeaux mixture.

Bordeaux mixture

Bordeaux mixture is a mixture of copper sulfate and hydrated lime in water. It has long residual action and has been used for years to control many diseases, including downy mildew and powdery mildew of grape. It can be mixed on-site but is also available as a dry wettable powder.

Fixed copper fungicides

Fixed copper formulations release copper ions more slowly and generally injure plant tissues less (safer to use) than Bordeaux mixture. But fixed copper use is still limited because of their potential to injure plants and lack of compatibility with other pesticides.

Some of the more common commercial formulations of fixed copper include:

Basic copper sulfate: Griffin Basicop, Basic Copper "53," Micro Flo Cuproxat, Tennessee Brand Tri-Basic Copper Sulfate, Tenn-Cop 5E, and Cuprofix Ultra 40DF.

Copper (Cupric) hydroxide: Agtrol Champion WP, Agtrol Champ flowable, Agtrol Champ 2F, Kocide 101, Kocide 3000DF, Kocide 2000D, Microflo BlueShield WP, and Microflo BlueShield DF.

Recommendations for copper fungicide use on grapes

Copper fungicides are highly effective against downy mildew and are moderately effective against powdery mildew. Copper fungicides are weak for controlling black rot, Botrytis bunch rot and Phomopsis blight.

To reduce the risk of phytotoxicity when using copper:

1. Do not make a complete season-long spray program with only copper fungicides.
2. Use fungicides other than copper whenever possible.
3. Delay copper use as late into the growing season as possible.
4. Avoid the use of copper sulfate alone. Always use a "fixed" copper formulation.
5. Remember that cool, wet weather enhances the risk of copper injury. Be especially certain to use adequate lime levels during such periods or switch to other fungicides.
6. Some products are incompatible with copper. Do not mix copper products with anything that will acidify the spray mixture (such as phosphorus acid fungicides).
7. Avoid copper and lime sprays on fruit destined for fresh market.

Late season applications of copper, captan, and sulfur

Broad-spectrum fungicides such as copper, captan, and sulfur should be avoided within 30-45 days of harvest. These products can interfere with the fermentation process by killing the yeast, resulting in the fermentation getting stuck. Stuck fermentations are hard to get going again and make the juices more susceptible to bacterial contamination or hydrogen sulfide production. Removing leaves around the clusters (leaf pulling) reduces the risk of fungal infections late in the season and thus reduces reliance on late season fungicides applications.

Note on insecticide resistance management

Insects have been known to develop resistance to insecticides after repeated exposure. For insecticide resistance management, avoid successive applications of insecticides in the same group or type of chemistry. The Insecticide Resistance Action Committee codes (IRAC codes) listed in each management section identify the various insecticide modes of action group. Rotating to insecticides with a different IRAC code should help avoid development of insecticide resistance.

Insect Management

Spotted-wing Drosophila

Spotted-wing Drosophila (SWD) is a serious invasive pest that attacks small fruit crops, some stone fruits (cherry, nectarine, peach), high tunnel tomatoes, and wild hosts (including pokeweed, autumn olive, crabapple, nightshade, Amur honeysuckle, and wild grape).

SWD is different than other fruit flies; the female has a stout, toothed ovipositor (egg layer) that enables her to lay eggs under the skin of ripening fruits that are otherwise healthy and sound. Soft-skinned fruit generally become vulnerable to attack as they begin to soften and turn color during ripening, usually in the final 7 to 10 days before harvest. The larvae tunnel and feed under the skin of the fruit and can reach 4 millimeters long. There is often a sunken area at the site where the eggs are laid, and damaged fruit may appear to collapse from the internal damage and rots.

SWD is able to complete its life cycle in just more than a week when temperatures are optimal, and there may be 10 or more generations per year. Growers need to monitor plantings for SWD in the final weeks before harvest. Traps for monitoring and detecting SWD are available. More information about SWD is available from Michigan State University Integrated Pest Management: <https://www.canr.msu.edu/swd/>.

Look for additional state labels that may allow for changes to rates and allowable number of applications

of various insecticides. When applying insecticides during the harvest period, carefully watch the pre-harvest intervals for the products you choose to apply.

Multi-colored Asian lady beetle

The multi-colored Asian lady beetle (MALB), a late-season vineyard inhabitant, can significantly reduce wine quality. These beetles are attracted to ripening grapes as a source of sugars in late summer and fall. They may congregate, often by the hundreds or thousands, in and among grape clusters from August through October.

Although they may cause direct yield loss, they more often reduce wine quality when sufficient numbers become trapped in the harvested grapes and are crushed along with them at the winery. When stressed, MALB secretes a defensive chemical that causes wine to smell “dirty,” (a musty, damp odor), masking the flavors and smells of the grapes.

As few as two MALB per lug of grapes can alter wine flavor and bouquet enough to be detected. Excessive numbers of MALB in grape clusters are most common in late-ripening varieties such as Cabernet Franc, Cabernet Sauvignon, Chambourcin, Riesling, Vidal, and Vignoles, but earlier grapes that are prone to cracking can also be infested.

Scout vineyards several days before harvest to determine the abundance of MALB. Belay 2.13SC, Venom 70SG, and Scorpion 35S are labeled specifically for control of this insect in grapes. Additional insecticides (including Baythroid and Mustang Maxx) have short pre-harvest intervals and, although not labeled specifically against MALB, have been effective in trials and vineyard use.

Grape root borer

This insect can be a serious pest in southern parts of our region. There are no insecticides currently labeled for use against this pest in grapes, but it can be managed with the biocontrol tactic of insect-parasitic nematodes. Research on the behavioral tactic of pheromone mating disruption has looked promising, but no mating disruption product is currently registered for control of this pest on grapes.

Spotted lanternfly

The spotted lanternfly is an invasive planthopper that is currently spreading throughout the Midwest. This insect feeds on plant sap causing wilting, dieback, and even death.

Currently, spotted lanternfly is believed to pose the greatest threat to the grape, hops, and hardwood industries. Know how to identify this pest and remain vigilant for its appearance in your vineyard and orchard systems. Consult your state entomologist and state department of Agriculture for guidance on identifying this pest, its preferred invasive host tree (tree of heaven), and potential use of monitoring traps.

Wasps in fruit plantings

Almost anywhere fruit is produced, wasps can become a nuisance or, in some cases, a severe pest to field workers. Unfortunately, little help is available for controlling wasps.

Wasps are generally attracted to the juice and soft fruit. Sanitation is key to preventing or at least reducing problems with wasps. Pick all ripe fruit and fruit debris regularly and thoroughly. Also remove any item that has food value (e.g., soft drinks, lunches, etc.) that pickers may bring in.

Table 6-10. Relative Disease Susceptibility and Chemical Sensitivity Among Grape Cultivars

The relative ratings in this chart apply to an average growing season under conditions usually favorable for disease development. Any given cultivar may be more or less severely affected depending on conditions.

| Cultivar | Susceptible Or Sensitive To ¹ | | | | | | | | | | | |
|-----------|--|--------------|----------------|--------------------|-----------|--------|------------|-------------|---------------------|---------------------|--------------------|----------------------|
| | Black Rot | Downy Mildew | Powdery Mildew | Botrytis Bunch Rot | Phomopsis | Eutypa | Crown Gall | Anthraxnose | Sulfur ² | Copper ³ | 2,4-D ⁴ | Dicamba ⁴ |
| Arandell | + | + | + | + | ++ | ? | ? | + | ? | ? | ++ | ? |
| Aromella | + | +++ | + | + | ++ | ? | ? | + | ? | ? | +++ | +++ |
| Aurore | +++ | ++ | ++ | +++ | + | +++ | ++ | + | No | ++ | ? | ? |
| Baco Noir | +++ | + | ++ | ++ | + | ++ | +++ | + | No | ? | ? | ? |
| Brianna | ? | + | ? | + | ? | ? | ? | ? | Yes | +++ | ++ | + |

(Continued)

Table 6-10. Relative Disease Susceptibility And Chemical Sensitivity Among Grape Cultivars (continued)

| Cultivar | Susceptible Or Sensitive To ¹ | | | | | | | | | | | |
|--------------------|--|------------------|----------------|--------------------|-----------|--------|------------|-------------|---------------------|---------------------|--------------------|----------------------|
| | Black Rot | Downy Mildew | Powdery Mildew | Botrytis Bunch Rot | Phomopsis | Eutypa | Crown Gall | Anthraxnose | Sulfur ² | Copper ³ | 2,4-D ⁴ | Dicamba ⁴ |
| Cabernet Franc | +++ | +++ | +++ | + | ? | ? | +++ | ++ | No | ? | + | +++ |
| Cabernet Sauvignon | +++ | +++ | +++ | + | +++ | +++ | +++ | ? | No | + | + | ? |
| Catawba | +++ | +++ | ++ | + | +++ | + | + | ++ | No | ++ | ++ | ++ |
| Cayuga White | + | ++ | + | + | ++ | + | ++ | +++ | No | + | + | +++ |
| Chambourcin | +++ | + | +++ | ++ | + | ? | ++ | + | Yes | ? | +++ | ++ |
| Chancellor | + | +++ | +++ | + | +++ | + | +++ | ++ | Yes | +++ | ++ | ? |
| Chardonel | ++ | ++ | ++ | ++ | +++ | ++ | ++ | + | No | ? | ++ | +++ |
| Chardonnay | ++ | +++ | +++ | +++ | +++ | ++ | +++ | +++ | No | + | ++ | +++ |
| Concord | +++ | + | ++ | + | +++ | +++ | + | + | Yes | + | +++ | ++ |
| Corot noir | + | +++ | + | + | ++ | + | + | + | No | ? | ++ | +++ |
| Cynthiana/Norton | + | ++ | + | + | + | ? | + | + | Yes | ? | +++ | +++ |
| DeChaunac | + | ++ | ++ | + | +++ | +++ | ++ | ++ | Yes | + | + | ++ |
| Delaware | ++ | +++ ⁵ | ++ | + | +++ | + | + | ++ | No | + | +++ | ? |
| Edelweiss | ? | ? | ? | ? | ? | ? | ? | ? | ? | ? | ++ | ? |
| Faith | + | + | + | + | + | + | + | + | ? | ? | + | ? |
| Foch | ++ | + | ++ | + | + | +++ | + | ++ | Yes | ++ | +++ | +++ |
| Fredonia | ++ | +++ | ++ | + | +++ | ? | + | +++ | No | ? | ++ | ++ |
| Frontenac | +++ | + | ++ | ++ | + | ? | ? | +++ | No | ++ | + | +++ |
| Frontenac Gris | ++ | + | ++ | ++ | + | ? | ? | ++ | No | ++ | + | + |
| Geneva Red | + | ++ | ++ | ++ | + | + | + | + | No | ? | + | +++ |
| Gewürztraminer | +++ | +++ | +++ | +++ | ? | ? | +++ | +++ | No | + | ? | ? |
| Gratitude | + | + | + | + | + | + | +++ | + | ? | ? | + | ? |
| Hope | + | + | + | + | + | + | + | + | ? | ? | + | ? |
| Joy | + | + | + | + | + | + | + | + | ? | ? | + | ? |
| Jupiter | ++ | +++ | +++ | + | + | ? | ? | + | ? | ? | + | ++ |
| LaCrescent | ++ | +++ | ++ | + | +++ | + | + | + | No | ? | +++ | +++ |
| LaCrosse | +++ | ++ | ++ | +++ | ++ | ? | ? | + | No | ++ | +++ | +++ |
| Lemberger | +++ | +++ | +++ | + | ? | +++ | +++ | ? | No | ? | ++ | ? |
| Leon Millot | + | ++ | +++ | + | + | + | ? | + | Yes | ++ | + | ? |
| Marquette | ++ | + | + | +++ | +++ | ? | + | +++ | No | ++ | +++ | + |
| Marquis | + | +++ | + | + | +++ | ? | ? | +++ | ? | ? | + | ? |
| Mars | + | + | + | + | + | ? | + | ++ | ? | ? | + | + |
| Merlot | ++ | +++ | +++ | ++ | + | +++ | +++ | ++ | No | ++ | ? | ? |
| Moore's Diamond | +++ | + | +++ | ++ | ? | ++ | ? | ? | No | ? | ? | ? |
| Niagara | +++ | +++ | ++ | + | +++ | + | ++ | ++ | No | + | +++ | ++ |
| Noiret | +++ | ++ | ++ | + | + | ? | ++ | + | No | ? | ++ | +++ |
| Petite Pearl | +++ | + | + | + | + | ? | + | + | ? | ? | + | ? |
| Pinot gris | +++ | +++ | +++ | ++ | ? | +++ | +++ | ? | No | ? | ? | ? |
| Pinot noir | +++ | +++ | +++ | +++ | ? | ? | +++ | ? | No | + | ? | ? |
| Reliance | +++ | +++ | ++ | + | ++ | ? | ? | +++ | No | + | + | ? |
| Riesling | +++ | +++ | +++ | +++ | ++ | ++ | +++ | ? | No | + | + | ++ |

(Continued)

Table 6-10. Relative Disease Susceptibility And Chemical Sensitivity Among Grape Cultivars (continued)

| Cultivar | Susceptible Or Sensitive To ¹ | | | | | | | | | | | |
|---------------|--|--------------|----------------|--------------------|-----------|--------|------------|-------------|---------------------|---------------------|--------------------|----------------------|
| | Black Rot | Downy Mildew | Powdery Mildew | Botrytis Bunch Rot | Phomopsis | Eutypa | Crown Gall | Anthraxnose | Sulfur ² | Copper ³ | 2,4-D ⁴ | Dicamba ⁴ |
| St. Croix | ? | ++ | ++ | ++ | +++ | ? | ? | + | No | ++ | ++ | ? |
| Seyval | ++ | ++ | +++ | +++ | ++ | + | ++ | + | No | + | ++ | +++ |
| Steuben | ++ | + | + | + | + | ? | + | +++ | No | ? | + | ++ |
| Sunbelt | + | ++ | ++ | + | + | ? | ? | + | ? | ? | +++ | ++ |
| Thompcord | + | +++ | + | + | + | ? | + | + | ? | ? | + | + |
| Traminette | + | ++ | + | + | +++ | ? | ++ | + | No | ? | ++ | ++ |
| Valvin Muscat | ++ | + | ++ | + | + | ? | + | ? | No | ? | +++ | + |
| Vanessa | +++ | ++ | ++ | + | + | ? | + | ? | ? | ? | + | ? |
| Vidal blanc | + | ++ | +++ | + | + | + | ++ | +++ | No | ? | ++ | +++ |
| Vignoles | + | ++ | +++ | +++ | ++ | ++ | ++ | +++ | No | ? | + | +++ |

¹ + = slightly susceptible or sensitive. ++ = moderately susceptible or sensitive. +++ = highly susceptible or sensitive. No = not sensitive. Yes = sensitive. ? = relative susceptibility or sensitivity not established.

² Slight to moderate sulfur injury may occur even on tolerant cultivars when temperatures are 85°F or higher during, or immediately following, the application.

³ Copper applied under cool, slow-drying conditions is likely to cause injury.

⁴ Herbicide sensitivity ratings based on observation and simulated drift studies in Indiana.

⁵ Berries not susceptible.

7. BLUEBERRY

Blueberry Spray Schedule

The shaded/colored boxes represent the crop stages where common pests in the Midwest are active and action (scouting or preventative sprays) may be needed or recommended.

| Stage | | | | | | |
|---------|---|----------|--------------------------|---------------|--------------|-----------------------------|
| Dormant | Green Tip | Pink Bud | Petal Fall | First Cover | Second Cover | Third And Additional Covers |
| | | | Cherry Fruitworm | | | |
| | | | Cranberry Fruitworm | | | |
| | | | | Plum Curculio | | |
| | | | | | | Blueberry Maggot |
| | | | | | | Brown Marmorated Stink Bug |
| | | | | | | Japanese Beetle |
| | | | | | | Spotted-Wing Drosophila |
| | | | Blueberry Stem Gall Wasp | | | |
| Major | Present in most years and usually causing economic damage if not managed. | | | | | |
| Minor | Often present but not causing economic damage and not requiring management. | | | | | |

How to read the spray schedule tables

Every blueberry growth stage has important notes on disease or insect management. In some cases, the reader will be directed to the special problems section at the end of the section or chapter. Please make sure to read thoroughly and contact your state Extension specialist with any specific questions.

Key to tables

E = excellent control

G = good control

F = fair control

[r] = fungicide/insecticide resistance possible

s = suppression only

i = ineffective

u = unknown efficacy

x = pest not on the label

¹ Efficacy data in this publication are based on trials conducted across various regions and does not necessarily reflect local efficacy differences or changes over time. Growers should contact their Extension specialist for the most recent or for state-specific information. The information on this publication is only a guide; the authors and their institutions assume no liability for

practices implemented based on this information. Always read and follow pesticide labels. The label is the law. Product registration may vary by state.

² F/IRAC code represents the mode of action of the fungicide/insecticide.

³ PHI refers to the pre-harvest interval, which is the number of days before harvest that the product may not be applied.

⁴ All fungicides/insecticides have a Restricted-Entry Interval (REI). The restricted-entry interval is the time immediately after a pesticide application when entry into the treated area is limited. Check labels for REI. Restrictions in REI may prohibit the use of certain pesticides during harvest.

Applicators must abide by both maximum amount of product per season AND maximum number of applications.

⁵ Max amt refers to the product's maximum amount/acre/year.

⁶ Max app refers to the product's maximum number of applications per year.

RUP refers to restricted use pesticide.

Blueberry Dormant to Delayed Dormant - Diseases

Apply after buds begin to break.

- Orondis Gold and Ridomil Gold SL are labeled for control of Phytophthora root rot of high bush blueberries; Ridomil Gold is labeled for highbush and low bush blueberry types. Apply to established plantings before the plants start growth in the spring. Apply to new plantings at time of planting.
- Sulforix should not be applied within 14 days of an oil spray.
 - Orondis Gold contains mefenoxam (FRAC 4), the active ingredient of Ridomil Gold (FRAC 4). Use only one of these products for control of Phytophthora and alternate with a FRAC 33.
- Ziram 76DF should be applied at loose bud and again 7 days later for Phomopsis cane blight. Ziram is currently in limited supply, and a shortage is anticipated in 2026 due to its upcoming EPA registration cancellation.

Table 7-1. Blueberry Diseases - Dormant Through Delayed Dormant¹

| Product And Formulation | Active Ingredient | FRAC Code ² | Phomopsis Cane Blight | Phytophthora Root Rot | REI ³ PHI ⁴ | Max Amt ⁵ Max App ⁶ |
|-------------------------|----------------------------|------------------------|-----------------------|-----------------------|-----------------------------------|---|
| Aliette WDG | | P07 (33) | 5 lb. | 0.5 lb. | 24h | 20 lb. |
| | aluminum tris | | s | G | 12h | 4 |
| Orondis Gold | | 49+4 | x | 28-55 oz. | 48 h | 110 fl. oz. |
| | oxathiapipralin+ mefenoxam | | x | E | 1 d | 2 |
| ProPhyt | | 33 | x | 4 pt. | 4h | NA |
| | potassium phosphite | | x | G | 0d | NA |
| Ridomil Gold SL | | 4 | x | 3.6 pt. | 48h | 7.2 pt. |
| | mefenoxam | | x | E | 0d | 2 |
| Sulforix | | M | 1-2 gal./100 gal. | x | 48h | 8 gal. |
| | calcium polysulfide | | u | x | NA | 4 |
| Ziram 76DF | | M3 | 3 lb. | x | 48h | NA |
| | ziram | | G | x | 30d | NA |

Footnotes: See how to read the spray schedule tables section at the beginning of the chapter.

Blueberry Green Tip - Diseases

- Apply when leaf buds swell and are showing 1/16-1/4 inch green tip.
- For control of **mummy berry**: Scout for fallen mummies producing tiny, trumpet-like “mushrooms,” particularly in wet areas. “Mushrooms” produce spores for 1-4 weeks, with the longer durations occurring under cooler conditions.
- Freezing temperatures may result in injury and increase the susceptibility of young leaves and shoots to Botryosphaeria infection. A spray within 24 hours of a frost event may improve efficacy of fungicides used.
- Tank mixes that include captan or Bravo with oils or EC formulated pesticides can cause phytotoxicity.
- The second application of Ridomil may be applied during periods of high disease pressure and wet conditions.

Table 7-2. Blueberry Diseases - Green Tip¹

| Product And Formulation Active Ingredient | FRAC Code ² | Botryosphaeria Canker | Mummy Berry | Phomopsis Cane Blight | REI ³ PHI ⁴ | Max Amt ⁵ Max App ⁶ |
|--|------------------------|--------------------------|------------------|--------------------------|--------------------------------------|--|
| Abound (SC) | 11 | 6-15.5 fl. oz. | 6-15.5 fl. oz. | 6-15.5 fl. oz. | 4h | 42 fl. oz. |
| azoxystrobin | | u | F | F | 0d | NA |
| Aliette WDG | P07 (33) | x | 5 lb. | 5 lb. | 12h | 20 lb. |
| aluminum tris | | x | s | G | 12h | 4 |
| Bravo Weather Stik | M3 | x | 3-4 pt. | 3-4 pt. | 12h | 12 pt. |
| chlorothalonil | | x | E | s | 42d | NA |
| Captan 80WG | M 5 | x | 1.25-3 lb. | 1.25-3 lb. | 48h | 43.75 lb. |
| captan | | x | s[E] | F | 0d | NA |
| Fontelis | 7 | x | 24 fl. oz. | 16-24 fl. oz. | 12h | 72 fl. oz. |
| penthiopyrad | | x | E | x | 0d | NA |
| Indar 2F | 3 | x | 6 fl. oz. | 6 fl. oz. | 12h | 24 fl. oz. |
| fenbuconazole | | x | E | G | 30d | 4 |
| Inspire Super | 3+9 | x | 16-20 fl. oz. | x | 12h | 80 fl. oz. |
| difenoconazole + cyprodinil | | x | E | x | 7d | NA |
| Kocide 3000 | M | x | 1.75 lb.-3.5 lb. | 1-2.25 lb. | 48h | 28 lb. |
| copper hydroxide | | x | F | s | 0d | NA |
| Luna Flex | 3+7 | x | 1.2-13.6 fl. oz. | 11.2-13.6 fl. oz. | 12h | 27.2 fl. oz. |
| fluopyram + difencolazole | | x | G | G | 7d | 2 |
| Luna Tranquility (SC) | 7+9 | x | 13.6-27 fl. oz. | 27 fl. oz. | 12h | 54.7 fl. oz. |
| fluopyram + pyrimethanil | | x | E-G | s | 0d | NA |
| Miravis Prime | 7+12 | x | 9-13.4 fl. oz. | 9-13.4 fl. oz. | 12h | 26.8 fl. oz. |
| pydiflumetofen+fludioxonil | | x | G | G | 0d | NA |
| Omega 500F | 29 | x | 20 fl. oz. | 20 fl. oz. | 12h | 120 fl. oz. |
| fluazinam | | x | F | u | 30d | NA |
| Pristine (38WG) | 11+7 | x | 18.5-23 oz. | 18.5-23 oz. | 24h | 92 oz. |
| pyraclostrobin + boscalid | | x | F | G | 0d | 4 |
| ProPhyt | P07 (33) | x | x | x | 4h | NA |
| potassium phosphite | | x | x | x | 0d | NA |
| Quadris Top | 11+3 | x | 12-14 fl. oz. | x | 12h | 56 fl. oz. |
| azoxystrobin + difenoconazole | | x | G | x | 7d | 4 |
| Quash SC | 3 | 2.5 oz. | 2.5 oz. | 2.5 oz. | 12h | 7.5 oz. |
| metconazole | | u | E | E | 7d | 3 |
| Quilt Xcel | 11+3 | 14-21 fl. oz. | 14-21 fl. oz. | 14-21 fl. oz. | 12h | 63 fl. oz. |
| azoxystrobin + propiconazole | | u | F | G | 30d | NA |
| Sulforix | M | 1-2 gal./100 gal. | x | 1-2 gal./100 gal. | 48h | 8 gal. |
| calcium polysulfide | | u | x | G | NA | 4 |
| Switch 62.5WG | 9+12 | x | 11-14 oz. | 11-14 oz. | 12h | 56 oz. |
| cyprodinil + fludioxonil | | x | F | F | 0d | NA |

(Continued)

Table 7-2. Blueberry Diseases - Green Tip¹ (continued)

| Product And Formulation Active Ingredient | FRAC Code ² | Botryosphaeria Canker | Mummy Berry | Phomopsis Cane Blight | REI ³ PHI ⁴ | Max Amt ⁵ Max App ⁶ |
|--|------------------------|--------------------------|----------------|--------------------------|--------------------------------------|--|
| Tilt | 3 | x | 6 fl. oz. | x | 12h | 30 fl. oz. |
| propiconazole | | x | G | x | 30d | 5 |
| Ziram 76DF | M3 | x | 3 lb. | 3 lb. | 48h | NA |
| ziram | | x | G | G | 30d | NA |

Footnotes: See how to read the spray schedule tables section at the beginning of the chapter.

Blueberry Pink Bud to Petal Fall - Diseases

Make applications as needed when flower petals show pink, at 25% bloom, and every 7-10 days when blooms are open through petal fall.

- **Mummy berry** management: Pre-bloom fungicide sprays should begin at green tip and continue sprays until all blooms have fallen.

- For early harvest berries: Be aware of the preharvest intervals (PHI) of Bravo, Indar2F, Quilt. Tilt, Ziram and generic counterparts.
- In the southeastern region, Botrytis and anthracnose populations have developed resistance to the fungicides Switch, Elevate, and Pristine.

Table 7-3. Blueberry Diseases - Pink Bud Through Petal Fall¹

| Product And Formulation Active Ingredient | FRAC code ² | Alternaria Fruit Rot | Anthraco- nose Fruit Rot | Botryos- phaeria Canker | Botrytis Blight | Mummy Berry | Phomopsis Cane Blight | REI ³ PHI ⁴ | Max Amt ⁵ Max App ⁶ |
|--|------------------------|-------------------------|--------------------------------|-------------------------------|--------------------|----------------|--------------------------|--------------------------------------|--|
| Abound (SC) | 11 | 6-15.5 fl. oz. | 6-15.5 fl. oz. | 6-15.5 fl. oz. | 6-15.5 fl. oz. | 6-15.5 fl. oz. | 6-15.5 fl. oz. | 4h | 46 fl. oz. |
| azoxystrobin | | E | E | u | s (P) | F | F | 0d | see label |
| Bravo Weather Stik | M3 | 3-4 pt. | 3-4 pt. | x | 3-4 pt. | 3-4 pt. | 3-4 pt. | 12h | 12 pt. |
| chlorothalonil | | u | s[G] | x | s | E | s | 42d | NA |
| Captan 80WG | M 5 | 1.25-3 lb. | 1.25-3 lb. | x | 1.25-3 lb. | 1.25-3 lb. | 1.25-3 lb. | 48h | 43.75 lb. |
| captan | | G | G | x | F | s[E] | F | 0d | NA |
| Elevate 50WDG | 17 | x | x | x | 1.5 lb. | 1.5 lb. | x | 12h | 6 lb. |
| fenhexamid | | x | x | x | E | F | x | 0d | NA |
| Fontelis | 7 | x | x | x | 16-24 fl. oz. | 24 fl. oz. | 16-24 fl. oz. | 12h | 72 fl. oz. |
| penthiopyrad | | x | x | x | E | E | E | 0d | NA |
| Indar 2F | 3 | 6 fl. oz. | 6 fl. oz. | x | 6 fl. oz. | 6 fl. oz. | 6 fl. oz. | 12h | 24 fl. oz. |
| fenbuconazole | | F | u | x | E | E | s | 30d | 4 |
| Inspire Super | 3+9 | 16-20 fl. oz. | 16-20 fl. oz. | x | 16-20 fl. oz. | 16-20 fl. oz. | x | 12h | 80 fl. oz. |
| difenoconazole + cyprodinil | | u | u | x | G-F | x | x | 7d | NA |
| Kenja 400SC | 7 | x | 13.5-15.5 fl. oz. | x | 13.5-15.5 fl. oz. | x | x | 12h | 54 fl. oz. |
| Isofetamid | | x | u | x | E | x | x | 7d | NA |
| Kocide 3000 | M | 1-2.25 lb. | 1-2.25 lb. | x | 1-2.25 lb. | x | x | 48h | 28 lb. |
| copper hydroxide | | F-P | F | x | F-P | x | x | 0d | NA |

(Continued)

Table 7-3. Blueberry Diseases - Pink Bud Through Petal Fall¹ (continued)

| Product And Formulation Active Ingredient | FRAC code ² | Alternaria Fruit Rot | Anthraco- nose Fruit Rot | Botryos- phaeria Canker | Botrytis Blight | Mummy Berry | Phomopsis Cane Blight | REI ³ PHI ⁴ | Max Amt ⁵ Max App ⁶ |
|--|---------------------------|-------------------------|--------------------------------|-------------------------------|----------------------|-------------------------|--------------------------|--------------------------------------|--|
| Luna Flex | 3+7 | 1.2-13.6 fl. oz. | 1.2-13.6 fl. oz. | x | 11.2-13.6 fl. oz. | 11.2-13.6 fl. oz. | 11.2-13.6 fl. oz. | 12h | 27.2 fl. oz. |
| fluopyram + difencolazole | | E | u | x | u | G | G | 7d | 2 |
| Luna Tranquility (SC) | 7+9 | 13.6-27 fl. oz. | 13.6-27 fl. oz. | x | 13.6-27 fl. oz. | 13.6-27 fl. oz. | 13.6-27 fl. oz. | 12h | 54.7 fl. oz. |
| fluopyram + pyrimethanil | | G-F | s | x | E | E-G | u | 0d | NA |
| Miravis Prime | 7+12 | 9-13.4 fl. oz. | 13.4 fl. oz. | x | 9-13.4 fl. oz. | 9-13.4 fl. oz. | 9-13.4 fl. oz. | 12h | 26.8 fl. oz. |
| pydiflumetofen+fludioxonil | | E | E | x | G | G | G | 0d | NA |
| Omega 500F | 29 | 20 fl. oz. | 20 fl. oz. | x | 20 fl. oz. | 20 fl. oz. | 20 fl. oz. | 12h | 120 fl. oz. |
| fluazinam | | E | E | x | u | F | u | 30d | NA |
| PhD/OSO | 19 | 6.2 oz. | 6.2 oz. | x | 6.2 oz. | 6.2 oz. | 6.2 oz. | 4h | NA |
| polyoxin D | | P | P | x | G | G | G-F | 0d | 6 |
| Pristine (38WG) | 11+7 | 18.5-23 oz. | 18.5-23 oz. | x | 18.5-23 oz. | 18.5-23 oz. | 18.5-23 oz. | 24h | 92 oz. |
| pyraclostrobin + boscalid | | G | E | x | E | F | G | 0d | 4 |
| Proline 480C | 3 | 5.7 fl. oz. | 5.7 fl. oz. | x | 5.7 fl. oz. | 5.7 fl. oz. | 5.7 fl. oz. | 12h | 11.4 oz. |
| prothioconazole | | F | u | x | E | E | E-G | 7d | 2 |
| ProPhyt | P07 (33) | 4 pt | 4 pt. | x | x | x | x | 4h | NA |
| potassium phosphite | | u | F | x | x | x | x | 0d | NA |
| Quadris Top | 11+3 | 12-14 fl. oz. | 12-14 fl. oz. | x | x | 12-14 fl. oz. | x | 12h | 56 fl. oz. |
| azoxystrobin + difenoconazole | | G | F | x | x | G | x | 7d | 4 |
| Quash SC | 3 | 2.5 oz. | 2.5 oz. | 2.5 oz. | 2.5 oz. | 2.5 oz. | 2.5 oz. | 12h | 7.5 oz. |
| metconazole | | E | E | u | F | E | E | 7d | 3 |
| Quilt Xcel | 11+3 | 14-21 fl. oz. | 14-21 fl. oz. | 14-21 fl. oz. | x | 14-21 fl. oz. | 14-21 fl. oz. | 12h | 63 fl. oz. |
| azoxystrobin + propiconazole | | E | E | u | x | F | G | 30d | NA |
| Sulforix | M | x | x | x | x | 1-2 gal./100 gal. | 1-2 gal./100 gal. | 48h | 8 gal. |
| calcium polysulfide | | x | x | x | x | G-F | u | NA | 4 |
| Switch 62.5WG | 9+12 | 11-14 oz. | 11-14 oz. | x | 11-14 oz. | 11-14 oz. | 11-14 oz. | 12h | 56 oz. |
| cyprodinil + fludioxonil | | E | G | x | E | F | F | 0d | NA |
| Tilt | 3 | x | x | x | x | 6 fl. oz. | x | 12h | 30 fl. oz. |
| propiconazole | | x | x | x | x | G | x | 30d | 5 |
| Ziram 76DF | M3 | 3 lb. | 3 lb. | x | 3 lb. | 3 lb. | 3 lb. | 48h | NA |
| ziram | | F | F | x | F | G | G | 30d | NA |

Footnotes: See how to read the spray schedule tables section at the beginning of the chapter.

Blueberry Pink Bud to Petal Fall - Insects

- Cherry fruitworm control by conventional insecticides (other than Esteem and Intrepid) starts at petal fall and are re-applied 10 days later.
- Cherry fruitworm control with Esteem: Apply when egg laying begins and again at petal fall.
- Cherry fruitworm control with Intrepid: First application is best at 400 degree days (base 50) after biofix (sustained catch of moths in pheromone trap). Second application at 100% petal fall.

Table 7-4. Blueberry Insects - Pink Bud Through Petal Fall¹

| Product And Formulation | Active Ingredient | IRAC Code ² | Cherry Fruitworm And Cranberry Fruitworm | REI ³ PHI ⁴ | Max Amt ⁵ Max App ⁶ |
|---|--|------------------------|--|-----------------------------------|---|
| Apta (1.34SC) | | 21A | 27 fl. oz. | 12h | 81 fl. oz. |
| | tolfenpyrad | | G | 3d | 3 |
| Asana XL (0.66EC) (RUP) | | 3A | 4.8-9.6 fl. oz. | 12h | 38.4 fl. oz. |
| | esfenvalerate | | G | 14d | NA |
| Assail 30SG | | 4A | 3.8-4.5 fl. oz. | 12h | 22.5 fl. oz. |
| | acetamiprid | | G | 1d | 5 |
| Avaunt eVo (30WDG) | | 22 | 3.5-6 oz. | 12h | 24 oz. |
| | indoxacarb | | G | 7d | 4 |
| <i>B.t. kurstaki</i> (DiPel DF, Javelin WG, etc.) | | 11 | 0.5-2 lb. | 4h | NA |
| | <i>Bacillus thuringiensis kurstaki</i> | | G | 0d | NA |
| Brigade 2EC (RUP) | | 3A | 2.1-6.4 fl. oz. | 12h | 32 fl. oz. |
| | bifenthrin | | G | 1d | 5 |
| Brigade WSB (10WP) (RUP) | | 3A | 5.3-16 oz. | 12h | 80 oz. |
| | bifenthrin | | G | 1d | 5 |
| Danitol 2.4EC (RUP) | | 3A | 10.6-16 oz. | 24h | 32 fl. oz. |
| | fenpropathrin | | E | 3d | 2 |
| Delegate WG (25WG) | | 5 | 3-6 oz. | 4h | 19.5 oz. |
| | spinetoram | | G | 3d | 6 |
| Diazinon AG600 WBC (RUP) | | 1B | 12.75 fl. oz. | 120h | 25.5 fl. oz. |
| | diazinon | | u | 7d | 2 |
| Entrust SC (2SC) | | 5 | 4-6 fl. oz. | 4h | 29 fl. oz. |
| | spinosad | | F | 1d | 6 |
| Esteem 35WP | | 7C | 5 oz. | 12h | 10 oz. |
| | pyriproxifen | | E | 7d | 2 |
| Exirel (0.83SE) | | 28 | 10-13.5 fl. oz. | 12h | 61.5 fl. oz. |
| | cyantraniliprole | | G | 3d | NA |
| Grandevo WDG | | UN | 1-3 lb. | 4h | NA |
| | <i>Chromobacterium subtsugae</i> | | E | 0d | NA |
| Imidan 70W | | 1B | 1.33 lb. | 24-72h | 713 lb. |
| | phosmet | | E | 3d | 5 |
| Intrepid 2F | | 18 | 10-16 fl. oz. | 4h | 48 fl. oz. |
| | methoxyfenozide | | F | 7d | 3 |
| Knack (0.86EC) | | 7C | 16 fl. oz. | 12h | 32 fl. oz. |

(Continued)

Table 7-4. Blueberry Insects - Pink Bud Through Petal Fall¹ (continued)

| Product And Formulation Active Ingredient | IRAC Code ² | Cherry Fruitworm And Cranberry Fruitworm | REI ³ PHI ⁴ | Max Amt ⁵ Max App ⁶ |
|--|------------------------|---|--------------------------------------|--|
| pyriproxyfen | | E | 7d | 2 |
| Lannate LV (2.4WSL) (RUP) | 1A | 1.5-3 pt. | 48h | 12 pt. |
| methomyl | | u | 3d | 4 |
| Malathion 8F | 1B | 1.25 pt. | 12h | NA |
| malathion | | u | 1d | 3 |
| Neemix 4.5 (0.39L), AzaDirect | UN | 4-16 fl. oz. | 4h | NA |
| azadirachtin | | u | 0d | NA |
| Pyganic 5EC | 3A | 4.5-15.6 fl. oz. | 12h | NA |
| pyrethrins | | u | 0d | 10 |
| Rimon 0.83EC | 15 | 20-30 fl. oz. | 12h | 90 fl. oz. |
| novaluron | | G | 8d | NA |
| Sevin XLR Plus (4F) | 1A | 1.5-2 qt. | 12h | 10 qt. |
| carbaryl | | G | 7d | 5 |
| Verdepryn 100SL (0.83SL) | 28 | 8.2-11 fl. oz. | 4h | 33 fl. oz. |
| cyclaniliprole | | G | 1d | 3 |

Footnotes: See how to read the spray schedule tables section at the beginning of the chapter.

Blueberry Diseases - Summer Cover

Apply first cover about 7-10 days after petal fall, and second cover about 10 days later.

Disease management notes

- Pre- and post-harvest rots can be greatly reduced by timely harvests and proper handling, followed by rapid, post-harvest cooling. Fungicides alone are not sufficient to control these diseases. Do not harvest or handle wet fruit.

- Do not use an adjuvant with Fontelis after petal fall.
- Applications of Abound, Quash and Quilt for control of other diseases may aid in the control Botryosphaeria (bot) canker.

Table 7-5. Blueberry Diseases - Summer Cover¹

| Product And Formulation Active Ingredient | FRAC Code ² | Alternaria Fruit Rot | Anthracnose Fruit Rot | Phomopsis Cane Blight | REI ³ PHI ⁴ | Max Amt ⁵ Max App ⁶ |
|--|------------------------|-------------------------|--------------------------|--------------------------|--------------------------------------|--|
| Abound (SC) | 11 | 6-15.5 fl. oz. | 6-15.5 fl. oz. | 6-15.5 fl. oz. | 4h | 46 fl. oz. |
| azoxystrobin | | G | G-E | F | 0d | varies |
| Aliette WDG | P07 (33) | 5 lb. | 5 lb. | 5 lb. | 12h | 20 lb. |
| aluminum tris | | u | u | 3-4 pt. | 0.5d | 4 |
| Bravo Weather Stik | M3 | 3-4 pt. | 3-4 pt. | 3-4 pt. | 12h | 12 pt. |
| chlorothalonil | | u | s | s | 42d | NA |
| Captan 80WG | M 5 | 1.25-3 lb. | 1.25-3 lb. | 1.25-3 lb. | 48h | 43.75 lb. |
| captan | | F | G | F | 0d | NA |
| Cevya | 3 | 4-5 fl. oz. | 4-5 fl. oz. | 4-5 fl. oz. | 12h | 15 fl. oz. |
| mefentrifluconazole | | u | u | u | 0d | NA |
| Fontelis | 7 | x | x | 24 fl. oz. | 12h | 72 fl. oz. |
| penthiopyrad | | x | x | E | 0d | NA |

(Continued)

Table 7-5. Blueberry Diseases - Summer Cover¹ (continued)

| Product And Formulation Active Ingredient | FRAC Code ² | Alternaria Fruit Rot | Anthracnose Fruit Rot | Phomopsis Cane Blight | REI ³ PHI ⁴ | Max Amt ⁵ Max App ⁶ |
|--|------------------------|-------------------------|--------------------------|--------------------------|--------------------------------------|--|
| Indar 2F | 3 | 6 fl. oz. | 6 fl. oz. | 6 fl. oz. | 12h | 24 fl. oz. |
| fenbuconazole | | F | u | G | 30d | 4 |
| Inspire Super | 3+9 | 16-20 fl. oz. | 16-20 fl. oz. | x | 12h | fl. oz. |
| difenoconazole + cyprodinil | | u | u | x | 7d | NA |
| Kenja 400SC | 7 | x | 13.5-15.5 fl. oz. | x | 12h | 54 fl. oz. |
| Isofetamid | | x | u | x | 7d | NA |
| Kocide 3000 | M | 1-2.25 lb. | 1-2.25 lb. | x | 48h | 28 lb. |
| copper hydroxide | | F-P | F | x | 0d | varies |
| Luna Flex | 3+7 | 6.8 fl. oz. | x | 11.2-13.6 fl. oz. | 12h | 27.2 fl. oz. |
| fluopyram + difencolazole | | E | x | G | 7d | 2 |
| Luna Tranquility (SC) | 7+9 | 13.6-27 fl. oz. | 13.6-27 fl. oz. | 13.6-27 fl. oz. | 12h | 54.7 fl. oz. |
| fluopyram + pyrimethanil | | G-F | G-F | u | 0d | NA |
| Miravis Prime | 7+12 | 9-13.4 fl. oz. | 13.4 fl. oz. | 9-13.4 fl. oz. | 12h | 26.8 fl. oz. |
| pydiflumetofen+fludioxonil | | G | G | G | 0d | NA |
| Omega 500F | 29 | 20 fl. oz. | 20 fl. oz. | 20 fl. oz. | 12h | 120 fl. oz. |
| fluazinam | | F | G | u | 30d | NA |
| PhD | 19 | 6.2 oz. | 6.2 oz. | 6.2 oz. | 4h | NA |
| polyoxin D | | P | P | G-F | 0d | 6 |
| Pristine (38WG) | 11+7 | 18.5-23 oz. | 18.5-23 oz. | 18.5-23 oz. | 24h | 92 oz. |
| pyraclostrobin + boscalid | | G | E | G | 0d | 4 |
| Proline 480C | 3 | 5.7 fl. oz. | 5.7 fl. oz. | 5.7 fl. oz. | 12h | 11.4 oz. |
| prothioconazole | | F | u | E-G | 7d | 2 |
| ProPhyt | P07 (33) | 4 pt. | 4 pt. | x | 4h | NA |
| potassium phosphite | | F-P | F | x | 0d | NA |
| Quadris Top | 11+3 | 12-14 fl. oz. | 12-14 fl. oz. | x | 12h | 56 fl. oz. |
| azoxystrobin + difenoconazole | | G | F | x | 7d | 4 |
| Quash SC | 3 | 2.5 oz. | 2.5 oz. | 2.5 oz. | 12h | 7.5 oz. |
| metconazole | | u | G | E | 7d | 3 |
| Quilt Xcel | 11+3 | x | 14-21 fl. oz. | 14-21 fl. oz. | 12h | 63 fl. oz. |
| azoxystrobin + propiconazole | | x | G | G | 30d | NA |
| Switch 62.5WG | 9+12 | 11-14 oz. | 11-14 oz. | 11-14 oz. | 12h | 56 oz. |
| cyprodinil + fludioxonil | | E | G | F | 0d | NA |
| Tilt | 3 | x | x | x | 12h | 30 fl. oz. |
| propiconazole | | x | x | x | 30d | 5 |
| Ziram 76DF | M3 | 3 lb. | 3 lb. | 3 lb. | 48h | NA |
| ziram | | F | G | G | 30d | NA |

Footnotes: See how to read the spray schedule tables section at the beginning of the chapter.

(Continued)

Blueberry First to Second Cover - Insects

- Control cranberry fruitworm 10 and 20 days after petalfall.
- Plum curculio adults and larvae have not been observed to damage blueberries in most southern portions of the region.
- Monitor for first emergence of blueberry maggot flies with traps. Emergence usually begins around July 1 in northern areas. Insecticide applications to protect berries may be needed until harvest. Blueberry maggot is not a common pest in the southern portion of the region.

Table 7-6. Blueberry Insects - First Through Second Cover¹

| Product And Formulation | Active Ingredient | IRAC Code ² | Cherry Fruitworm/ Cranberry Fruit- worm | Plum Curculio | REI ³ PHI ⁴ | Max Amt ⁵ Max App ⁶ |
|--|--|------------------------|---|-------------------|--------------------------------------|--|
| Apta (1.34SC) | | 21A | 27 fl. oz. | 27 fl. oz. | 12h | 81 fl. oz. |
| | tolfenpyrad | | G | G | 3d | 3 |
| Asana XL (0.66EC) (RUP) | | 3A | 4.8-9.6 fl. oz. | x | 12h | 38.4 fl. oz. |
| | esfenvalerate | | G | x | 14d | NA |
| Assail 30SC | | 4A | 3.8-4.5 fl. oz. | x | 12h | 22.5 fl. oz. |
| | acetamiprid | | G | x | 1d | 5 |
| Avaunt eVo (30WDG) | | 22 | 3.5-6 oz. | 6 oz. | 12h | 24 oz. |
| | indoxacarb | | G | E | 7d | 4 |
| B.t. kurstaki (DiPel DF, Javelin WG, etc.) | | 11 | 0.5-2 lb. | x | 4h | NA |
| | <i>Bacillus thuringiensis kurstaki</i> | | G | x | 0d | NA |
| Brigade 2EC (RUP) | | 3A | 2.1-6.4 fl. oz. | 2.1-6.4 fl. oz. | 12h | 32 fl. oz. |
| | bifenthrin | | G | G | 1d | 5 |
| Brigade WSB (10WP) (RUP) | | 3A | 5.3-16 oz. | 5.3-16 oz. | 12h | 80 oz. |
| | bifenthrin | | G | G | 1d | 5 |
| Danitol 2.4EC (RUP) | | 3A | 10.6-16 oz. | 10.6-16 oz. | 24h | 32 fl. oz. |
| | fenpropathrin | | E | G | 3d | 2 |
| Delegate WG (25WG) | | 5 | 3-6 oz. | x | 4h | 19.5 oz. |
| | spinetoram | | G | x | 3d | 6 |
| Diazinon AG600 WBC (RUP) | | 1B | 12.75 fl. oz. | x | 120h | 25.5 fl. oz. |
| | diazinon | | u | x | 7d | 2 |
| Entrust SC (2SC) | | 5 | 4-6 fl. oz. | x | 4h | 29 fl. oz. |
| | spinosad | | F | x | 1d | 6 |
| Esteem 35WP | | 7C | 5 oz. | x | 12h | 10 oz. |
| | pyriproxyfen | | E | x | 7d | 2 |
| Exirel (0.83SE) | | 28 | 10-13.5 fl. oz. | 13.5-20.5 fl. oz. | 12h | 61.5 fl. oz. |
| | cyantraniliprole | | G | G | 3d | NA |
| Grandevo WDG | | UN | 1-3 lb. | x | 4h | NA |
| | <i>Chromobacterium subtsugae</i> | | E | x | 0d | NA |
| Imidan 70W | | 1B | 1.3 lb. | 1.3 lb. | 24-72h | 7.1 lb. |
| | phosmet | | E | E | 3d | 5 |
| Intrepid 2F | | 18 | 10-16 fl. oz. | x | 4h | 48 fl. oz. |
| | methoxyfenozide | | F | x | 7d | 3 |

(Continued)

Table 7-6. Blueberry Insects - First Through Second Cover¹ (continued)

| Product And Formulation | Active Ingredient | IRAC Code ² | Cherry Fruitworm/ Cranberry Fruit- worm | Plum Curculio | REI ³ PHI ⁴ | Max Amt ⁵ Max App ⁶ |
|-------------------------------|-------------------|------------------------|---|----------------|--------------------------------------|--|
| Knack (0.86EC) | | 7C | 16 fl. oz. | x | 12h | 32 fl. oz. |
| | pyriproxyfen | | E | x | 7d | 2 |
| Lannate LV (2.4WSL) (RUP) | | 1A | 1.5-3 pt. | x | 48h | 12 pt. |
| | methomyl | | u | x | 3d | 4 |
| Malathion 8F | | 1B | 1.25 pt. | 1.25 pt. | 12h | NA |
| | malathion | | u | F | 1d | 3 |
| Neemix 4.5 (0.39L), AzaDirect | | UN | 4-16 fl. oz. | x | 4h | NA |
| | azadirachtin | | u | x | 0d | NA |
| Pyganic 5EC | | 3A | 4.5-15.6 fl. oz. | x | 12h | NA |
| | pyrethrins | | u | x | 0d | 10 |
| Rimon 0.83EC | | 15 | 20-30 fl. oz. | x | 12h | 90 fl. oz. |
| | novaluron | | G | x | 8d | NA |
| Sevin XLR Plus (4F) | | 1A | 1.5-2 qt. | x | 12h | 10 qt. |
| | carbaryl | | G | x | 7d | 5 |
| Verdepryn 100SL (0.83SL) | | 28 | 8.2-11 fl. oz. | 8.2-11 fl. oz. | 4h | 33 fl. oz. |
| | cyclaniliprole | | G | G | 1d | 3 |

Footnotes: See how to read the spray schedule tables section at the beginning of the chapter.

Blueberry Third To Summer Covers - Insects

Apply about 10 days after previous cover and repeat as needed. Be sure to check PHIs.

- Pre- and post-harvest rots can be greatly reduced by timely harvests and proper handling, followed by rapid, post-harvest cooling. Fungicides alone are not sufficient to control these diseases. Do not harvest or handle wet fruit.

- Sprays for blueberry maggot should begin as soon as the adults are observed in traps and continue until harvest.

Table 7-7. Blueberry Insects - Third Through Summer Covers¹

| Product And Formulation | Active Ingredient | IRAC Code ² | Blueberry Maggot | Brown Marmorated Stink Bug | Japanese Beetle | Spotted-Wing Drosophila | REI ³ PHI ⁴ | Max Amt ⁵ Max App ⁶ |
|-------------------------|---------------------|------------------------|------------------|----------------------------|-----------------|-------------------------|--------------------------------------|--|
| Admire Pro (4.6F) | | 4A | 2.1-2.8 fl. oz. | x | 2.1-2.8 fl. oz. | x | 12h | 14 fl. oz. |
| | imidacloprid | | F | x | F | x | 3d | 5 |
| Altacor (35WDG) | | 28 | x | x | 3-4.5 oz. | x | 4h | 9 oz. |
| | chlorantraniliprole | | x | x | i | x | 1d | NA |
| Apta (1.34SC) | | 21A | 27 fl. oz. | x | x | x | 12h | 81 fl. oz. |
| | tolfenpyrad | | F | x | x | x | 3d | 3 |
| Asana XL (0.66EC) (RUP) | | 3A | 9.6 fl. oz. | x | 4.8-9.6 fl. oz. | x | 12h | 38.4 fl. oz. |
| | esfenvalerate | | G | x | G | x | 14d | NA |

(Continued)

Table 7-7. Blueberry Insects - Third Through Summer Covers¹ (continued)

| Product And Formulation Active Ingredient | IRAC Code ² | Blueberry Maggot | Brown Marmorated Stink Bug | Japanese Beetle | Spotted-Wing Drosophila | REI ³ PHI ⁴ | Max Amt ⁵ Max App ⁶ |
|--|---------------------------|---------------------|----------------------------------|--------------------|----------------------------|--------------------------------------|--|
| Assail 30SG | 4A | 3.8-4.5 fl. oz. | x | 3.8-4.5 fl. oz. | x | 12h | 26.7 oz. |
| acetamiprid | | G | x | u | x | 1d | 5 |
| Aza-Direct | UN | x | 1-3.5 pt. | 1-3.5 pt. | x | 4h | NA |
| azadirachtin | | x | F | F | x | 0d | NA |
| Beetle GONE! Ag | UN | x | x | 1-17.5 lb. | x | 4h | NA |
| <i>Bacillus thuringiensis galleriae</i> | | x | x | u | x | 0d | NA |
| Brigade 2EC (RUP) | 3A | 2.1-6.4 fl. oz. | x | 2.1-6.4 fl. oz. | x | 12h | 32 fl. oz. |
| bifenthrin | | G | x | E | x | 1d | 5 |
| Brigade WSB (10WP) (RUP) | 3A | 5.3-16 oz. | x | x | x | 12h | 80 oz. |
| bifenthrin | | G | x | x | x | 1d | 5 |
| Danitol 2.4EC (RUP) | 3A | 10.6-16 fl. oz. | 10.6-16 fl. oz. | 10.6-16 oz. | 10.6-16 oz. | 24h | 32 fl. oz. |
| fenpropathrin | | G | G | G | E | 3d | 2 |
| Delegate WG (25WG) | 5 | 3-6 oz. | x | x | 3-6 oz. | 4h | 19.5 oz. |
| spinetoram | | F | x | x | E | 3d | 6 |
| Diazinon AG600 WBC (RUP) | 1B | 12.75 fl. oz. | x | x | x | 120h | 25.5 fl. oz. |
| diazinon | | G | x | x | x | 7d | 2 |
| Entrust SC (2SC) | 5 | x | x | x | 4-6 fl. oz. | 4h | 29 fl. oz. |
| spinosad | | x | x | x | G | 1d | 6 |
| Exirel (0.83SE) | 28 | 13.5-20 fl. oz. | x | x | 13.5-20.5 fl. oz. | 12h | 61.5 fl. oz. |
| cyantraniliprole | | F | x | x | E | 3d | NA |
| Grandevo WDG | UN | x | 2-3 lb. | 2-3 lb. | 2-3 lb. | 4h | NA |
| <i>Chromobacterium subsugae</i> | | x | u | s | G | 0d | NA |
| Imidan 70W | 1B | 1.3 lb. | x | 1.3 lb. | 1.3 lb. | 24-72h | 7.1 lb. |
| phosmet | | E | x | G | E | 3d | 5 |
| Lannate LV (2.4WSL) (RUP) | 1A | 0.75-1.5 pt. | 1.5-3 pt. | x | 1.5-3 pt. | 48h | 12 pt. |
| methomyl | | G | G | x | E | 3d | 4 |
| Malathion 8F | 1B | 1.25 pt. | x | 1.25 pt. | x | 12h | NA |
| malathion | | G | x | F | x | 1d | 3 |
| Mustang Maxx (0.83EC) (RUP) | 3A | x | x | x | 4 fl. oz. | 12h | 24 fl. oz. |
| zeta-cypermethrin | | x | x | x | E | 1d | NA |
| Platinum 75SG | 4A | x | x | 1.66-4 oz. | x | 12h | 4 oz. |
| thiamethoxam | | x | x | u | x | 75d | NA |
| Pyganic 5EC | 3A | 4.5-15.6 fl. oz. | 4.5-15.6 fl. oz. | 4.5-15.6 fl. oz. | 4.5-15.6 fl. oz. | 12h | NA |
| pyrethrins | | F | u | u | F | 0d | 10 |
| Rimon 0.83EC | 15 | 20-30 fl. oz. | x | x | 20-30 fl. oz. | 12h | 90 fl. oz. |
| novaluron | | G | x | x | u | 8d | NA |
| Sevin XLR Plus (4F) | 1A | 1.5-2 qt. | x | 1-2 qt. | x | 12h | 10 qt. |
| carbaryl | | G | x | G | x | 7d | 5 |

(Continued)

Table 7-7. Blueberry Insects - Third Through Summer Covers¹ (continued)

| Product And Formulation Active Ingredient | IRAC Code ² | Blueberry Maggot | Brown Marmorated Stink Bug | Japanese Beetle | Spotted-Wing Drosophila | REI ³ PHI ⁴ | Max Amt ⁵ Max App ⁶ |
|--|---------------------------|---------------------|----------------------------------|--------------------|----------------------------|--------------------------------------|--|
| Sivanto Prime (1.67SC) | 4D | 12-14 fl. oz. | x | x | x | 4h | 28 fl. oz. |
| flupyradifurone | | G | x | x | x | 3d | NA |
| Verdepryn 100SL (0.83SL) | 28 | 8.2-11 fl. oz. | 8.2-11 fl. oz. | 8.2-11 fl. oz. | 8.2-11 fl. oz. | 4h | 33 fl. oz. |
| cyclaniliprole | | G | s | u | E | 1d | 3 |

Footnotes: See how to read the spray schedule tables section at the beginning of the chapter.

Blueberry Postharvest - Diseases

Table 7-8. Blueberry Diseases - Postharvest¹

| Product And Formulation Active Ingredient | FRAC Code ² | Phomopsis Cane Blight | Phytophthora Root Rot | Powdery Mildew | REI ³ PHI ⁴ | Max Amt ⁵ Max App ⁶ |
|--|------------------------|--------------------------|--------------------------|-------------------|--------------------------------------|--|
| Abound (SC) | 11 | 6-15.5 fl. oz. | 6-15.5 fl. oz. | 6-15.5 fl. oz. | 4h | 46 fl. oz. |
| azoxystrobin | | F | u | x | 0d | NA |
| Aliette WDG | P07 (33) | 5 lb. | 5 lb. | x | 12h | 20 lb. |
| aluminum tris | | s | G | x | 0.5d | 4 |
| Bravo Weather Stik | M3 | 3-4 pt. | x | 3-4 pt. | 12h | 12 pt. |
| chlorothalonil | | s | x | s | 42d | NA |
| Captan 80WG | M 5 | 1.25-3 lb. | x | 1.25-3 lb. | 48h | 43.75 lb. |
| captan | | s | x | G | 0d | NA |
| Indar 2F | 3 | 6 fl. oz. | x | 6 fl. oz. | 12h | 24 fl. oz. |
| fenbuconazole | | G | x | E | 30d | 4 |
| Inspire Super | 3+9 | x | x | 16-20 fl. oz. | 12h | 80 fl. oz. |
| difenoconazole + cyprodinil | | x | x | u | 7d | NA |
| Kenja 400SC | 7 | x | x | 13.5-15.5 fl. oz. | 12h | 54 fl. oz. |
| Isofetamid | | x | x | G | 7d | NA |
| Kocide 3000 | M | x | x | x | 48h | 28 lb. |
| copper hydroxide | | x | x | x | 0d | varies |
| Luna Flex | 3+7 | 11.2-13.6 fl. oz. | x | 11.2-13.6 fl. oz. | 12h | 27.2 fl. oz. |
| fluopyram + difencolazole | | G | x | E | 7d | 2 |
| Luna Tranquility (SC) | 7+9 | 13.6-27 fl. oz. | x | 13.6-27 fl. oz. | 12h | 54.7 fl. oz. |
| fluopyram + pyrimethanil | | u | x | E | 0d | NA |
| Mettle 125ME | 3 | x | x | 3-5 fl. oz. | 12h | 20 fl. oz. |
| tetraconazole | | x | x | E | 0d | 5 |
| Miravis Prime | 7+12 | 9-13.4 fl. oz. | x | 9-13.4 fl. oz. | 12 h | 26.8 fl. oz. |
| pydiflumetofen+fludioxonil | | G | x | u | 0d | NA |
| Omega 500F | 29 | 20 fl. oz. | x | 20 fl. oz. | 12h | 120 fl. oz. |
| fluazinam | | u | x | G | 30d | NA |
| Orondis Gold | 49+4 | x | 20-55 oz. | x | 48h | 110 fl. oz. |
| oxathiapipralin+ mefenoxam | | x | E | x | 1d | 2 |
| PhD | 19 | 6.2 oz. | x | 6.2 oz. | 4h | NA |
| polyoxin D | | G-F | x | s | 0d | 6 |

(Continued)

Table 7-8. Blueberry Diseases - Postharvest¹ (continued)

| Product And Formulation Active Ingredient | FRAC Code ² | Phomopsis Cane Blight | Phytophthora Root Rot | Powdery Mildew | REI ³ PHI ⁴ | Max Amt ⁵ Max App ⁶ |
|--|------------------------|--------------------------|--------------------------|-------------------|--------------------------------------|--|
| Pristine (38WG) | 11+7 | 18.5-23 oz. | x | 18.5-23 oz. | 24h | 92 oz. |
| pyraclostrobin + boscalid | | G | x | E | 0d | 4 |
| Procure 480SC | 3 | x | x | 4-8 fl. oz. | 12h | 32 oz. |
| triflumizole | | x | x | E | 0d | 8 |
| Proline 480C | 3 | 5.7 fl. oz. | x | 5.7 fl. oz. | 12h | 11.4 oz. |
| prothioconazole | | E-G | x | G | 7d | 2 |
| Prolivo 300 SC | 50 | x | x | 4-5 fl. oz. | 4h | 16 fl. oz. |
| pyriofenone | | x | x | E | 0d | NA |
| ProPhyt | P07 (33) | x | 4 pt. | x | 4h | NA |
| potassium phosphite | | x | G | x | 0d | NA |
| Quadris Top | 11+3 | x | x | 12-14 fl. oz. | 12h | 56 fl. oz. |
| azoxystrobin + difenoconazole | | x | x | G | d | 4 |
| Quash SC | 3 | 2.5 oz. | x | 2.5 oz. | 12h | 7.5 oz. |
| metconazole | | E | x | E | 7d | 3 |
| Quilt Xcel | 11+3 | 14-21 fl. oz. | x | 14-21 fl. oz. | 12h | 63 fl. oz. |
| azoxystrobin + propiconazole | | G | x | E | 30d | NA |
| Ridomil Gold SL | 4 | x | 3.6 pt. | x | 48h | 3.6 pt. |
| mefenoxam | | x | E | x | 0d | NA |
| Sulforix | M | 1-2 gal./100 gal. | x | x | 48h | 8 gal. |
| calcium polysulfide | | u | x | x | NA | 4 |
| Switch 62.5WG | 9+12 | 11-14 oz. | x | x | 12h | 56 oz. |
| cyprodinil + fludioxonil | | F | x | x | 0d | NA |
| Tilt | 3 | x | x | 6 fl. oz. | 12h | 30 fl. oz. |
| propiconazole | | x | x | E | 30d | 5 |
| Torino | U6 | x | x | 3.4 oz. | 4 h | 6.8 oz. |
| cyflufenamid | | x | x | E | 0d | 2 |
| Ziram 76DF | M3 | 3 lb. | x | 3 lb. | 48h | NA |
| ziram | | G | x | u | 30d | NA |

Footnotes: See how to read the spray schedule tables section at the beginning of the chapter.

Special Comments on Blueberry Schedule

Spotted lanternfly

The spotted lanternfly is an invasive planthopper that is spreading throughout the Midwest. This insect feeds on plant sap, causing wilting, dieback and even death.

Currently spotted lanternfly is believed to pose the greatest threat to the blueberry, grape, hop, stone fruit, and hardwood industries. Know how to identify this pest and remain vigilant for its appearance in your vineyard and orchard systems.

For more on blueberry production

For additional information on blueberry production and management, the University of Kentucky has published the Midwest Blueberry Production Guide, available at: <http://www2.ca.uky.edu/agcomm/pubs/ID/ID210/ID210.pdf>

8. RASPBERRY AND BLACKBERRY

Raspberry And Blackberry Spray Schedule

The shaded/colored boxes represent the crop stages where common pests in the Midwest are active. Scouting and/or preventative sprays may be necessary or recommended. Weather and degree day accumulation will impact the exact timing of pest appearance in the orchard.

| Stage | | | | |
|-----------------------|---------------------|-----------------------|-------------------------|-----------------------|
| Delayed Dormant | Pre-bloom | Bloom Thru Petal Fall | Post-bloom Thru Harvest | Postharvest |
| Raspberry Crown Borer | | | | Raspberry Crown Borer |
| | | | Broad Mite | |
| | | | Green June Beetle | |
| | | | Japanese Beetle | |
| | | | Spotted Wing Drosophila | |
| | | | Stink Bugs | |
| | Raspberry Fruitworm | | | |
| | Raspberry Sawfly | | | |
| | Rose Chafer | | | |
| | Strawberry Clipper | | | |
| | | | Plant Bugs | |
| | | | Rednecked Crane Borer | |
| | | | Thrips | |
| | | | Sap Betles | |
| | | | Spider Mites | |

How to read the spray schedule tables

Every raspberry and blackberry growth stage has important notes on disease or insect management. In some cases, the reader will be directed to the special problems section at the end of the section or chapter. Please make sure to read thoroughly and contact your state Extension specialist with any specific questions.

Key to tables

E = excellent control

G = good control

F = fair control

[r] = fungicide/insecticide resistance possible

s = suppression only

i = ineffective

u = unknown efficacy

x = pest not on the label

¹Efficacy data in this publication are based on trials conducted across various regions and does not neces-

sarily reflect local efficacy differences or changes over time. Growers should contact their Extension specialist for the most recent or for state-specific information. The information on this publication is only a guide; the authors and their institutions assume no liability for practices implemented based on this information. Always read and follow pesticide labels. The label is the law. Product registration may vary by state.

² F/IRAC code represents the mode of action of the fungicide/insecticide.

³ PHI refers to the pre-harvest interval, which is the number of days before harvest that the product may not be applied.

⁴ All fungicides/insecticides have a Restricted-Entry Interval (REI). The restricted-entry interval is the time immediately after a pesticide application when entry into the treated area is limited. Check labels for REI. Restrictions in REI may prohibit the use of certain pesticides during harvest.

Applicators must abide by both maximum amount of product per season AND maximum number of applications.

⁵ Max amt refers to the product's maximum amount/acre/year.

⁶ Max app refers to the product's maximum number of applications per year.

RUP refers to restricted use pesticide.

Raspberry and Blackberry Preplant

- Do not plant raspberries following potatoes or alfalfa or where Verticillium is known to occur.
- For Phytophthora root rot prevention an application of Orondis Gold or Ridomil Gold can be made

at planting. A second application of Orondis Gold can be made at least 7-days after the first application.

Raspberry and Blackberry Dormant to Delayed Dormant - Diseases

- Sanitation is a cornerstone of disease and insect management. Remove and destroy old, infected and infested floricanes after harvest to aid in the management of fungal diseases and borers.
- Apply application of line sulfur or Sulforix if anthracnose, spur blight or cane blight were problematic the previous season.

Table 8-1. Raspberry And Blackberry Diseases - Dormant Through Delayed Dormant¹

| Product And Formulation Active Ingredient | FRAC Code ² | Anthracnose | Cane Blight/ Spur Blight | Phytophthora Root Rot | REI ³ PHI ⁴ | MaxAmt ⁵ MaxApp ⁶ |
|--|------------------------|----------------------|-----------------------------|--------------------------|--------------------------------------|--|
| Aliette WDG | P07 (33) | x | x | 5 lb. | 12h | N/A |
| aluminum tris | | x | x | E | 60d | 4 |
| Badge SC | M | 1-2.25 pt. | 1-2.25 pt. | x | 48h | 35.2 pt. |
| copper sulfate + oxychloride | | F | F | u | 0d | N/A |
| Cuproxat FL | M | 2.5-6 pt. | 2.5-5 pt. | x | 12h | varies |
| tribasic copper sulfate | | F | F | x | 0d | N/A |
| Cabrio EG (20EG) | 11 | 14 oz. | 14 oz. | x | 12h | 56 oz. |
| pyraclostrobin | | E | E | u | 0d | N/A |
| Kocide 3000 | M | 1.75 lb. or 0.75 lb. | 1.75 lb. or 0.75 lb. | x | 48h | 28.6 lb. |
| copper hydroxide | | u | F | x | 0d | N/A |
| Lime Sulfur | M | 6-12 gal./100 gal. | 12 gal./100 gal. | x | 48h | 46 |
| calcium polysulfide | | E | G | x | 0d | 16 |
| Nordox 75G | M | 1.25-2.5 lb. | x | x | 24h | 24 lb. |
| cuprous oxide | | F | F | x | 0d | N/A |
| Orondis Gold | 49+4 | x | x | 25-56 fl. oz. | 48h | 220 fl. oz. |
| oxathiapiprolin+ mefenoxam | | 0 | x | G-E | 1d | varies |
| Orondis Gold200 | 49+4 | x | x | 25-56 fl. oz. | 4h | 19.2 fl. oz. |
| oxathiapiprolin | | x | x | G | 1d | 2 |
| ProPhyte | 33 | x | x | 4 pt. | 4h | 2 |
| phosphorous acid | | x | x | E | 0d | 8 pt. |
| Ridomil Gold SL | 4 | x | x | 3.6 pt. | 48h | 7.2 pt. |
| mefenoxam | | x | x | G-E | 45d | 1 |
| Sulforix | M | 3 gal./100 gal. | 3 gal./100 gal. | x | 48h | N/A |
| calcium polysulfide | | E | G | x | 0d | varies |

Footnotes: See how to read the spray schedule tables section at the beginning of the chapter.

Raspberry and Blackberry Dormant To Delayed Dormant - Insects

- Apply insecticides after egg hatch in late October or early November or wait until late March.
- Apply as a soil drench directed at the crown of the plants in a minimum of 50 gal. of water per acre prior to a significant rainfall or irrigation.

- See Rednecked Cane Borer (page 218-219) about pruning to remove last year's galls.

Table 8-2. Raspberry And Blackberry Insects - Dormant Through Delayed Dormant¹

| Product And Formulation Active Ingredient | IRAC Code ³ | Raspberry Crown Borer | REI ³ PHI ⁴ | Max Amt ⁵ Max App ⁶ |
|--|------------------------|--------------------------|--------------------------------------|--|
| Altacor (35WG) | 28 | 1.5-2.2 oz. | 4h | 4.6 oz. |
| chlorantraniliprole | | G | 3d | 3 |
| Brigade WSB (10WP) (RUP) | 3A | 16 oz. | 12h | 32 oz. |
| bifenthrin | | E | 3d | NA |
| Hero (1.24EC) (RUP) | 3A | 10.3 fl. oz. | 12h | 27.4 fl. oz. |
| bifenthrin + zeta-permethrin | | u | 3d | 2 |

Footnotes: See how to read the spray schedule tables section at the beginning of the chapter.

Raspberry and Blackberry Pre-bloom - Diseases

Apply when blossom buds begin to separate from one another (early bud break) or before flowers open)

Delayed prebloom (early bud break) disease management notes

For management of Phytophthora:

- Orondis Gold, Orondis Gold 200, or Ridomil Gold can be applied to established plantings when tips of buds show green.
- Phosphorous acid products (ProPhyt, Phostrol, Fosphite, K-phite, and Rampart) or Alliette (Aluminum tris) can be applied when plants produce new growth of 1-3 inches.

Table 8-3. Raspberry And Blackberry Diseases - Prebloom¹

| Product And Formulation Active Ingredient | FRAC Code ² | Anthracnose | Cane Blight/Spur Blight | Raspberry Leaf Spot/ Septoria Leaf Spot | Rusts (Orange And Late Leaf) | Powdery Mildew | REI ³ PHI ⁴ | Max Amt ⁵ Max App ⁶ |
|--|---------------------------|----------------------|-------------------------------|--|------------------------------------|-------------------|--------------------------------------|--|
| Abound (SC) | 11 | 6-15.5 fl. oz. | 6-15.5 fl. oz. | 6-15.5 fl. oz. | 10-15.5 fl. oz. | 6-15.5 fl. oz. | 4h | 92.3 fl. oz. |
| azoxystrobin | | E | E | E | E | E | 0d | 9 |
| Cabrio EG (20EG) | 11 | 14 oz. | 14 oz. | 14 oz. | 15 oz. | 14 oz. | 12h | 56 oz. |
| pyraclostrobin | | E | E | E | s(E) | E | 0d | N/A |
| Captan 80WDG | M | 2.5 lb. | 2.5 lb. | 2.5 lb. | x | x | 48h | 12.5 lb. |
| captan | | G | G | s | x | x | 3d | N/A |
| Captex 4L | M | 0.75-1 qts./100 gal. | 0.75-1 qts./100 gal. | x | x | x | 72h | 35 qt. |
| captan | | G | G | x | x | x | 3d | N/A |
| CaptEvate 68WDG | M+17 | 3.5 lb. | 3.5 lb. | x | x | x | 48h | 21 lb. |
| captan + fenhexamid | | G | G | x | x | x | 30d | N/A |

(Continued)

Table 8-3. Raspberry And Blackberry Diseases - Prebloom¹ (continued)

| Product And Formulation Active Ingredient | FRAC Code ² | Anthracnose | Cane Blight/Spur Blight | Raspberry Leaf Spot/ Septoria Leaf Spot | Rusts (Orange And Late Leaf) | Powdery Mildew | REI ³ PHI ⁴ | Max Amt ⁵ Max App ⁶ |
|--|---------------------------|---------------|-------------------------------|--|------------------------------------|-------------------|--------------------------------------|--|
| Cevya | 3 | x | x | 4-5 fl. oz. | x | x | 12h | 15 fl. oz. |
| mefentrifluconazole | | x | x | G | x | x | 0d | 3 |
| Fontelis | 7 | x | 14-24 fl. oz. | x | 14-24 fl. oz. | 14-24 fl. oz. | 12h | 72 fl. oz. |
| penthiopyrad | | x | u | x | u | s(G) | 0d | N/A |
| JMS Stylet Oil | M | x | x | x | 3-6 qt. | 3-6 qt. | 4h | N/A |
| mineral oil | | x | x | x | F | F | NL | N/A |
| Kocide 3000 | M | 0.75 lb. | 0.75 lb. | x | 0.75 lb. | x | 48h | 28.6 lb. |
| copper hydroxide | | u | F | x | u | x | 0d | N/A |
| Luna Privilege | 7 | x | x | 4.8-6.4 fl. oz. | x | 4.8-6.4 fl. oz. | 12 h | 13.7 fl. oz. |
| fluopyram | | x | x | G | x | E | 0d | N/A |
| Luna Tranquility (SC) | 7+9 | x | x | 16-27 fl. oz. | x | 13.6-27 fl. oz. | 12h | 54.7 fl. oz. |
| fluopyram + pyrimethanil | | x | x | G-E | x | G | 0d | N/A |
| Merivon | 7+11 | 4-11 fl. oz. | 4-11 fl. oz. | 4-11 fl. oz. | 8-11 fl. oz. | 4-11 fl. oz. | 12 h | 33 fl. oz. |
| pyraclostrobin + fluxopyroxad | | u | E | E | s | G | 0 | 3 |
| Pristine 38WG | 11+7 | 18.5-23 oz. | 18.5-23 oz. | x | 18.5-23 oz. | 18.5-23 oz. | 12h | 92 oz. |
| pyraclostrobin + boscalid | | E | E | E | s | E | 0d | 4 |
| Prolio 300SC | 50 | x | x | x | x | 4-5 fl. oz. | 4h | 16 fl. oz. |
| pyriofenone | | x | x | x | x | E | 0d | 3 or 4 |
| Quilt Xcel | 11+3 | 14-21 fl. oz. | 14-21 fl. oz. | 14-21 fl. oz. | 14-21 fl. oz. | 14-21 fl. oz. | 12h | 63 fl. oz. |
| azoxystrobin + propiconazole | | G | G | G | G | G | 30d | 3 |
| Rally 40WSP | 3 | x | x | x | 1.25-3 oz. | 1.25-3 oz. | 24h | 10 oz. |
| myclobutanil | | x | x | x | E | E | 0d | N/A |
| Sulfur 80WDG | M | x | x | x | x | 6-15 lb. | 24h | varies |
| sulfur | | x | x | x | x | u | 0d | N/A |
| Switch 62.5WG | 9+12 | 11-14 oz. | x | x | x | x | 12h | 56 oz. |
| cyprodinil + fludioxonil | | u | x | x | x | x | 0d | 2 |
| Tanos (DW) | 11+27 | x | 8-10 oz. | 8-10 oz. | x | x | 12h | 72 oz. |
| famoxadone + cymoxanil | | s | G | G | x | x | 0d | N/A |
| Tilt (EC) | 3 | x | 6 fl. oz. | x | 6 fl. oz. | 6 fl. oz. | 12h | 30 fl. oz. |
| propiconazole | | x | u | x | E | E | 30d | 5 |

Footnotes: See how to read the spray schedule tables section at the beginning of the chapter.

Raspberry and Blackberry Pre-bloom - Insects

Table 8-4. Raspberry And Blackberry Insects - Prebloom¹

| Product And Formulation Active Ingredient | IRAC Code ² | Leaf Roller | Raspberry Fruitworm | Raspberry Sawfly | Rose Chafer | Strawberry Clipper | REI ³ PHI ⁴ | Max Amt ⁵ Max App ⁶ |
|--|---------------------------|-------------------|------------------------|---------------------|-------------------|-----------------------|--------------------------------------|--|
| Actara (25WDG) | 4A | x | x | x | x | 3 oz. | 12h | 6 oz. |
| thiamethoxam | | x | x | x | x | G | 3d | NA |
| Agree WG (Dipel, etc.) | 1I | 1-2 lb. | x | x | x | x | 4h | NA |
| <i>B. thuringiensis</i> | | F | x | x | x | x | 0d | NA |
| Altacor eVo (35WG) | 28 | 1.5-2.2 oz. | x | x | x | x | 4h | 4.6 oz. |
| chlorantraniliprole | | E | x | x | x | x | 3d | 3 |
| Asana XL (0.66EC) (RUP) | 3A | 4.8-9.6 fl. oz. | x | x | x | x | 12h | 28.8 fl. oz. |
| esfenvalerate | | E | x | x | x | x | 7d | NA |
| Assail 30SG | 4A | x | 4.5-5.3 oz. | x | x | x | 12h | 26.7 oz. |
| acetamiprid | | x | u | x | x | x | 1d | 5 |
| Brigade WSB (10WP) (RUP) | 3A | 8-16 oz. | x | x | x | x | 12h | 32 oz. |
| bifenthrin | | E | x | x | x | x | 3d | NA |
| Confirm 2F | 18 | 16 fl. oz. | x | x | x | x | 4h | 64 fl. oz. |
| tebufenozide | | E | x | x | x | x | 14d | NA |
| Danitol 2.4EC (RUP) | 3A | 10.6-16 fl. oz. | x | x | x | x | 24h | 32 fl. oz. |
| fenpropathrin | | E | x | x | x | x | 3d | NA |
| Delegate WG (25WG) | 5 | 3-6 oz. | 3-6 oz. | 3-6 oz. | x | x | 4h | 19.5 oz. |
| spinetoram | | E | E | G | x | x | 1d | 6 |
| Entrust SC (2SC) | 5 | 4-6 fl. oz. | 4-6 fl. oz. | 4-6 fl. oz. | x | x | 4h | 29 fl. oz. |
| spinosad | | G | G | G | x | x | 1d | 6 |
| Hero (1.24EC) (RUP) | 3A | 4 to 10.3 fl. oz. | x | x | x | x | 12h | 274 fl. oz. |
| bifenthrin + zeta-permethrin | | G | x | x | x | x | 3d | 2 |
| Intrepid 2F | 18 | 10 to 16 fl. oz. | x | x | x | x | 4h | 48 fl. oz. |
| methoxyfenozide | | G | x | x | x | x | 3d | 3 |
| Malathion 5EC | 1B | x | x | x | 3 pt. | x | 12h | 9.6 pt. |
| malathion | | x | x | x | G | x | 1d | 3 |
| Mustang Maxx (0.83EC) (RUP) | 3A | 4 fl. oz. | x | x | x | x | 12h | 24 fl. oz. |
| zeta-cypermethrin | | E | x | x | x | x | 1d | 6 |
| Neemix 4.5 (0.39L) | UN | 7-16 fl. oz. | x | x | 7-16 fl. oz. | x | 4h | NA |
| azadirachtin | | u | x | x | u | x | 0d | NA |
| Pyganic 5EC | 3A | 4.5-15.61 fl. oz. | 4.5-15.61 fl. oz. | 4.5-15.61 fl. oz. | 4.5-15.61 fl. oz. | 4.5-15.61 fl. oz. | 12h | NA |
| pyrethrins | | F | F | i | F | u | 0d | 10 |
| Sevin XLR Plus (4F) | 1A | 1-2 qt. | 2 qt. | 2 qt. | 1-2 qt. | 1-2 qt. | 12h | 10 qt. |
| carbaryl | | G | i | G | G | G | 7d | NA |
| Surround WP (95WP) | UN | 25-50 lb. | x | x | 25-50 lb. | x | 4h | NA |
| kaolin | | u | x | x | u | x | 0d | NA |

Footnotes: See how to read the spray schedule tables section at the beginning of the chapter.

Raspberry and Blackberry Bloom To Petal Fall - Diseases

Apply when first flowers open through when petals fall.

- Botrytis gray mold:
 - Promoting good air circulation in the canopy is critical for Botrytis gray mold prevention and management as it promotes rapid drying of the leaves and flowers and allows for good fungicide spray coverage.
 - For best management of Botrytis, plan on making three fungicide applications during this period. Apply the first as blooms begin to open, not later than 5% bloom. Make the second at full bloom. Follow with a third as petals begin to fall.
- Botrytis is prone to developing resistance to many fungicides. Rotating modes of action and tank mixing with a multisite fungicide (FRAC M) is highly recommended if more than one fungicide is made during this period.
- Do not mix phosphorus acids with copper after fruit develop to avoid phytotoxicity.

Table 8-5. Raspberry And Blackberry Diseases - Bloom Through Petal Fall¹

| Product And Formulation Active Ingredient | FRAC Code ² | Anthraco-nose | Cane Blight/Spur Blight | Raspberry Leaf Spot/ Septoria Leaf Spot | Botrytis Fruit Rot | Rusts (Orange And Late Leaf) | Powdery Mildew | REI ³ PHI ⁴ | Max Amt ⁵ Max App ⁶ |
|---|------------------------|---------------------|-------------------------|---|---------------------|------------------------------|-------------------|-----------------------------------|---|
| Abound (SC) | 11 | 6-15.5 fl. oz. | 6-15.5 fl. oz. | 6-15.5 fl. oz. | x | 10-15.5 fl. oz. | 6-15.5 fl. oz. | 4h | 92.3 fl. oz. |
| azoxystrobin | | E | E | E | x | E | E | 0d | 9 |
| Cabrio EG (20EG) | 11 | 14 oz. | 14 oz. | 14 oz. | 14 oz. | 14 oz. | 14 oz. | 12h | 56 oz. |
| pyraclostrobin | | E | E | E | s | s | E | 0d | NA |
| Captan 80WDG | M4 | 2.5 lb. | 2.5 lb. | 2.5 lb. | 2.5 lb. | x | x | 48h | 12.5 lb. |
| captan | | G | G | s | G | x | x | 3d | NA |
| Captec 4L | M4 | 0.75-1 qt./100 gal. | 2.5 lb. | 2.5 lb. | 0.75-1 qt./100 gal. | x | x | 72h | 35 qt. |
| captan | | G | G | F | G | x | x | 3d | NA |
| CaptEstate 68WDG | M+17 | 3.5 lb. | 3.5 lb. | x | 3.5 lb. | x | x | 48h | 21 lb. |
| captan + fenhexamid | | G | G | x | E | x | x | 30d | NA |
| Cevya | 3 | x | x | 4-5 fl. oz. | x | x | x | 12h | 15 fl. oz. |
| mefentrifluconazole | | x | x | G | x | x | x | 0d | 3 |
| Elevate 50WDG | 17 | x | x | x | 1.5 lb. | x | x | 12h | 6 lb. |
| fenhexamid | | x | x | x | E | x | x | 0d | NA |
| Fontelis | 7 | x | x | x | 1-2 pt. | x | x | 12h | 72 fl. oz. |
| penthiopyrad | | x | x | x | E | x | x | 0d | NA |
| JMS Stylet Oil | M | x | x | x | x | 3-6 qt. | 3-6 qt. | 4h | NA |
| mineral oil | | x | x | x | x | F | F | NL | NA |
| Kenja 400SC | 7 | x | x | x | 13.5-15.5 fl. oz. | x | 13.5-15.5 fl. oz. | 12h | 54 fl. oz. |
| Isofetamid | | u | x | x | E | x | E | 0d | NA |

(Continued)

Table 8-5. Raspberry And Blackberry Diseases - Bloom Through Petal Fall¹ (continued)

| Product And Formulation Active Ingredient | FRAC Code ² | Anthrac-nose | Cane Blight/Spur Blight | Raspberry Leaf Spot/ Septoria Leaf Spot | Botrytis Fruit Rot | Rusts (Orange And Late Leaf) | Powdery Mildew | REI ³ PHI ⁴ | Max Amt ⁵ Max App ⁶ |
|---|------------------------|-----------------|-------------------------|---|--------------------|------------------------------|-----------------|-----------------------------------|---|
| Luna Tranquility (SC) | 7+9 | x | x | 16-27 fl. oz. | 16-27 fl. oz. | x | 13.6-27 fl. oz. | 12h | 54.7 fl. oz. |
| fluopyram + pyrimethanil | | G | x | G-E | E | x | G | 0d | NA |
| Merivon | 7+11 | 4-11 fl. oz. | 4-11 fl. oz. | 4-11 fl. oz. | 8-11 fl. oz. | 4-11 fl. oz. | 4-11 fl. oz. | 12 h | 33 fl. oz. |
| pyraclostrobin + fluxopyroxad | | E | E | E | s | s | G | 0 | 3 |
| OSO 5%SC | 19 | 6.5-13 fl. oz. | x | x | 6.5-13 fl. oz. | 6.5-13 fl. oz. | 6.5-13 fl. oz. | 4h | 78 fl. oz. |
| polyoxin D | | u | x | x | E | u | G | 0d | 6 |
| Pristine 38WG | 11+7 | 18.5-23 oz. | 18.5-23 oz. | x | 18.5-23 oz. | 18.5-23 oz. | 18.5-23 oz. | 12h | 92 oz. |
| pyraclostrobin + boscalid | | E | E | x | E | s | E | 0d | 4 |
| Prolivo 300SC | 50 | x | x | x | x | x | 4-5 fl. oz. | 4h | 16 fl. oz. |
| pyriofenone | | x | x | x | x | x | E | 0d | 3 or 4 |
| Quilt Xcel | 11+3 | 14-21 fl. oz. | 14-21 fl. oz. | 14-21 fl. oz. | x | 14-21 fl. oz. | 14-21 fl. oz. | 12h | 63 fl. oz. |
| azoxystrobin + propiconazole | | G | G | G | x | G | G | 30d | 3 |
| Rally 40WSP | 3 | x | x | 1.25-3 oz. | x | 1.25-3 oz. | 1.25-3 oz. | 24h | 10 oz. |
| myclobutanil | | x | x | G | x | E | E | 0d | NA |
| Rovral 4F | 2 | x | x | x | 1-2 pt. | x | x | 24h | 8 pt. |
| iprodione | | x | x | x | E | x | x | 0d | 4 |
| Sulfur 80WDG | M | x | x | x | x | x | 6-15 lb. | 24h | varies |
| sulfur | | x | x | x | x | x | F | 0d | NA |
| Switch 62.5WG | 9+12 | 4.8-6.4 fl. oz. | 4.8-6.4 fl. oz. | x | 4.8-6.4 fl. oz. | x | x | 12h | 56 oz. |
| cyprodinil + fludioxonil | | G | G | x | E | x | x | 0d | 2 |
| Tanos (DW) | 11+27 | x | 8-10 oz. | 8-10 oz. | x | x | x | 12h | 72 oz. |
| famoxadone + cymoxanil | | x | G | G | x | x | x | 0d | NA |
| Tilt (EC) | 3 | x | 6 fl. oz. | x | x | 6 fl. oz. | 6 fl. oz. | 12h | 30 fl. oz. |
| propiconazole | | x | u | x | x | E | E | 30d | 5 |

Footnotes: See how to read the spray schedule tables section at the beginning of the chapter.

Raspberry and Blackberry First Bloom To Petal Fall - Insects

Save the bees! Do not apply insecticides during bloom to protect bees and pollinators.

Raspberry and Blackberry Post-bloom To Harvest - Diseases

Disease management notes

Fungicide applications for Botrytis gray mold management at this time can help reduce the incidence of postharvest rot.

- Botrytis is prone to developing resistance to many fungicides. Rotating modes of action and tank mixing with a multisite fungicide (FRAC M) is highly recommended if more than one fungicide is made during this period.

Table 8-6. Raspberry And Blackberry Diseases - Postbloom Through Harvest¹

| Product And Formulation Active Ingredient | FRAC Code ² | Anthracnose | Cane Blight/ Spur Blight | Raspberry Leaf Spot/ Septoria Leaf Spot | Botrytis Fruit Rot | Rusts (Orange And Late Leaf) | Powdery Mildew | Phytophthora Root Rot | REI ³ PHI ⁴ | Max Amt ⁴ Max App ⁵ |
|---|------------------------|---------------------|--------------------------|---|---------------------|------------------------------|-------------------|-----------------------|-----------------------------------|---|
| Abound (SC) | 11 | 6-15.5 fl. oz. | 6-15.5 fl. oz. | 6-15.5 fl. oz. | x | 10-15.5 fl. oz. | 6-15.5 fl. oz. | x | 4h | 92.3 fl. oz. |
| azoxystrobin | | E | E | E | x | E | E | x | 0d | 9 |
| Aliette WDG | P07 (33) | x | x | x | x | x | x | 5 lb. | 12h | 4 |
| aluminum tris | | x | x | x | x | x | x | E | 60d | NA |
| Badge SC | M | 1-2.25 pt. | 1-2.25 pt. | x | x | x | x | x | 48h | 35.2 pt. |
| copper sulfate + oxychloride | | F | F | x | x | x | x | x | 0d | NA |
| Cuproxat FL | M | 2.5-6 pt. | 2.5-5 pt. | 2.5-5 pt. | x | 2.5-5 pt. | x | x | 12h | varies |
| tribasic copper sulfate | | F | F | F | x | F | x | x | 0d | NA |
| Cabrio EG (20EG) | 11 | 14 oz. | 14 oz. | 14 oz. | 14 oz. | 14 oz. | 14 oz. | x | 12h | 56 oz. |
| pyraclostrobin | | E | E | E | s | s | E | u | 0d | NA |
| Captan 80WDG | M4 | 2.5 lb. | 2.5 lb. | 2.5 lb. | 2.5 lb. | x | x | x | 48h | 12.5 lb. |
| captan | | G | G | s | G | x | x | x | 3d | NA |
| Captec 4L | M4 | 0.75-1 qt./100 gal. | 0.75-1 qt./100 gal. | x | 0.75-1 qt./100 gal. | x | x | x | 72h | 35 qt. |
| captan | | G | G | x | G | x | x | x | 3d | NA |
| CaptEvote 68WDG | M+17 | 3.5 lb. | 3.5 lb. | x | 3.5 lb. | x | x | x | 48h | 21 lb. |
| captan + fenhexamid | | G | G | x | E | x | x | x | 30d | NA |
| Cevya | 3 | x | x | 4-5 fl. oz. | x | x | x | x | 12h | 15 fl. oz. |
| mefentrifluconazole | | x | x | G | x | x | x | x | 0d | 3 |
| Elevate 50WDG | 17 | x | x | x | 1.5 lb. | x | x | x | 12h | 6 lb. |
| fenhexamid | | x | x | x | E | x | x | x | 0d | NA |
| Fontelis | 7 | x | x | x | 1-2 pt. | x | x | x | 12h | 72 fl. oz. |
| penthiopyrad | | x | x | x | E | x | x | x | 0d | NA |
| JMS Stylet Oil | M | x | x | x | x | 3-6 qt. | 3-6 qt. | x | 4h | NA |
| mineral oil | | x | x | x | x | F | F | x | NL | NA |
| Kenja 400SC | 7 | x | x | x | 13.5-15.5 fl. oz. | x | 13.5-15.5 fl. oz. | x | 12h | 54 fl. oz. |
| Isofetamid | | x | x | x | E | x | E | x | 0d | NA |

(Continued)

Table 8-6. Raspberry And Blackberry Diseases - Postbloom Through Harvest¹ (continued)

| Product And Formulation Active Ingredient | FRAC Code ² | Anthracnose | Cane Blight/ Spur Blight | Raspberry Leaf Spot/ Septoria Leaf Spot | Botrytis Fruit Rot | Rusts (Orange And Late Leaf) | Powdery Mildew | Phytophthora Root Rot | REI ³ PHI ⁴ | Max Amt ⁴ Max App ⁵ |
|---|------------------------|----------------|--------------------------|---|--------------------|------------------------------|-----------------|-----------------------|-----------------------------------|---|
| Kocide 3000 | M | x | 0.75 lb. | 0.75 lb. | 0.75 lb. | 0.75 lb. | 0.75 lb. | x | 48h | 28.6 lb. |
| copper hydroxide | | x | F | u | u | u | u | x | 0d | NA |
| Luna Tranquility (SC) | 7+9 | x | x | 16-27 fl. oz. | 16-27 fl. oz. | x | 13.6-27 fl. oz. | x | 12h | 54.7 fl. oz. |
| fluopyram + pyrimethanil | | x | x | G-E | E | x | G | x | 0d | NA |
| Merivon | 7+11 | 4-11 fl. oz. | 4-11 fl. oz. | 4-11 fl. oz. | 8-11 fl. oz. | 4-11 fl. oz. | 4-11 fl. oz. | x | 12 h | 33 fl. oz. |
| pyraclostrobin + fluxapyroxad | | E | E | E | s | s | G | x | 0 | 3 |
| Nordox 75G | M | 1.25-2.5 lb. | x | x | x | 1.25-2.5 lb. | x | x | 24h | 24 lb. |
| cuprous oxide | | F | x | x | x | u | x | x | 0d | NA |
| Orondis Gold | 49+4 | x | x | x | x | x | x | 13.7-110 fl. oz. | 48 h | 220 fl. oz. |
| oxathiapiprolin + mefenoxam | | x | x | x | x | x | x | E | 1d | 2 |
| OSO 5%SC | 19 | 6.5-13 fl. oz. | x | x | 6.5-13 fl. oz. | 6.5-13 fl. oz. | 6.5-13 fl. oz. | x | 4h | 78 fl. oz. |
| polyoxin D | | u | x | x | E | u | G | x | 0d | 6 |
| Pristine 38WG | 11+7 | 18.5-23 oz. | 18.5-23 oz. | x | 18.5-23 oz. | 18.5-23 oz. | 18.5-23 oz. | x | 12h | 92 oz. |
| pyraclostrobin + boscalid | | E | E | x | E | S | E | x | 0d | 4 |
| Prolivo 300SC | 50 | x | x | x | x | x | 4-5 fl. oz. | x | 4h | 3 or 4 |
| pyriofenone | | x | x | x | x | x | E | x | 0d | NA |
| ProPhyt | P07 (33) | x | x | x | x | x | x | 4 pt. | 4h | varies |
| phosphorous acid | | x | x | x | x | x | x | E | 0d | 4 |
| Quilt Xcel | 11+3 | 14-21 fl. oz. | 14-21 fl. oz. | 14-21 fl. oz. | x | 14-21 fl. oz. | 14-21 fl. oz. | x | 12h | 63 fl. oz. |
| azoxystrobin + propiconazole | | G | G | G | x | G | G | x | 30d | 3 |
| Rally 40WSP | 3 | x | x | x | x | 1.25-3 oz. | 1.25-3 oz. | x | 24h | 10 oz. |
| myclobutanil | | x | x | x | x | E | E | x | 0d | NA |
| Ridomil Gold SL | 4 | x | x | x | x | x | x | 3.6 pt. | 48h | 3.6 pt. |
| mefenoxam | | x | x | x | x | x | x | E | 45d | 1 |
| Rovral 4F | 2 | x | x | x | 2 pt. | x | x | x | 24h | 8 pt. |
| iprodione | | x | x | x | E | x | x | x | 0d | 4 |
| Sulfur 80-WDG | M | x | x | x | x | x | 6-15 lb. | x | 24h | varies |
| sulfur | | x | x | x | x | x | F | x | 0d | NA |

(Continued)

Table 8-6. Raspberry And Blackberry Diseases - Postbloom Through Harvest¹ (continued)

| Product And Formulation Active Ingredient | FRAC Code ² | Anthracnose | Cane Blight/ Spur Blight | Raspberry Leaf Spot/ Septoria Leaf Spot | Botrytis Fruit Rot | Rusts (Orange And Late Leaf) | Powdery Mildew | Phytophthora Root Rot | REI ³ PHI ⁴ | Max Amt ⁴ Max App ⁵ |
|---|------------------------|-------------|--------------------------|---|--------------------|------------------------------|----------------|-----------------------|-----------------------------------|---|
| Switch 62.5WG | 9+12 | 11-14 oz. | x | x | x | x | x | x | 12h | 56 oz. |
| cyprodinil + fludioxonil | | u | x | x | x | x | x | x | 0d | 2 |
| Tanos (DW) | 11+27 | 8-10 oz. | 8-10 oz. | 8-10 oz. | x | x | x | x | 12h | 72 oz. |
| famoxadone + cymoxanil | | s | G | G | x | x | x | x | 0d | NA |
| Tilt (EC) | 3 | x | 6 fl. oz. | x | x | 6 fl. oz. | 6 fl. oz. | x | 12h | 30 fl. oz. |
| propiconazole | | x | u | x | x | E | E | x | 30d | 5 |

Footnotes: See how to read the spray schedule tables section at the beginning of the chapter.

Raspberry and Blackberry Post-bloom To Harvest - Insects

Check the insecticide label for toxicity to bees and timing of application before selecting insecticide compounds for the Japanese beetle and other insect pests if growing fall-bearing varieties.

Table 8-7. Raspberry And Blackberry Insects - Postbloom Through Harvest¹

| Product And Formulation Active Ingredient | IRAC Code ² | Broad Mite | Green June Japanese Beetle | Plant Bugs | Rednecked Cane Borer | Sap Beetle | Stink Bug | Spotted Wing Drosophila | Thrips | two-Spotted Spider Mite | REI ³ PHI ⁴ | Max Amt ⁵ Max app ⁶ |
|---|------------------------|-------------|----------------------------|-------------|----------------------|-------------|-----------|-------------------------|--------------|-------------------------|-----------------------------------|---|
| Acramite 50WS | 20D | x | x | x | x | x | x | x | x | 0.75-1 lb. | 12h | NA |
| bifenazate | | x | x | x | x | x | x | x | x | G | 1d | 2 |
| Actara (25WDG) | 4A | x | 3 oz. | 3 oz. | x | x | 3 oz. | x | x | x | 12h | 6 oz. |
| thiamethoxam | | x | G | G | x | x | G | x | x | x | 3d | NA |
| Admire Pro (4.6F) | 4A | x | x | x | 10.5-14 fl. oz. | x | x | x | 7-14 fl. oz. | x | 12h | 14 fl. oz. |
| imidacloprid | | x | x | x | G | x | x | x | F | x | 7d | NA |
| Agri-Mek SC (0.7SC) (RUP) | 6 | 3.5 fl. oz. | x | x | x | x | x | x | x | 1.75-3.5 fl. oz. | 12h | 10.25 fl. oz. |
| abamectin | | E | x | x | x | x | x | x | x | E | 7d | NA |
| Assail 30SG | 4A | x | 4.5-5.3 oz. | 4.5-5.3 oz. | x | 4.5-5.3 oz. | x | x | 4.5-5.3 oz. | x | 12h | 26.7 oz. |
| acetamiprid | | x | G | G | x | G | x | x | u | x | 1d | 5 |

(Continued)

Table 8-7. Raspberry And Blackberry Insets - Postbloom Through Harvest¹ (continued)

| Product And Formulation Active Ingredient | IRAC Code ² | Broad Mite | Green June Japanese Beetle | Plant Bugs | Rednecked Cane Borer | Sap Beetle | Stink Bug | Spotted Wing Drosophila | Thrips | two-Spotted Spider Mite | REI ³ PHI ⁴ | Max Amt ⁵ Max app ⁶ |
|--|---------------------------|------------|-------------------------------|--------------------------|-------------------------|------------|-----------------------------|-----------------------------|--------------------------|----------------------------|--------------------------------------|---|
| BeetleGone! | 1I | x | 1-17.5 lb. | x | x | x | x | x | x | x | 4h | NA |
| <i>B. thuringiensis</i> | | x | u | x | x | x | x | x | x | x | 0d | NA |
| Brigade WSB (10WP) (RUP) | 3A | x | x | x | x | x | x | 8-16 oz. | x | 16 oz. | 12h | 32 oz. |
| bifenthrin | | x | x | x | x | x | x | E | x | F | 3d | NA |
| Danitol 2.4EC (RUP) | 3A | x | 10.6-16 fl. oz. | 10.6-16 fl. oz. | x | x | 10.6- 16 fl. oz. | 10.6-16 fl. oz. | x | 16 fl. oz. | 24h | 32 fl. oz. |
| fenpropathrin | | x | E | E | x | x | E | E | x | F | 3d | NA |
| Delegate WG (25WG) | 5 | x | x | x | x | x | x | 3-6 oz. | 3-6 oz. | x | 4h | 19.5 oz. |
| spinetoram | | x | x | x | x | x | x | E | E | x | 1d | 6 |
| Entrust SC (2SC) | 5 | x | x | x | x | x | x | 4-6 fl. oz. | x | x | 4h | 29 fl. oz. |
| spinosad | | x | x | x | x | x | x | G | x | x | 1d | 6 |
| Hero (1.24EC) (RUP) | 3A | x | x | x | x | x | x | x | x | 10.3 fl. oz. | 12h | 27.4 fl. oz. |
| bifenthrin + zeta-permethrin | | x | x | x | x | x | x | x | x | F | 3d | 2 |
| Kanemite 15SC | 20B | x | x | x | x | x | x | x | x | 31 fl. oz. | 12h | 62 fl. oz. |
| acequinocyl | | x | x | x | x | x | x | x | x | G | 1d | 2 |
| M-Pede | UN | 1-2% | x | x | x | x | x | x | x | 1-2% | 12h | NA |
| potassium salts of fatty acids | | E | x | x | x | x | x | x | x | u | 0d | NA |
| Magister (1.7SC) | 21A | x | x | x | x | x | x | x | x | 32-36 fl. oz. | 12h | 36 fl. oz. |
| fenazaquin | | x | x | x | x | x | x | x | x | u | 7d | 1 |
| Malathion 5EC | 1B | x | 3 pt. | x | x | x | x | x | 3 pt. | x | 12h | 9.6 pt. |
| malathion | | x | G | x | x | x | x | x | G | x | 1d | 3 |
| Mustang Maxx (0.83EC) (RUP) | 3A | x | 4 fl. oz. | x | x | x | x | 4 fl. oz. | x | x | 12h | 24 fl. oz. |
| zeta-cypermethrin | | x | E | x | x | x | x | E | x | x | 1d | 6 |
| Neemix 4.5 (0.39L) | UN | x | fl. oz. | x | x | x | x | x | 7-16 fl. oz. | x | 4h | NA |
| azadirachtin | | x | u | x | x | x | x | x | u | x | 0d | NA |
| Pyganic 5EC | 3A | x | 4.5-15.61 fl. oz. | 4.5- 15.61 fl. oz. | x | x | 4.5- 15.61 fl. oz. | 4.5- 15.61 fl. oz. | 4.5- 15.61 fl. oz. | x | 12h | NA |
| pyrethrins | | x | i | F | x | x | i | F | i | x | 0d | 10 |
| Savey 50DF | 10A | x | x | x | x | x | x | x | x | 4-6 oz. | 12h | 6 oz. |
| hexythiazox | | x | x | x | x | x | x | x | x | E | 3d | 1 |

(Continued)

Table 8-7. Raspberry And Blackberry Insets - Postbloom Through Harvest¹ (continued)

| Product And Formulation Active Ingredient | IRAC Code ² | Broad Mite | Green June Japanese Beetle | Plant Bugs | Rednecked Cane Borer | Sap Beetle | Stink Bug | Spotted Wing Drosophila | Thrips | two-Spotted Spider Mite | REI ³ PHI ⁴ | Max Amt ⁵ Max app ⁶ |
|--|---------------------------|------------|-------------------------------|--------------|-------------------------|------------|-----------|----------------------------|--------------|----------------------------|--------------------------------------|---|
| Sevin XLR Plus (4F) | 1A | x | 1-2 qt. | 1.5-2 qt. | x | x | x | x | x | x | 12h | 10 qt. |
| carbaryl | | x | G | G | x | x | i | x | x | x | 7d | NA |
| Surround WP (95WP) | UN | x | 25-50 lb. | x | x | x | x | x | 25-50 lb. | x | 4h | NA |
| kaolin | | x | u | x | x | x | x | x | u | x | 0d | NA |
| Zeal (72WP) | 10B | x | x | x | x | x | x | x | x | 2-3 oz. | 12h | 3 oz. |
| etoxazole | | x | x | x | x | x | x | x | x | E | 0d | 1 |

Footnotes: See how to read the spray schedule tables section at the beginning of the chapter.

Raspberry And Blackberry Postharvest - Diseases

Disease management notes

- Apply fall spray of any fungicide after old canes are removed.
- When applying any fungicide after harvest, consider maximum rate allowed.

- For best control of Phytophthora, the last fall application of Aliette, Ridomil Gold or Orondis Gold is recommended. Refer to the fungicide labels for the best timing for each of these products.

Table 8-8. Raspberry And Blackberry Diseases - Postharvest¹

| Product And Formulation Active Ingredient | FRAC Code ² | Anthracnose | Cane Blight/Spur Blight | Raspberry Leaf Spot/ Septoria Leaf Spot | Botrytis Fruit Rot | Rusts (Orange And Late Leaf) | Powdery Mildew | REI ³ PHI ⁴ | Max Amt ⁵ Max Amt ⁶ |
|--|---------------------------|---------------------|-------------------------------|--|-----------------------|------------------------------------|-------------------|--------------------------------------|--|
| Abound (SC) | 11 | 6-15.5 fl. oz. | 6-15.5 fl. oz. | 6-15.5 fl. oz. | x | 10-15.5 fl. oz. | 6-15.5 fl. oz. | 4h | 92.3 fl. oz. |
| azoxystrobin | | E | E | E | x | E | E | 0d | 9 |
| Cabrio EG (20EG) | 11 | 14 oz. | 14 oz. | 14 oz. | 14 oz. | 14 oz. | 14 oz. | 12h | 56 oz. |
| pyraclostrobin | | E | E | E | s | s | E | 0d | NA |
| Captan 80WDG | M4 | 2.5 lb. | 2.5 lb. | x | 2.5 lb. | x | x | 48h | 12.5 lb. |
| captan | | G | G | s | G | x | x | 3d | NA |
| Captec 4L | M4 | 0.75-1 qt./100 gal. | 0.75-1 qt./100 gal. | x | 0.75-1 qt./100 gal. | x | x | 72h | 35 qt. |
| captan | | G | G | x | G | x | x | 3d | NA |
| CaptEstate 68WDG | M+17 | 3.5 lb. | 3.5 lb. | x | 3.5 lb. | x | x | 48h | 21 lb. |
| captan + fenhexamid | | G | G | x | E | G | x | 30d | NA |
| Cevya | 3 | x | x | 4-5 fl. oz. | x | x | x | 12h | 15 fl. oz. |
| mefentrifluconazole | | x | x | G | x | x | x | 0d | 3 |

(Continued)

Table 8-8. Raspberry And Blackberry Diseases - Postharvest¹ (continued)

| Product And Formulation Active Ingredient | FRAC Code ² | Anthraco | Cane Blight/Spur Blight | Raspberry Leaf Spot/ Septoria Leaf Spot | Botrytis Fruit Rot | Rusts (Orange And Late Leaf) | Powdery Mildew | REI ³ PHI ⁴ | Max Amt ⁵ Max Amt ⁶ |
|--|------------------------|----------------|-------------------------|--|--------------------|------------------------------|-------------------|--------------------------------------|--|
| Elevate 50WDG | 17 | x | x | x | 1.5 lb. | x | x | 12h | 6 lb. |
| fenhexamid | | x | x | x | E | x | x | 0d | NA |
| Fontelis | 7 | x | x | x | 1-2 pts. | x | x | 12h | 72 fl. oz. |
| penthiopyrad | | x | x | x | E | x | G | 0d | NA |
| JMS Stylet Oil | M | x | x | x | x | 3-6 qt. | 3-6 qt. | 4h | NA |
| mineral oil | | x | x | x | x | F | F | NL | NA |
| Kenja 400SC | 7 | x | x | x | 13.5-15.5 fl. oz. | x | 13.5-15.5 fl. oz. | 12h | 54 fl. oz. |
| Isofetamid | | u | x | x | G | u | E | 0d | NA |
| Luna Tranquility (SC) | 7+9 | x | x | 16-27 fl. oz. | 16-27 fl. oz. | x | 13.6-27 fl. oz. | 12h | 54.7 fl. oz. |
| fluopyram + pyrimethanil | | x | x | G-E | E | x | G | 0d | NA |
| OSO 5%SC | 19 | 6.5-13 fl. oz. | x | x | 6.5-13 fl. oz. | 6.5-13 fl. oz. | 6.5-13 fl. oz. | 4h | 78 fl. oz. |
| polyoxin D | | u | x | x | F-G | u | G | 0d | 6 |
| Pristine 38WG | 11+7 | 18.5-23 oz. | 18.5-23 oz. | x | 18.5-23 oz. | 18.5-23 oz. | 18.5-23 oz. | 12h | 92 oz. |
| pyraclostrobin + boscalid | | E | E | E | E[r] | S | E | 0d | 4 |
| Prolivo 300SC | 50 | x | x | x | x | x | 4-5 fl. oz. | 4h | 16 fl. oz. |
| pyriofenone | | x | x | x | x | x | E | 0d | 3 or 4 |
| Quilt Xcel | 11+3 | 14-21 fl. oz. | 14-21 fl. oz. | 14-21 fl. oz. | x | 14-21 fl. oz. | 14-21 fl. oz. | 12h | 63 fl. oz. |
| azoxystrobin + propiconazole | | G | G | G | G | G | G | 30d | 3 |
| Rally 40WSP | 3 | x | x | x | x | 1.25-3 oz. | 1.25-3 oz. | 24h | 10 oz. |
| myclobutanil | | x | x | x | x | E | E | 0d | NA |
| Rovral 4F | 2 | x | x | x | 2 pt. | x | x | 24h | 8 pt. |
| iprodione | | x | x | x | E[r] | x | x | 0d | 4 |
| Sulfur 80WDG | M | x | x | x | x | x | 6-15 lb. | 24h | varies |
| sulfur | | x | x | x | x | x | u | 0d | NA |
| Switch 62.5WG | 9+12 | 11-14 oz. | x | x | 11-14 oz. | x | x | 12h | 56 oz. |
| cyprodinil + fludioxonil | | u | x | x | G | x | x | 0d | 2 |
| Tanos (DW) | 11+27 | x | 8-10 oz. | 8-10 oz. | x | x | x | 12h | 72 oz. |
| famoxadone + cymoxanil | | x | G | G | x | x | x | 0d | NA |
| Tilt (EC) | 3 | x | fl. oz. | x | x | 6 fl. oz. | 6 fl. oz. | 12h | 30 fl. oz. |
| propiconazole | | x | u | x | x | E | E | 30d | 5 |

Footnotes: See how to read the spray schedule tables section at the beginning of the chapter.

Raspberry and Blackberry Postharvest - Insects

Insect management notes

- Apply insecticides for **borer** control during October and November.

Table 8-9. Raspberry And Blackberry Insects - Postharvest¹

| Product And Formulation Active Ingredient | IRAC Code ² | Raspberry Crown Borer | REI ³ PHI ⁴ | Max Amt ⁵ Max App ⁶ |
|--|------------------------|--------------------------|--------------------------------------|--|
| Altacor eVo (35WG) | 28 | 1.5-2.2 oz. | 4h | 4.6 oz. |
| chlorantranilprole | | G | 3d | 3 |
| Brigade WSB (10WP) (RUP) | 3A | 16 oz. | 12h | 32 oz. |
| bifenthrin | | E | 3d | NA |
| Hero (1.24EC) (RUP) | 3A | 10.3 fl. oz. | 12h | 274 fl. oz. |
| bifenthrin + zeta-permethrin | | u | 3d | 2 |

Footnotes: See how to read the spray schedule tables section at the beginning of the chapter.

Special Comments on Raspberry and Blackberry Schedule

Spotted-wing Drosophila

See page 181 (under Grapes).

Broad mite

The broad mite damages the bud, leaf, or flower. This feeding injects a toxin that stunts growth, curls and bronzes leaves, and often kills terminal and lateral leaf and flower buds. Broad mites have damaged floricane-fruiting blackberry cultivars. Infested floricanes have delayed bud break and low vigor in spring. Broad mites can be found on terminal floricane leaves from April through harvest. Primocane terminals can also become infested. These symptoms are similar to those of fire blight.

The mite overwinters under blackberry bud scales and in the soil and in litter under plants. Eggs are oval and spotted (0.08 mm long). Broad mites are oval and vary from small white immatures to amber adults (0.2mm) with white hourglass mark on back of females.

From late May through fall in Arkansas, you can find a buildup of broad mites on the terminal leaves of emerging primocanes. In more northern states, broad mite numbers increase and damage appears later in the summer or early fall.

The only recommended miticide is Agri-Mek. One application has reduced and maintained broad mite numbers to near zero for up to a month. Additional applications may be needed if mite numbers resurge. You

can reapply Agri-Mek once and then you must rotate to a different mode of action. Other products that have significantly reduced broad mite on blackberry include Microthiol Disperss wettable sulfur (10 pounds per acre), 2% JMS Stylet-Oil, and 1% M-Pede. For these products, check safety to blooms by testing a few plants prior to broad application, and do not apply if temperatures are expected to exceed 90°F. See labels for use and rate recommendations.

For more information about broad mites, including photos, see Managing Broad Mite in Southeastern Caneberry Plantings at: <https://smallfruits.org/2021/04/managing-broad-mite-in-southeastern-caneberry-plantings>

Brown marmorated stink bug

The brown marmorated stink bug (BMSB) has an extremely wide host range and is a pest of all small fruit. BMSB is attracted to fruits throughout much of the growing season. It has piercing sucking mouthparts, which cause injury that may appear as sunken areas on the fruit.

Actara, Brigade, and Danitol have shown good efficacy in trials; however, multiple applications may be needed for reinfestations.

Raspberry cane maggot

The raspberry cane maggot causes wilted tips in May. Cut off wilted tips a few inches below the girdle when first seen. Destroy the removed tips.

Rednecked cane borer

Scout for galls before or during the dormant period. Prune out galled canes and burn, bury, or otherwise destroy them to kill overwintered larvae. If more than 5 percent of all canes have galls, an insecticide application immediately after bloom may be warranted.

Adults begin to emerge in May or June. Begin scouting plantings during bloom by looking for adult beetles active during daylight hours. Begin insecticide application(s) after bloom has ended and bees are no longer present. Apply Admire Pro via drip or trickle chemigation or in a soil drench in a minimum of 500 gallons of water per acre. Do not apply pre-bloom or during bloom or when bees are actively foraging.

Phytophthora root rot

Ridomil Gold SL, Ridomil Gold, and Orondis Gold 200 are all labeled for control of Phytophthora root rot on brambles. See the labels for more detailed information on application rates and timing.

Note: Do not apply Ridomil within 45 days before harvest, or illegal residues may result. See the label for more detailed information.

Many phosphorous acid fungicides are registered for Phytophthora root rot control on blackberry and raspberry, and they all essentially have the same active ingredient. All are foliar sprays. They are highly systemic and move rapidly into leaves and are translocated in the plant to the crown and roots. Recommendations for use vary among products. See labels for use recommendations and restrictions.

Blackberry rosette or double blossom

Rosette is caused by the fungus *Cercospora rubi*. It is a serious disease of blackberry in the southern Midwest (Arkansas, Kentucky, Missouri, and Oklahoma). It is a minor disease of raspberries. FRAC 11 fungicides (e.g., Abound, Pristine, Quilt Xcel) and Switch (cyprodinil + fludioxonil) are highly effective; however, chemical control of this disease under heavy disease pressure has not been successful.

One cultural practice for infected sites is to mow the planting down before flowering to eliminate spore release and infection of emerging primocanes. Although this sacrifices one year of production, the practice may provide short-term control.

Varieties vary in susceptibility. Apache, Ouachita, and Triple Crown are resistant. Chester, Hull, and Navaho are tolerant. Chickasaw, Choctaw, Kiowa, Shawnee, and Illini Hardy are highly susceptible. Other cultivars differ in susceptibility, but all become infected over time.

For more information on blackberry rosette refer to <https://plantpathology.mgcafe.uky.edu/files/ppfs-fr-s-03.pdf>.

Blackberry downy mildew

Blackberry downy mildew is caused by the fungus-like water mold *Peronospora sparsa*. The pathogen spreads systemically and infects both the leaves and fruit. Stunting can occur in infected plants even when foliar symptoms are not visible. The disease is most severe during wet weather. Purchase plants from a reputable nursery and inspect them for signs or symptoms of downy mildew before planting. Early symptoms include light green to yellow leaves with brown to red spots, stunting, and red streaking on the stems and petioles. Fungicides containing mefenoxam, oxythiapiprolin or potassium phosphite provide the best level of control. Applications of potassium phosphite can result in phosphorous deficiencies; a balanced nutritional program therefore should be followed and monitored.

Orange rust

All cultivars of black and purple raspberries and most erect and trailing blackberries are very susceptible to orange rust. Unlike all other fungi infecting brambles, this fungus grows systemically throughout the roots, crowns, and shoots of infected plants and is perennial in belowground plant parts. Plants do not die but become stunted and weakened, producing little to no fruit. Key control methods include cultural practices such as removing infected plants early in the spring and eradicating nearby wild brambles. Alternate Rally with Abound (or another axozystrobin product), Cabrio, or Pristine in the spray program to prevent fungicide resistance development.

Raspberry leaf spot and Septoria leaf spot of blackberry and raspberry

The incidence of raspberry leaf spot and Septoria leaf spot is increasing across the Midwest. If not controlled, these diseases can result in severe defoliation of the plant.

The strobilurin fungicides (Abound, Cabrio, Pristine) provide good control of both diseases. Abound is registered for control of raspberry leaf spot and Septoria leaf spot. Some fungicide trials have shown that Captan and Rally also provide some level of control. Post-harvest (late-season) applications are important for controlling these leaf diseases. Most defoliation resulting from these diseases occurs later in the season (post-harvest).

Fungicide resistance management

Elevate, Rovral, Switch, and Pristine should not be used alone for season-long control of Botrytis fruit rot, because some Botrytis cinerea strains may develop resistance to these fungicides. Adding (tank mixing) Captan to Elevate, Rovral, Switch, or Pristine should

enhance disease control and help prevent fungicide resistance development. Rotating these fungicides in blocks of one or two sprays is a good resistance management strategy.

9. STRAWBERRY

Strawberry Spray Schedule

The boxes represent the crop stages where common pests in the Midwest are active. Scouting and/or preventative sprays may be necessary or recommended. Weather and degree day accumulation will impact the exact timing of pest appearance in the orchard.

| Stage | | | |
|-----------------------|-------|-------------------------|------------------------------|
| Pre-bloom | Bloom | Post-bloom Thru Harvest | Post-bloom And New Plantings |
| Tarnished Plant Bug | | Tarnished Plant Bug | |
| Spittlebug | | Spittlebug | |
| Spider Mite | | Spider Mite | |
| | | Slugs | |
| | | Sap Beetles | |
| | | Spotted wing Drosophila | |
| Eastern Flower Thrips | | Eastern Flower Thrips | |
| Cyclamen Mite | | | |
| Bud Clipper | | | |
| | | Leafroller | |
| | | Leafhopper | |
| | | | Root Weevils |

How to read the spray schedule tables

Every strawberry growth stage has important notes on disease or insect management. In some cases, the reader will be directed to the special problems section at the end of the section or chapter. Please make sure to read thoroughly and contact your state Extension Specialist with any specific questions.

Key to tables

E = excellent control

G = good control

F = fair control

[r] = fungicide/insecticide resistance possible

s = suppression only

i = ineffective

u = unknown efficacy

x = pest not on the label

¹ Efficacy data in this publication are based on trials conducted across various regions and does not necessarily reflect local efficacy differences or changes over time. Growers should contact their Extension Specialist for the most recent or for state-specific information. The information on this publication is only a guide; the

authors and their institutions assume no liability for practices implemented based on this information. Always read and follow pesticide labels. The label is the law. Product registration may vary by state.

² F/IRAC code represents the mode of action of the fungicide/insecticide.

³ PHI refers to the pre-harvest interval, which is the number of days before harvest that the product may not be applied.

⁴ All fungicides/insecticides have a Restricted-Entry Interval (REI). The restricted-entry interval is the time immediately after a pesticide application when entry into the treated area is limited. Check labels for REI. Restrictions in REI may prohibit the use of certain pesticides during harvest.

Applicators must abide by both maximum amount of product per season AND maximum number of applications.

⁵ Max amt refers to the product's maximum amount/acre/year.

⁶ Max app refers to the product's maximum number of applications per year.

RUP refers to restricted use pesticide.

Strawberry Pre-plant - Diseases

New Plantings

- For *Phytophthora* spp. prevention, an application of Orondis Gold or Ridomil Gold can be made at planting. A second application of Orondis Gold can be made at least 7-days after the first application.

Fungicide Dips

- Recent research on the role of fungicide dips in disease management have shown inconsistent results. The information is included but not necessarily recom-

mended. Starting with disease-free plants and plugs are essential for successful strawberry production and these treatments will not rescue infected plants.

- Wash excess soil from roots prior to dipping. Dip or expose plants for label recommended time. Drain transplants after dip and plant immediately. Do not reuse solution and dispose of solution as per local regulations.
- When dipping with Abound and Switch, submerge plants for 2-5 min.; when using, Aliette and Rampart, dip for 15-30 min.
- Labeled rates of Aliette differ between dips and foliar application.

Table 9-1. Strawberry Diseases - Preplant¹

| Product And Formulation | Active Ingredient | FRAC Code ² | Anthracnose Crown Rot | Phytophthora (Crown Rot, Red Stele, Leather Rot) | REI ³ PHI ⁴ | Max Amt ⁵ Max App ⁶ |
|-------------------------|---------------------------|------------------------|-----------------------|--|-----------------------------------|---|
| Abound (SC) | | 11 | 5-8 fl. oz. | 5-8 fl. oz. | 4h | 61.5 fl. oz. |
| | azoxystrobin | | G[r] | E | 0d | NA |
| Aliette WDG | | P07 (33) | x | 2.5 lb./100 gal. | 24h | 30 lb. |
| | aluminum tris | | x | E | 12d | NA |
| Phostrol | | 33 | x | 2.5-5 pt./100 gal. | 4h | NA |
| | phosphorous acid | | x | E | NA | NA |
| Prophyte | | P07 (33) | x | 2 pt./100 gal. | 4h | varies |
| | phosphorous acid | | x | x | NA | 4 |
| Orandis Gold | | 49+4 | x | 20-62 fl. oz. | 48h | 124 fl. oz. |
| | Oxathiapiprolin+mefenoxam | | x | E | 14d | 2 |
| Ridomil Gold | | 4 | x | 1 pt. | 48h | 3 pt. |
| | mefenoxam | | x | E | 0d | 3 |
| Switch 62.5WG | | 9+12 | 5-8 oz./100 gal. | x | 12h | 56 oz. |
| | cyprodinil + fludioxonil | | G | x | 0d | NA |

Footnotes: See how to read the spray schedule tables section at the beginning of the chapter.

Strawberry Prebloom - Diseases

Apply when new leaves are expanding and blossom buds are visible.

Disease management notes

- Thiram SC is recommended when the risk of Neopestalotiopsis disease is moderate to high. After two applications alternate with Switch (FRAC 12+9).
- Ridomil is labeled for control of red stele, caused by *Phytophthora fragariae*, and leather rot, caused by *Phytophthora cactorum*. Treatment for perennial strawberries includes one application in the spring

after the ground thaws and before first bloom, and a second application in the fall. Follow label instructions carefully as Ridomil can cause stunting in young plants of some varieties. For supplemental control of leather rot, an application may be made at fruit set.

- Several phosphorous acid fungicides are labeled for control of red stele and leather rot. They all have essentially the same active ingredient. These products include Agri-Fos, Aliette, ProPhyt, Phostrol, and Rampart. These materials are highly systemic as foliar sprays for leather rot control or as root dip for red stele control. Rates, recommendations for use, and prices vary among products.

Table 9-2. Strawberry Diseases - Prebloom¹

| Product And Formulation Active Ingredient | FRAC Code ² | Angular Leaf Spot | Anthraxnose Crown | Leaf Blight Phomopsis | Leaf Scorch (Diplocarpon) | Leaf Spot (Mycosphaerella) | Powdery Mildew | Red Stele (Phytophthora) | REI ³ PHI ⁴ | Max Amt ⁵ Max App ⁶ |
|--|---------------------------|-------------------|----------------------|--------------------------|------------------------------|-------------------------------|-----------------|-----------------------------|--------------------------------------|--|
| Abound (SC) | 11 | x | 5-8 fl. oz. | x | x | x | 6-15.5 fl. oz. | x | 4h | NA |
| azoxystrobin | | x | G[r] | x | x | x | G | x | 0d | NA |
| Actigard 50WG | 21 | 0.5-0.75 oz. | x | x | x | x | x | x | 12h | 6 oz. |
| Acibenzolar-S-methyl: | | s | x | x | x | x | x | x | 0d | NA |
| Aftershock/Evito 480SC | 11 | x | 2-5.7 fl. oz. | x | x | x | 2-5.7 fl. oz. | x | 12h | 22.8 fl. oz. |
| fluoxastrobin | | x | G | x | x | x | G | x | 1d | 4 |
| Aliette WDG | 33 | x | x | x | x | x | x | 2.5-5 lb. | 24h | 30 lb. |
| aluminum tris | | x | x | x | x | x | x | E | 12d | NA |
| Badge SC | M | 1-2.5 pt. | x | 1 – 2.5 pt. | 2-2.5 pt. | x | 1-2.5 pt. | x | 48h | 28.9 pt. |
| copper sulfate + oxychloride | | F | x | F | F | x | F | x | 0d | NA |
| Cabrio EG (20EG) | 11 | x | 12-14 oz. | x | x | 12-14 oz. | 12-14 oz. | 12-14 oz. | 12h | 70 oz. |
| pyraclostrobin | | x | E | x | x | G | E | E | 0d | NA |
| Captan (80WDG) | M | x | 1.8-3.75 lb. | 1.8-3.75 lb. | 1.8-3.75 lb. | 1.8-3.75 lb. | x | x | 24h | 30 lb. |
| captan | | x | E | G | G | G | x | x | 0d | NA |
| Cuprofix Ultra 40 disperss | M | 1.2-2.5 lb. | x | 1.2-2.5 lb. | 1.2-2.5 lb. | 1.2-2.5 lb. | x | x | 12h | 20.5 lb. |
| copper sulfate | | F | x | F | F | F | x | x | 0d | NA |
| Cuproxtat FL | M | 2.5-5 pt. | x | 2.5-5 pt. | 2.5-5 pt. | 2.5-5 pt. | x | x | 48h | NA |
| tribasic copper sulfate | | F | x | F | F | F | x | x | NA | NA |
| Elevate 50WDG | 17 | x | x | x | x | x | x | x | 4h | 6 lb. |
| fenhexamid | | x | x | x | x | x | x | x | 0d | NA |
| Evito 480SC | 11 | x | x | x | x | x | 2.0-5.7 fl. oz. | x | 12h | 22.8 fl. oz. |
| fluoxastrobin | | x | x | x | x | x | E | x | 1d | 4 |
| Flint Extra | 11 | x | 2.5-3 fl. oz. | 2.5-3 fl. oz. | x | x | 2.5-3 fl. oz. | x | 12h | 18 fl. oz. |
| trifloxystrobin | | x | G | G | x | x | G | x | 0d | 6 |
| Fontelis (SC) | 7 | x | 16-24 oz. | x | x | x | 16-24 oz. | x | 12h | 72 fl. oz. |
| penthiopyrad | | x | G | x | x | x | E | x | 0d | NA |
| Inspire | 3 | x | 7 fl. oz. | x | x | x | 7 fl. oz. | x | 12h | 28 fl. oz. |
| difenoconazole | | x | u | x | x | x | u | x | 0d | 4 |
| Inspire Super | 3+9 | x | 16-20 fl. oz. | x | x | x | 16-20 fl. oz. | x | 12h | 80 fl. oz. |
| difenoconazole+cyprodinil | | x | u | x | x | x | F | x | 0d | 3 or 4 |
| Intuity (SC) | 11 | x | x | x | x | x | 6 fl. oz. | x | 12h | 12 fl. oz. |
| mandestrobin | | x | x | x | x | x | s(E) | x | 0d | 2 |

(Continued)

Table 9-2. Strawberry Diseases - Prebloom¹ (continued)

| Product And Formulation Active Ingredient | FRAC Code ² | Angular Leaf Spot | Anthraxnose Crown | Leaf Blight Phomopsis | Leaf Scorch (Diplocarpon) | Leaf Spot (Mycosphaerella) | Powdery Mildew | Red Stele (Phytophthora) | REI ³ PHI ⁴ | Max Amt ⁵ Max App ⁶ |
|--|---------------------------|-------------------|----------------------|--------------------------|------------------------------|-------------------------------|----------------------|-----------------------------|--------------------------------------|--|
| Kenja 400SC | 7 | x | 13.5-15.5 fl. oz. | x | x | x | x | x | 12h | 54 fl. oz. |
| isofetamid | | x | i | x | x | x | x | x | 0d | NA |
| Kocide 3000 | M | 0.75-1.25 lb. | x | 0.75-1.25 lb. | 0.75-1.25 lb. | 0.75-1.25 lb. | x | x | 48h | 27.3 lb. |
| copper hydroxide | | F | x | F | F | F | x | x | 0d | NA |
| Luna Flex | 7+3 | x | 12.0-13.6 fl. oz. | x | x | x | 12.0-13.6 fl. oz. | x | 12h | 13.7 fl. oz. |
| fluopyram+difenoconazole | | x | u | x | x | x | F-G | x | 0d | 2 |
| Luna Privilege | 7 | x | 3.2-6.8 fl. oz. | x | x | 3.2-6.8 fl. oz. | 3.2-6.8 fl. oz. | x | 12h | 13.7 fl. oz. |
| fluopyram | | x | G | x | x | G | G | x | NL | 2 |
| Luna Sensation (SC) | 7+11 | x | 4-7.6 fl. oz. | x | x | 4-7.6 fl. oz. | 4-7.6 fl. oz. | E | 12h | 27.1 fl. oz. |
| fluopyram + trifloxystrobin | | x | E | x | x | G | E | 4-7.6 fl. oz. | 0d | NA |
| Luna Tranquility (SC) | 7+9 | x | x | 16-27 fl. oz. | x | 16-27 fl. oz. | 16-27 fl. oz. | x | 12h | 54.7 fl. oz. |
| fluopyram + pyrimethanil | | x | x | G | x | G | E | x | 1d | NA |
| Merivon | 7+11 | x | 4-8 fl. oz. | 4-8 fl. oz. | 4-8 fl. oz. | 4-8 fl. oz. | 4-8 fl. oz. | x | 12h | 33 fl. oz. |
| pyraclostrobin + fluxapyroxad | | x | E | E | G | E | F | x | 0d | 3 |
| Mettle 125ME | 3 | x | x | 3-5 oz. | 3-5 oz. | 3-5 oz. | 3-5 oz. | x | 12h | 20 fl. oz. |
| tetraconazole | | x | x | E | G | G | E | x | 0d | 4 |
| Miravis Prime | 7+12 | x | 9.1-13.4 oz. | 9.1-13.4 oz. | x | x | 9.1-13.4 oz. | x | 12h | 27.2 oz. |
| pydiflumetofen + fludioxonil | | x | E | G | x | x | G | x | 0 d | 4 |
| Nordox 75G | M | x | x | 1.25-2.5 lb. | 1.25-2.5 lb. | 1.25-2.5 lb. | x | x | 12h | NA |
| cuprous oxide | | x | x | F | F | F | x | x | NA | NA |
| Orondis Gold 200 | 49+4 | x | x | x | x | x | x | 20-62 fl. oz. | 48h | 124 fl. oz. |
| oxathiapiprolin + mefenoxam | | x | x | x | x | x | x | E | 28d | 2 |
| PhD | 19 | x | 6.2 oz. | x | x | x | 6.2 oz. | x | 4h | NA |
| polyoxin D | | x | G | x | x | x | G | x | 0d | NA |
| Prophyt | 33 | x | x | x | x | x | x | 2 pt./100 gal. | 4h | 4 pt. |
| phosphorous acid | | x | x | x | x | x | x | G | NA | varies |
| Phostrol | 33 | x | x | x | x | x | x | G | 4 h | NA |
| phosphorous acid | | x | x | x | x | x | x | 2.5-5 pt./100 gal. | NA | NA |

(Continued)

Table 9-2. Strawberry Diseases - Prebloom¹ (continued)

| Product And Formulation Active Ingredient | FRAC Code ² | Angular Leaf Spot | Anthraxnose Crown | Leaf Blight Phomopsis | Leaf Scorch (Diplocarpon) | Leaf Spot (Mycosphaerella) | Powdery Mildew | Red Stele (Phytophthora) | REI ³ PHI ⁴ | Max Amt ⁵ Max App ⁶ |
|--|---------------------------|-------------------|----------------------|--------------------------|------------------------------|-------------------------------|----------------------|-----------------------------|--------------------------------------|--|
| Protocol (L) | 1+3 | x | 1.3 pt. | x | x | 1.3 pt. | 1.3 pt. | x | 24h | 5.3 pt. |
| thiophanate-methyl + propiconazole | | x | G[r] | x | x | G[r] | G[r] | x | 1d | NA |
| Pristine | 11+7 | x | 18.5-23 oz. | 18.5-23 oz. | x | 18.5-23 oz. | 18.5-23 oz. | 18.5-23 oz. | 12h | 115 oz. |
| pyraclostrobin + boscalid | | x | G | G | x | G | E | E | 0d | 5 |
| ProBlad Verde | M12 | x | x | x | x | x | 20.5-24.4 fl. oz. | x | 4 h | NA |
| Banda de Lupinus alba doce | | x | x | x | x | x | F | x | 1d | 5 |
| Procure 480SC | 3 | x | x | x | x | x | 4-8 fl. oz. | x | 12h | 32 fl. oz. |
| triflumizole | | x | x | x | x | x | G | x | 1d | NA |
| Quadris Top (SC) | 3+11 | x | 12-14 fl. oz. | x | x | 12-14 fl. oz. | 12-14 fl. oz. | x | 12h | 56 fl. oz. |
| difenoconazole + azoxystrobin | | x | G | x | x | G | G | x | 0d | 4 |
| Quilt Xcel (SE) | 11+3 | x | 14 fl. oz. | x | x | 14 fl. oz. | 15 fl. oz. | x | 12h | 56 fl. oz. |
| azoxystrobin + propiconazole | | x | G | x | x | G | E | x | 0d | 4 |
| Quintec 28F | 13 | x | x | x | x | x | 4-6 fl. oz. | x | 24h | 24 fl. oz. |
| quinoxyfen | | x | x | x | x | x | E | x | 1d | 4 |
| Rally 40WSP | 3 | x | x | 2.5-5 oz. | x | 2.5-5 oz. | 2.5-5 oz. | x | 24h | 30 oz. |
| myclobutanil | | x | x | F | x | G | E | x | 0d | NA |
| Ridomil Gold SL | 4 | x | x | x | x | x | x | E[r] | 48h | 1.5 lb. |
| mefenoxam | | x | x | x | x | x | x | 1 pt. | 0d | 3 |
| Rovral 4F | 2 | x | 2 pt. | 2 pt. | x | 2 pt. | x | x | 24h | NA |
| iprodione | | x | u | u | x | u | x | x | 0d | 1 |
| Switch 62.5WG | 9+12 | x | 11-14 oz. | x | x | 11-14 oz. | x | x | 12h | 56 oz. |
| cyprodinil + fludioxonil | | x | G | x | x | F | x | x | 0d | NA |
| Thiram SC | M | x | x | x | x | 2.0-2.5 qt. | x | x | 24h | 29.7 qt. |
| thiram | | x | x | x | x | i | x | x | 1d | 12 |
| Tilt (EC) | 3 | x | x | x | x | 4 fl. oz. | 5 fl. oz. | x | 12h | 16 fl. oz. |
| propiconazole | | x | x | x | x | G | G | x | 0d | NA |
| Topguard EQ (SC) | 3+11 | x | 5-8 oz. | x | x | x | 5-8 oz. | 5-8 oz. | 12h | 32 fl. oz. |
| azoxystrobin + flutriafol | | x | G | x | x | x | E | u | 0d | 4 |
| Topsin-M WSB | 1 | x | 0.75-1 lb. | 0.75-1 lb. | x | 0.75-1 lb. | 0.75-1 lb. | x | 24h | 4 lb. |
| thiophanate-methyl | | x | G | i | x | G | i[r] | x | 1d | NA |
| Torino (SC) | U6 | x | x | x | x | x | 3.4 fl. oz. | x | 4h | 7.2 oz. |
| cyflufenamid | | x | x | x | x | x | E | x | 3d | 2 |

Footnotes: See how to read the spray schedule tables section at the beginning of the chapter.

Strawberry Prebloom - Insects

- Do not apply insecticides during bloom to protect bees and other pollinators.
- Watch for clipper when flower buds start coming out of the crown and when temperatures approach 65°F. Treat if the number of clipped buds per row-yard is 3 or more primary buds, or 30 or more secondary or tertiary buds. Infestations begin at field edge so border spray is often sufficient.
- For **Eastern flower thrips**: Sample when first blossom buds begin to open. Threshold is 2-10 thrips per blossom. Treat before widespread bloom draws pollinators.
- For **spittle bug, tarnished plant bug**: If a problem, apply when buds first become visible, and make a second application just before the first bloom opens.
- Admire Pro cannot be applied within 10 days prior to bloom or when bees are foraging.
- Diazinon AG600 allows for one foliar and one soil application per year.

Table 9-3. Strawberry Insect - Prebloom¹

| Product And Formulation Active Ingredient | IRAC Code ² | Clipper | Cyclamen Mite | Eastern Flower Thrips | Spider Mite | Spittlebug | Tarnished Plant Bug | Thrips | REI ³ PHI ⁴ | Max Amt ⁵ Max App ⁶ |
|--|------------------------|-----------------|-----------------|-----------------------|-------------------------|---------------|---------------------|-------------|--------------------------------------|--|
| Acramite 50WS | 20D | x | x | x | 0.75-1 lb. | x | x | x | 12h | NA |
| bifenazate | | x | x | x | E | x | x | x | 1d | 1 |
| Actara (25WDG) | 4A | x | x | x | x | x | 4 oz. | x | 12h | 12 oz. |
| thiamethoxam | | x | x | x | x | x | s | x | 3d | NA |
| Admire Pro (4.6F) | 4A | x | x | x | x | 1.3 fl. oz. | x | x | 12h | 3.9/14 fl. oz. |
| imidacloprid | | x | x | x | x | G | x | x | 7 or 14d | NA |
| Agri-Mek SC (0.7SC) (RUP) | 6 | x | 3.5 fl. oz. | x | 3.5 fl. oz. | x | x | x | 12h | 14 fl. oz. |
| abamectin | | x | s | x | E | x | x | x | 3d | NA |
| Apta (1.34SC) | 21A | x | x | 27 fl. oz. | x | x | 27 fl. oz. | 27 fl. oz. | 12h | 81 fl. oz. |
| tolfenpyrad | | x | x | x | u | x | u | u | 1d | 3 |
| Assail 30SG | 4A | x | x | 4-6.9 oz. | x | 1.9-6.9 oz. | 1.9-6.9 oz. | 4-6.9 oz. | 12h | 13.8 oz. |
| acetamiprid | | x | x | G | x | G | G | E | 1d | 2 |
| Beleaf 50SG | 29 | x | x | x | x | x | 2.8 oz. | x | 12h | 8.4 oz. |
| flonicamid | | x | x | x | x | x | E | x | 0d | 2 |
| Brigade eVo | 3A | 2.56-12.8 oz. | x | x | 6.4-12.8 oz. | 2.56-12.8 oz. | 2.56-12.8 oz. | x | 12h | 32 oz. |
| bifenthrin | | E | x | x | F | E | E | x | 0d | 3 |
| Danitol 2.4EC (RUP) | 3A | 16-21.3 fl. oz. | 16-21.3 fl. oz. | x | 16-21.3 fl. oz. | 10.6 fl. oz. | 10.6 fl. oz. | x | 24h | See label |
| fenpropathrin | | E | u | x | F | u | u | x | 3d | See label |
| Diazinon AG600 WBC (RUP) | 1B | x | 25.5 oz. | x | 12.75 fl. oz. /100 gal. | x | x | x | 3d | 25.5 fl. oz. |
| diazinon | | x | G | x | F | x | x | x | 5d | 1 |
| Dibrom 8E (RUP) | 1B | x | x | 1 pt. | 1 pt. | 1 pt. | 1 pt. | x | 2d | 5 pt. |
| naled | | x | x | u | u | u | u | x | 1d | 5 |
| Entrust SC (2SC) | 5 | x | x | 4-6 fl. oz. | x | x | x | 4-6 fl. oz. | 4h | 18 fl. oz. |
| spinosad | | x | x | G | x | x | x | x | 1d | 3 |

(Continued)

Table 9-3. Strawberry Insect - Prebloom¹ (continued)

| Product And Formulation Active Ingredient | IRAC Code ² | Clipper | Cyclamen Mite | Eastern Flower Thrips | Spider Mite | Spittlebug | Tarnished Plant Bug | Thrips | REI ³ PHI ⁴ | Max Amt ⁵ Max App ⁶ |
|---|------------------------|---------|---------------|-----------------------|---------------|-------------|---------------------|----------|-----------------------------------|---|
| Exirel (0.83SE) | 28 | x | x | 13.5-20.5 fl. oz. | x | x | x | x | 12h | 61.5 fl. oz. |
| cyantraniliprole | | x | x | s | x | x | x | x | 1d | NA |
| Grandevo WDG | UN | x | 2-3 lb. | 2-3 lb. | 2-3 lb. | x | 2-3 lb. | x | 4h | NA |
| <i>Chromobacterium subtsugae</i> | | x | u | u | u | x | u | x | 0d | NA |
| Kanemite 15SC | 20B | x | x | x | 21-31 fl. oz. | x | x | x | 12h | 62 fl. oz. |
| acequinocyl | | x | x | x | E | x | x | x | 1d | 2 |
| Malathion 5EC | 1B | x | x | 1.5-3.2 pt. | 1.5-3.2 pt. | 1.5-3.2 pt. | 1.5-3.2 pt. | x | 12h | 12.8 pt. |
| malathion | | x | x | s | u | u | u | x | 3d | 4 |
| Nealta (1.67SC) | 25 | x | x | x | 13.7 fl. oz. | x | x | x | 12h | 27.4 fl. oz. |
| cyflumetofen | | x | x | x | G | x | x | x | 1d | 2 |
| Nexter (75WP) | 21 | x | x | x | 4.4-10.6 oz. | x | x | x | 12h | 21.34 oz. |
| pyridaben | | x | x | x | G | x | x | x | 1d | 2 |
| Oberon 2SC | 23 | x | x | x | 16 fl. oz. | x | x | x | 12h | 16 fl. oz. |
| spiromesifen | | x | x | x | G | x | x | x | 3d | 1 |
| Portal XLO (0.4EC) | 21A | x | 2 pt. | x | 2 pt. | x | x | x | 12h | 4 pt. |
| fenpyroximate | | x | G | x | E | x | x | x | 1d | 2 |
| Radiant SC (1SC) | 5 | x | x | 6-10 fl. oz. | x | x | x | 6-10 oz. | 4h | 30 fl. oz. |
| spinetoram | | x | x | G | x | x | x | x | 1d | 3 |
| Rimon 0.83EC | 15 | x | x | 6-12 fl. oz. | x | x | 9-12 fl. oz. | x | 12h | 36 fl. oz. |
| novaluron | | x | x | u | x | x | E | x | 1d | NA |
| Savey 50DF | 10A | x | x | x | 6 oz. | x | x | x | 12h | 6 oz. |
| hexythiazox | | x | x | x | E | x | x | x | 3d | 1 |
| Sevin XLR Plus (4F) | 1A | 1-2 qt. | x | x | x | 1-2 qt. | 1.5-2 qt. | x | 12h | 10 qt. |
| carbaryl | | G | x | x | x | x | G | x | 7d | 5 |
| Transform WG | 4C | x | x | 2.2 oz. | x | x | 1.5-2.2 oz. | x | 24h | 8.5 oz. |
| sulfoxaflor | | x | x | s | x | x | u | x | 1d | 4 |
| Vendex 50WP (RUP) | 12B | x | x | x | 1.5-2 lb. | x | x | x | 2d | 4 lb. |
| fenbutatin-oxide | | x | x | x | G | x | x | x | 1d | 2 |
| Zeal (72WP) | 10B | x | x | x | 2-3 oz. | x | x | x | 12h | 3 oz. |
| etoxazole | | x | x | x | E | x | x | x | 1d | 1 |

Footnotes: See how to read the spray schedule tables section at the beginning of the chapter.

Strawberry Prebloom - Diseases

Apply from 5-10% bloom until flowers have finished blooming.

Disease management notes

- Carefully examine FRAC codes to confirm that fungicide rotations use different FRAC classes. For example, Abound, Cabrio, Pristine, Luna Sensation, Merivon and one component of Quilt Xcel all have the same FRAC code (FRAC 11) and cannot be alternated with each other as a fungicide resistance management strategy.
- Anthraco**se can be severe on both green and ripe (red) strawberry fruit. The disease is favored by moderate to warm temperatures accompanied by rainfall before and during harvest. A preventative spray program (7-10 days) with Captan beginning at 10% bloom is recommended. During periods of high disease pressure the addition of FRAC 11 or FRAC 3+11 (i.e., Quadris Top or Quilt Xcel) fungicides.
- Captan and copper both pose a risk of phytotoxicity at bloom. Take care to avoid tank mixing, particularly with other products that have EC or SC formulations.
- Fungicide resistance to the FRAC 7 and 11 fungicides has been reported in *Botrytis* in many strawberry producing states but has only confirmed in Ohio (FRAC 11 only) the Midwest. Resistance to FRAC 1 has been a longstanding issue.

Table 9-4. Strawberry Diseases - Bloom¹

| Product And Formulation Active Ingredient | FRAC Code ² | Anthraco nose Crown | Anthraco nose Fruit Rot | Gray Mold (<i>Botrytis</i>) | Leaf Blight Phomopsis | Leaf Scorch (<i>Diplocarpon</i>) | Leaf Spot (<i>Mycosphaerella</i>) | Powdery Mildew | REI ³ PHI ⁴ | Max Amt ⁵ Max App ⁶ |
|--|---------------------------|---------------------------|-------------------------------|----------------------------------|--------------------------|---------------------------------------|--|--------------------|--------------------------------------|--|
| Abound (SC) | 11 | 5-8 fl. oz. | x | 6-15.5 fl. oz. | x | x | x | 6-15.5 fl. oz. | 4h | 61.5 fl. oz. |
| azoxystrobin | | G[r] | x | s | x | x | x | G | 0d | NA |
| Aftershock/Evito 480SC | 11 | 2-5.7 fl. oz. | 2-5.7 fl. oz. | x | x | x | x | 2-5.7 fl. oz. | 12h | 22.8 fl. oz. |
| fluoxastrobin | | G | G | x | x | x | x | G | 1d | NA |
| Cabrio EG (20EG) | 11 | 12-14 oz. | 12-14 oz. | 12-14 oz. | x | x | 12-14 oz. | 12-14 oz. | 12h | 70 oz. |
| pyraclostrobin | | E | G | G | x | x | G | E | 0d | NA |
| Captan (80WDG) | M | 1.8-3.75 lb. | 1.8-3.75 lb. | 1.8-3.75 lb. | 1.8-3.75 lb. | 1.8-3.75 lb. | 1.8-3.75 lb. | x | 24h | 30 lb. |
| captan | | E | G | G | G | G | G | x | 0d | NA |
| Elevate 50WDG | 17 | x | x | 1.5 lb. | x | x | x | x | 4h | 6 lb. |
| fenhexamid | | x | x | E[r] | x | x | x | x | 0d | NA |
| Evito 480SC | 11 | x | 2.0-5.7 fl. oz. | x | x | x | x | 2.0-5.7 fl. oz. | 12h | 22.8 fl. oz. |
| fluoxastrobin | | x | E | s | x | x | x | E | 0d | 4 |
| Flint Extra | 11 | 2.5-3 fl. oz. | 2.5-3 fl. oz. | 2.5-3 fl. oz. | 2.5-3 fl. oz. | x | x | 2.5-3 fl. oz. | 12h | 18 fl. oz. |
| trifloxystrobin | | G | E | G | G | x | x | G | 0d | NA |
| Fontelis (SC) | 7 | 16-24 oz. | 16-24 oz. | 16-24 oz. | x | x | x | 16-24 oz. | 12h | 72 fl. oz. |
| penthiopyrad | | G | E | E | x | x | x | E | 0d | NA |
| Inspire | 3 | 7 fl. oz. | 7 fl. oz. | x | x | x | x | 7 fl. oz. | 12h | 28 fl. oz. |
| difenoconazole | | u | G | x | x | x | x | u | 0d | 4 |

(Continued)

Table 9-4. Strawberry Diseases - Bloom¹ (continued)

| Product And Formulation Active Ingredient | FRAC Code ² | Anthraco- nose Crown | Anthraco- nose Fruit Rot | Gray Mold (Botrytis) | Leaf Blight Phomopsis | Leaf Scorch (Diplocarpon) | Leaf Spot (Mycosphaerella) | Powdery Mildew | REI ³ PHI ⁴ | Max Amt ⁵ Max App ⁶ |
|--|---------------------------|----------------------------|--------------------------------|-------------------------|--------------------------|------------------------------|-------------------------------|--------------------------|--------------------------------------|--|
| Inspire Super | 3+9 | 16-20 fl. oz. | 16-20 fl. oz. | 16-20 fl. oz. | x | x | x | 16-20 fl. oz. | 12h | 13.7 fl. oz. |
| difenoconazole+cyrodinil | | u | G | F | x | x | x | F | 0d | 2 |
| Intuity (SC) | 11 | x | x | 6 fl. oz. | x | x | x | 6 fl. oz. | 12h | 12 fl. oz. |
| mandestrobin | | x | x | E | x | x | x | s-E | 0d | 2 |
| Kenja 400SC | 7 | 13.5-15.5 fl. oz. | 13.5-15.5 fl. oz. | 13.5-15.5 fl. oz. | x | x | x | x | 12h | 54 fl. oz. |
| isofetamid | | i | i | E | x | x | x | x | 0d | NA |
| Luna Flex | 7+3 | 12-13.6 fl. oz. | 12-13.6 fl. oz. | 12-13.6 fl. oz. | x | x | 12-13.6 fl. oz. | 12-13.6 fl. oz. | 12 h | 13.7 fl. oz. |
| fluopyram+difenoconazole | | u | u | F | x | x | u | F-G | 0d | 2 |
| Luna Sensation (SC) | 7+11 | 4-7.6 fl. oz. | 4-7.6 fl. oz. | 4-7.6 fl. oz. | x | x | 4-7.6 fl. oz. | 4-7.6 fl. oz. | 12h | 27.1 fl. oz. |
| fluopyram + trifloxystrobin | | E | x | G | x | x | G | E | 0d | NA |
| Luna Tranquility (SC) | 7+9 | x | x | 16-27 fl. oz. | 16-27 fl. oz. | x | 16-27 fl. oz. | 16-27 fl. oz. | 12h | 54.7 fl. oz. |
| fluopyram + pyrimethanil | | x | x | E | G | x | G | E | 1d | NA |
| Merivon | 7+11 | 4-8 fl. oz. | 4-8 fl. oz. | 4-8 fl. oz. | 4-8 fl. oz. | 4-8 fl. oz. | 4-8 fl. oz. | 4-8 fl. oz. | 12h | 33 fl. oz. |
| pyraclostrobin + fluxapyroxad | | E | E | E | E | G | E | F | 0d | 3 |
| Mettle 125ME | 3 | x | x | x | 3-5 oz. | 3-5 oz. | 3-5 oz. | 3-5 oz. | 12h | 20 fl. oz. |
| tetraconazole | | x | x | x | G | G | G | E | 0d | 4 |
| Miravis Prime | 7+12 | 9.1-13.4 oz. | 9.1-13.4 oz. | 9.1-13.4 oz. | 9.1-13.4 oz. | x | x | 9.1-13.4 oz. | 12 h | 27.2 oz. |
| pydiflumetofen+fludioxonil | | E | E | E | G | x | x | G | 0 d | 4 |
| Nordox 75G | M | x | x | x | 3-5 lb. | 3-5 lb. | 3-5 lb. | x | 12h | NA |
| cuprous oxide | | x | x | x | F | F | F | x | NA | NA |
| PhD | 19 | 6.2 oz. | 6.2 oz. | 6.2 oz. | x | x | x | 6.2 oz. | 4h | NA |
| polyoxin D | | G | G | E | x | x | x | G | 0d | NA |
| Protocol (L) | 1+3 | 1.3 pt. | 1.3 pt. | 1.3 pt. | x | x | 1.3 pt. | 1.3 pt. | 24h | 5.3 pt. |
| thiophanate-methyl + propiconazole | | G[r] | G[r] | G[r] | x | x | G[r] | G[r] | 1d | NA |
| Pristine | 11+7 | 18.5-23 oz. | 18.5-23 oz. | 18.5-23 oz. | 18.5-23 oz. | x | 18.5-23 oz. | 18.5-23 oz. | 12h | 115 oz. |
| pyraclostrobin + boscalid | | G | E | E | G | x | G | E | 0d | 5 |
| ProBlad Verde | M12 | x | x | 24.4-36.6 fl. oz. | x | x | x | 20.5- 24.4 fl. oz. | 4 h | NA |
| Banda de Lupinus alba doce | | x | x | F | x | x | x | F | 1d | 5 |

(Continued)

Table 9-4. Strawberry Diseases - Bloom¹ (continued)

| Product And Formulation Active Ingredient | FRAC Code ² | Anthracnose Crown | Anthracnose Fruit Rot | Gray Mold (Botrytis) | Leaf Blight Phomopsis | Leaf Scorch (Diplocarpon) | Leaf Spot (Mycosphaerella) | Powdery Mildew | REI ³ PHI ⁴ | Max Amt ⁵ Max App ⁶ |
|--|---------------------------|----------------------|--------------------------|-------------------------|--------------------------|------------------------------|-------------------------------|------------------|--------------------------------------|--|
| Procure 480SC | 3 | x | x | x | x | x | x | 4-8 fl. oz. | 12h | 32 fl. oz. |
| triflumizole | | x | x | x | x | x | x | G | 1d | NA |
| Quadris Top (SC) | 3+11 | 12-14 fl. oz. | 12-14 fl. oz. | 12-14 fl. oz. | x | x | 12-14 fl. oz. | 12-14 fl. oz. | 12h | 56 fl. oz. |
| difenoconazole + azoxystrobin | | G | E | G | x | x | G | G | 0d | 4 |
| Quilt Xcel (SE) | 11+3 | 14 fl. oz. | 14 fl. oz. | 14 fl. oz. | x | x | 14 fl. oz. | 14 fl. oz. | 12h | 56 fl. oz. |
| azoxystrobin + propiconazole | | G | G | F | x | x | x | E | 0d | 4 |
| Quintec 2.08F | 13 | x | x | x | x | x | x | 4-6 fl. oz. | 24h | 24 fl. oz. |
| quinoxifen | | x | x | x | x | x | x | E | 1d | 4 |
| Rally 40WSP | 3 | x | x | x | 2.5-5 oz. | x | 2.5-5 oz. | 2.5-5 oz. | 24h | 30 oz. |
| myclobutanil | | x | x | x | F | x | G | E | 0d | NA |
| Rovral 4F | 2 | x | x | 2 pt. | x | x | 2 pt. | x | 24h | 1 |
| iprodione | | x | x | G[r] | x | x | G | x | 0d | N/A |
| Scala SC | 9 | x | x | 18 fl. oz. | x | x | x | x | 12h | 54 fl. oz. |
| pyrimethanil | | x | x | E[r] | x | x | x | x | 1d | NA |
| Switch 62.5WG | 9+12 | 11-14 oz. | 11-14 oz. | 11-14 oz. | x | x | 11-14 oz. | x | 12h | 56 oz. |
| cyprodinil + fludioxonil | | G | E | E | x | x | F | x | 0d | NA |
| Thiram SC | M | x | 2.0-2.2 qt. | 1.5-2.5 qt. | x | x | x | x | 24h | 29.7 qt. |
| thiram | | F | G | G | x | x | x | x | 1d | 12 |
| Tilt (EC) | 3 | x | x | x | x | x | 4 fl. oz. | 4 fl. oz. | 12h | 16 fl. oz. |
| propiconazole | | x | x | x | x | x | G | G | 0d | NA |
| Topguard EQ (SC) | 3+11 | 5-8 oz. | 5-8 oz. | x | x | x | x | 5-8 oz. | 12h | 32 fl. oz. |
| azoxystrobin + flutriafol | | G | G | x | x | x | x | E | 0d | 4 |
| Topsin-M WSB | 1 | 0.75-1 lb. | 0.75-1 lb. | 0.75-1 lb. | 0.75-1 lb. | x | 0.75-1 lb. | 0.75-1 lb. | 24h | 4 lb. |
| thiophanate-methyl | | G | E | G[r] | i | x | G | G-P[r] | 1d | NA |
| Torino (SC) | U6 | x | x | x | x | x | x | 3.4 fl. oz. | 4h | 6.8 oz. |
| cyflufenamid | | x | x | x | x | x | x | E | 3d | 2 |

Footnotes: See how to read the spray schedule tables section at the beginning of the chapter.

Strawberry Bloom - Insects

- Do not apply insecticides during bloom to protect bees and other pollinators.

Strawberry Postbloom To Harvest - Diseases

Apply every 7-10 days as needed. Be sure to check PHIs.

- Abound, Cabrio, and Pristine also provide protection against leather rot when applied in a protectant program.
- Leather rot (*Phytophthora cactorum*) can be managed using the same fungicides used for anthracnose. Phosphorous acids like ProPhyt and Phostrol also provide good control.

Table 9-5. Strawberry Diseases - Postbloom Through Harvest¹

| Product And Formulation Active Ingredient | FRAC Code ² | Anthracnose Crown Rot | Anthracnose Fruit Rot | Gray Mold (Botrytis) | Leaf Blight Phomopsis | Leaf Scorch (Diplocarpon) | Leaf Spot (Mycosphaerella) | Powdery Mildew | Leather Rot (Phytophthora) | REI ³ PHI ⁴ | Max Amt ⁵ Max App ⁶ |
|--|---------------------------|--------------------------|--------------------------|-------------------------|--------------------------|------------------------------|-------------------------------|----------------|-------------------------------|--------------------------------------|--|
| Abound (SC) | 11 | 5-8 fl. oz. | x | 6-15.5 fl. oz. | x | x | x | 6-15.5 fl. oz. | 6-15.5 fl. oz. | 4h | 61.5 fl. oz. |
| azoxystrobin | | G[r] | x | s-G[r] | x | x | x | G | E | 0d | NA |
| Aftershock/Evito 480SC | 11 | 2-5.7 fl. oz. | 2-5.7 fl. oz. | x | x | x | x | 2-5.7 fl. oz. | x | 12h | 22.8 fl. oz. |
| fluoxastrobin | | G | G | x | x | x | x | G | x | 1d | NA |
| Aliette WDG | P07 (33) | x | x | x | x | x | x | x | 2.5-5 lb. | 12h | 30 lb. |
| aluminum tris | | x | x | x | x | x | x | x | E | 1d | NA |
| Cabrio EG (20EG) | 11 | 12-14 oz. | 12-14 oz. | 12-14 oz. | x | x | 12-14 oz. | 12-14 oz. | 12-14 oz. | 12h | 70 oz. |
| pyraclostrobin | | E | G | G | x | x | G | E | E | 0d | NA |
| Captan (80WDG) | M | 1.8-3.75 lb. | 1.87-3.75 lb. | 1.8-3.75 lb. | 1.87-3.75 lb. | 1.8-3.75 lb. | 1.8-3.75 lb. | x | x | 24h | 30 lb. |
| captan | | E | G | G | G | G | G | x | x | 0d | NA |
| Elevate 50WDG | 17 | x | x | 1.5 lb. | x | x | x | x | x | 4h | 6 lb. |
| fenhexamid | | x | x | E[r] | x | x | x | x | x | 0d | NA |
| Evito 480SC | 11 | x | 2-5.7 fl. oz. | x | x | x | x | 2-5.7 fl. oz. | x | 12h | 22.8 fl. oz. |
| fluoxastrobin | | x | E | s | x | x | x | E | x | 1d | 4 |
| Flint Extra | 11 | 2.5-3 fl. oz. | 2.5-3 fl. oz. | 2.5-3 fl. oz. | 2.5-3 fl. oz. | x | x | 2.5-3 fl. oz. | x | 12h | 18 fl. oz. |
| trifloxystrobin | | G | E | G | G | x | x | G | x | 0d | NA |
| Fontelis (SC) | 7 | 16-24 oz. | 16-24 oz. | 16-24 oz. | x | x | x | 16-24 oz. | x | 12h | 72 fl. oz. |
| penthiopyrad | | G | E | E | x | x | x | E | x | 0d | NA |
| Inspire | 3 | 7 fl. oz. | 7 fl. oz. | x | x | x | x | 7 fl. oz. | x | 12h | 28 fl. oz. |
| difenoconazole | | u | G | x | x | x | x | u | x | 0h | 4 |

(Continued)

Table 9-5. Strawberry Diseases - Postbloom Through Harvest¹ (continued)

| Product And Formulation Active Ingredient | FRAC Code ² | Anthraxnose Crown Rot | Anthraxnose Fruit Rot | Gray Mold (Botrytis) | Leaf Blight Phomopsis | Leaf Scorch (Diplocarpon) | Leaf Spot (Mycosphaerella) | Powdery Mildew | Leather Rot (Phytophthora) | REI ³ PHI ⁴ | Max Amt ⁵ Max App ⁶ |
|--|---------------------------|--------------------------|--------------------------|-------------------------|--------------------------|------------------------------|-------------------------------|--------------------------|-------------------------------|--------------------------------------|--|
| Inspire Super | 3+9 | 16-20 fl. oz. | 16-20 fl. oz. | 16-20 fl. oz. | x | x | x | 16-20 fl. oz. | x | 12h | 13.7 fl. oz. |
| difenoconazole+cyprodinil | | u | G | F | x | x | x | F | x | 0d | 2 |
| Intuity (SC) | 11 | x | x | 6 fl. oz. | x | x | x | 6 fl. oz. | x | 12h | 12 fl. oz. |
| mandestrobin | | x | x | E | x | x | x | s-E | x | 0d | 2 |
| Kenja 400SC | 7 | 13.5-15.5 fl. oz. | 13.5-15.5 fl. oz. | 13.5-15.5 fl. oz. | x | x | x | x | x | 12h | 54 fl. oz. |
| isofetamid | | i | i | E | x | x | x | x | x | 0d | NA |
| Luna Flex | 7+3 | 12.0-13.6 fl. oz. | 12.0-13.6 fl. oz. | 12.0-13.6 fl. oz. | x | x | 12.0- 13.6 fl. oz. | 12.0- 13.6 fl. oz. | x | 12 h | 13.7 fl. oz. |
| fluopyram + difeno- conazole | | u | u | F | x | x | u | F-G | x | 0d | 2 |
| Luna Sensation (SC) | 7+11 | 4-7.6 fl. oz. | 4-7.6 fl. oz. | 4-7.6 fl. oz. | x | x | 4-7.6 fl. oz. | 4-7.6 fl. oz. | 4-7.6 fl. oz. | 12h | 271 fl. oz. |
| fluopyram + trifloxystrobin | | E | x | G | x | x | G | E | E | 0d | NA |
| Luna Tranquility (SC) | 7+9 | x | x | 16-27 fl. oz. | 16-27 fl. oz. | x | 16-27 fl. oz. | 16-27 fl. oz. | x | 12h | 54.7 fl. oz. |
| fluopyram + pyrimethanil | | x | x | E | G | x | G | E | x | 1d | NA |
| Merivon | 7+11 | 4-8 fl. oz. | 4-8 fl. oz. | 4-8 fl. oz. | 4-8 fl. oz. | 4-8 fl. oz. | 4-8 fl. oz. | 4-8 fl. oz. | x | 12h | 33 fl. oz. |
| pyraclostrobin + fluxopy- roxad | | E | E | E | E | G | E | F | x | 0d | 3 |
| Mettle 125ME | 3 | x | x | x | 3-5 oz. | 3-5 oz. | 3-5 oz. | 3-5 oz. | x | 12h | 20 fl. oz. |
| tetraconazole | | x | x | x | G | G | G | E | x | 0d | 4 |
| Miravis Prime | 7+12 | 9.1-13.4 oz. | 9.1-13.4 oz. | 9.1-13.4 oz. | 9.1-13.4 oz. | x | x | 9.1-13.4 oz. | x | 12 h | 27.2 oz. |
| pydiflumetofen+fludioxonil | | E | E | E | G | x | x | G | x | 0 d | 4 |
| Nordox 75G | M | x | x | x | 1.25-2.5 lb. | 1.25-2.5 lb. | 1.25- 2.5 lb. | x | x | 12h | NA |
| cuprous oxide | | x | x | x | F | F | F | x | x | NA | NA |
| Orondis Gold | 49+4 | x | x | x | x | x | x | x | 20-62 fl. oz.** | 48 h | 124.0 fl. oz. |
| oxathiapiprolin+ mefenox- am | | x | x | x | x | x | x | x | E | 28d | 2 |
| PhD | 19 | 6.2 oz. | 6.2 oz. | 6.2 oz. | x | x | x | 6.2 oz. | s | 4h | NA |
| polyoxin D | | G | G | E | x | x | x | G | x | 0d | NA |
| Prophyt | P07 (33) | x | x | x | x | x | x | x | 2 pt./ 100 gal. | 4h | varies |
| phosphorous acid | | x | x | x | x | x | x | x | G | NA | 4 |

(Continued)

Table 9-5. Strawberry Diseases - Postbloom Through Harvest¹ (continued)

| Product And Formulation Active Ingredient | FRAC Code ² | Anthraxnose Crown Rot | Anthraxnose Fruit Rot | Gray Mold (Botrytis) | Leaf Blight Phomopsis | Leaf Scorch (Diplocarpon) | Leaf Spot (Mycosphaerella) | Powdery Mildew | Leather Rot (Phytophthora) | REI ³ PHI ⁴ | Max Amt ⁵ Max App ⁶ |
|--|---------------------------|--------------------------|--------------------------|-------------------------|--------------------------|------------------------------|-------------------------------|------------------|-------------------------------|--------------------------------------|--|
| Phostrol | P07 (33) | x | x | x | x | x | x | x | 2.5 - 5.0 pt./100 gal. | 4 h | NA |
| phosphorous acid | | x | x | x | x | x | x | x | G | NA | NA |
| Protocol (L) | 1+3 | 1.3 pt. | 1.3 pt. | 1.3 pt. | x | x | 1.3 pt. | 1.3 pt. | x | 24h | 5.3 pt. |
| thiophanate-methyl + propiconazole | | G[r] | G[r] | G[r] | x | x | G[r] | G[r] | x | 1d | NA |
| Pristine | 11+7 | 18.5-23 oz. | 18.5-23 oz. | 18.5-23 oz. | 18.5-23 oz. | x | 18.5-23 oz. | 18.5-23 oz. | 18.5-23 oz. | 12h | 115 oz. |
| pyraclostrobin + boscalid | | G | E | E | G | x | G | E | E | 0d | 5 |
| Procure 480SC | 3 | x | x | x | x | x | x | 4-8 fl. oz. | x | 12h | 32 fl. oz. |
| triflumizole | | x | x | x | x | x | x | G | x | 1d | NA |
| Quadris Top (SC) | 3+11 | 12-14 fl. oz. | 12-14 fl. oz. | 12-14 fl. oz. | x | x | 12-14 fl. oz. | 12-14 fl. oz. | x | 12h | 56 fl. oz. |
| difenoconazole + azox- ystrobin | | G | E | G | x | x | G | G | x | 0d | 4 |
| Quilt Xcel (SE) | 11+3 | 14 fl. oz. | 14 fl. oz. | 14 fl. oz. | x | x | 14 fl. oz. | 14 fl. oz. | x | 12h | 56 fl. oz. |
| azoxystrobin + propi- conazole | | G | G | F | x | x | x | E | x | 0d | 4 |
| Quintec 2.08F | 13 | x | x | x | x | x | x | 4-6 fl. oz. | x | 24h | 24 fl. oz. |
| quinoxifen | | x | x | x | x | x | x | E | x | 1d | 4 |
| Rally 40WSP | 3 | x | x | x | 2.5-5 oz. | x | 2.5-5 oz. | 2.5-5 oz. | x | 24h | 30 oz. |
| myclobutanil | | x | x | x | F | x | G | E | x | 0d | NA |
| Rally 40WSP | 4 | x | x | x | 2.5-5 oz. | x | 2.5-5 oz. | 2.5-5 oz. | x | 24h | 30 oz. |
| myclobutanil | | x | x | x | F | x | G | E | x | 0d | NA |
| Ridomil Gold SL | 4 | x | x | x | x | x | x | x | 1 pt. | 48h | 1 |
| mefenoxam | | x | x | x | x | x | x | x | E[r] | 0d | N/A |
| Scala SC | 9 | x | x | 18 fl. oz. | x | x | x | x | x | 12h | 54 fl. oz. |
| pyrimethanil | | x | x | E[r] | x | x | x | x | x | 1d | NA |
| Switch 62.5WG | 9+12 | 11-14 oz. | 11-14 oz. | 11-14 oz. | x | x | 11-14 oz. | x | x | 12h | 56 oz. |
| cyprodinil + fludioxonil | | G | E | E | x | x | F | x | x | 0d | NA |
| Thiram SC | M | x | 2-2.2 qt. | 1.5-2.5 qt. | x | x | x | x | x | 24h | 29.7 qt. |
| thiram | | x | G | G | x | x | x | x | x | 1d | 12 |
| Tilt (EC) | 3 | x | x | x | x | x | 4 fl. oz. | 4 fl. oz. | x | 12h | 16 fl. oz. |
| propiconazole* | | x | x | x | x | x | G | G | x | 0d | NA |

(Continued)

Table 9-5. Strawberry Diseases - Postbloom Through Harvest¹ (continued)

| Product And Formulation Active Ingredient | FRAC Code ² | Anthraco- nose Crown Rot | Anthraco- nose Fruit Rot | Gray Mold (Botrytis) | Leaf Blight Phomopsis | Leaf Scorch (Diplocarpon) | Leaf Spot (Mycosphaerella) | Powdery Mildew | Leather Rot (Phytophthora) | REI ³ PHI ⁴ | Max Amt ⁵ Max App ⁶ |
|--|---------------------------|--------------------------------|--------------------------------|-------------------------|--------------------------|------------------------------|-------------------------------|----------------|-------------------------------|--------------------------------------|--|
| Topguard EQ (SC) | 3+11 | 5-8 oz. | 5-8 oz. | x | x | x | x | 5-8 oz. | 5-8 oz. | 12h | 32 fl. oz. |
| azoxystrobin + flutriafol | | G | G | x | x | x | x | E | G | 0d | 4 |
| Topsin-M WSB6 | 1 | 0.75-1 lb | 0.75-1 lb | 0.75-1 lb | 0.75-1 lb. | x | 0.75-1 lb. | 0.75-1 lb. | x | 24h | 4 lb. |
| thiophanate-methyl | | G | E | G[r] | i | x | G | X | x | 1d | NA |
| Torino (SC) | U6 | x | x | x | x | x | x | 3.4 fl. oz. | x | 4h | 6.8 oz. |
| cyflufenamid | | x | x | x | x | x | x | E | x | 3d | 2 |

Footnotes: See how to read the spray schedule tables section at the beginning of the chapter.

Strawberry Postbloom To Harvest - Insects

- Apply every 7-10 days as needed. Be sure to check PHIs.
- Make every effort to protect bees by spraying when bees are not active, typically after sunset or early morning before 6 am.
- Strawberry root weevil larvae** damage strawberry roots. The weevils lack mobility, so infestations do not spread rapidly. Be sure that nursery stock is not infested before planting. Plow under old plantings soon after harvest, and locate new plantings 300 feet away. Adult weevils can be killed by one or more foliar sprays of Brigade. Platinum is labeled for soil application to control root weevil larvae.
- Leafhoppers** damage strawberry foliage, feed for a short time, then leave. Damaged leaves can become crinkled and turn yellow to brown at the margins. Damage is often detected after leafhoppers have left the field. Carbaryl (Sevin XLR Plus) is labeled for control of this pest. Brigade, Danitol, and Diazinon do not list leafhoppers on their labels but should also provide control.
- In June-bearing strawberry, **spotted-wing Drosophila** is not considered a major pest in the Midwest as harvest concludes before spotted-wing Drosophila populations start ramping up. In day-neutral strawberry, spotted-wing Drosophila can be a major pest and should be managed.
- No insecticides are registered for **strawberry root-worm**. This pest is best managed by post-harvest site rotation.
- Diazinon AG600 allows for one foliar and one soil application per year.

Table 9-6. Strawberry Insects - Postbloom Through Harvest¹

| Product And Formulation Active Ingredient | IRAC Code ² | Eastern Flower Thrips | Leafhopper | Leafroller | Sap Beetle | Slugs | Spider Mite | Spittlebug | Spotted wing Drosophila | Tarnished Plant Bug | REI ³ PHI ⁴ | Max Amt ⁵ Max App ⁶ |
|--|---------------------------|--------------------------|------------|------------|------------|-------|-------------|------------|----------------------------|------------------------|--------------------------------------|--|
| Acramite 50WS | 20D | x | x | x | x | x | 0.75-1 lb. | x | x | x | 12h | NA |
| bifenazate | | x | x | x | x | x | E | x | x | x | 1d | 2 |
| Actara (25WDG) | 4A | x | 4 oz. | x | x | x | x | x | x | 4 oz. | 12h | 12 oz. |
| thiamethoxam | | x | E | x | x | x | x | x | x | s | 3d | NA |

(Continued)

Table 9-6. Strawberry Insects - Postbloom Through Harvest¹ (continued)

| Product And Formulation Active Ingredient | IRAC Code ² | Eastern Flower Thrips | Leafhopper | Leafroller | Sap Beetle | Slugs | Spider Mite | Spittlebug | Spotted wing Drosophila | Tarnished Plant Bug | REI ³ PHI ⁴ | Max Amt ⁵ Max App ⁶ |
|--|------------------------|-----------------------|-----------------|------------------------|-----------------|--------|------------------------|-----------------|-------------------------|---------------------|--------------------------------------|--|
| Admire Pro (4.6F) | 4A | x | x | x | x | x | x | 1.3 fl. oz. | x | x | 12h | 3.9-14 fl. oz. |
| imidacloprid | | x | x | x | x | x | x | G | x | x | 7 or 14d | NA |
| Agri-Mek SC (0.7SC) (RUP) | 6 | x | x | x | x | x | 3.5 fl. oz. | x | x | x | 12h | 14 fl. oz. |
| abamectin | | x | x | x | x | x | E | x | x | x | 3d | NA |
| Apta (1.34SC) | 21A | 27 fl. oz. | x | x | x | x | x | x | 27 fl. oz. | 27 fl. oz. | 12h | 81 fl. oz. |
| tolfenpyrad | | u | x | x | x | x | x | x | s | u | 1d | 3 |
| Assail 30SG | 4A | 4-6.9 oz. | 4-6.9 oz. | 4-6.9 oz. | 4-6.9 oz. | x | x | 1.9-6.9 oz. | x | 4-6.9 oz. | 12h | 13.8 oz. |
| acetamiprid | | G | G | G | G | x | x | G | x | G | 1d | 2 |
| Beleaf 50SG | 29 | x | x | x | x | x | x | x | x | 2.8 oz. | 12h | 8.4 oz. |
| flonicamid | | x | x | x | x | x | x | x | x | E | 0d | 2 |
| Brigade eVo | 3A | x | x | 2.56-12.8 oz. | 2.56-12.8 oz. | x | 6.4-12.8 oz. | 2.56-12.8 oz. | x | 2.56-12.8 oz. | 12h | 32 oz. |
| bifenthrin | | x | x | G | E | x | F | E | x | E | 0d | 3 |
| Danitol 2.4EC (RUP) | 3A | x | 16-21.3 fl. oz. | 10.66-21.33 fl. oz. | 16-21.3 fl. oz. | x | 16-21.3 fl. oz. | 16-21.3 fl. oz. | 16 fl. oz. | 16-21.3 fl. oz. | 24h | see label |
| fenpropathrin | | x | G | E | u | x | F | u | E | u | 3d | see label |
| Deadline MP's (4% bait) | UN | x | x | x | x | 25 lb. | x | x | x | x | 12h | 75 lb. |
| metaldehyde | | x | x | x | x | G | x | x | x | x | 0d | 3 |
| Diazinon AG600 WBC (RUP) | 1B | x | x | 12.75 fl. oz./100 gal. | x | x | 12.75 fl. oz./100 gal. | x | x | x | 3d | 64 fl. oz. |
| diazinon | | x | x | G | x | x | F | x | x | x | 5d | 2 |
| Dibrom 8E (RUP) | 1B | 1 pt. | x | 1 pt. | x | x | 1 pt. | 1 pt. | x | 1 pt. | 2d | 4.7 lbs. |
| naled | | u | x | u | x | x | u | u | x | u | 1d | 5 |
| Entrust SC (2SC) | 5 | 4-6 fl. oz. | x | 4-6 fl. oz. | x | x | x | x | x | x | 4h | 18 fl. oz. |
| spinosad | | G | x | G | x | x | x | x | x | x | 1d | 3 |
| Exirel (0.83SE) | 28 | 13.5-20.5 fl. oz. | x | x | x | x | x | x | 13.5-20.5 fl. oz. | x | 12h | 61.5 fl. oz. |
| cyantraniliprole | | s | x | x | x | x | x | x | E | x | 1d | NA |
| Grandevo WDG | UN | 2-3 lb. | x | 1-3 lb. | x | x | 2-3 lb. | x | 2-3 lb. | 2-3 lb. | 4h | NA |
| <i>Chromobacterium subsugae</i> | | u | x | G | x | x | u | x | u | u | 0d | NA |

(Continued)

Table 9-6. Strawberry Insects - Postbloom Through Harvest¹ (continued)

| Product And Formulation Active Ingredient | IRAC Code ² | Eastern Flower Thrips | Leafhopper | Leafroller | Sap Beetle | Slugs | Spider Mite | Spittlebug | Spotted wing Drosophila | Tarnished Plant Bug | REI ³ PHI ⁴ | Max Amt ⁵ Max App ⁶ |
|---|------------------------|-----------------------|--------------|--------------|--------------|-----------|---------------|-------------|-------------------------|---------------------|-----------------------------------|---|
| Javelin WG | 11A | x | x | 0.25-1.5 lb. | x | x | x | x | x | x | 4h | UN |
| <i>Bacillus thuringiensis</i> | | x | x | G | x | x | x | x | x | x | 0d | NA |
| Kanemite 15SC | 20B | x | x | x | x | x | 21-31 fl. oz. | x | x | x | 12h | 62 fl. oz. |
| acequinocyl | | x | x | x | x | x | E | x | x | x | 1d | 2 |
| Magister SC (1.7SC) | 21A | x | x | x | x | x | 32-36 fl. oz. | x | x | x | 12h | 36 fl. oz. |
| fenazaquin | | x | x | x | x | x | u | x | x | x | 1d | 1 |
| Malathion 5EC | 1B | 1.5-3.2 pt. | 1.5-3.2 pt. | 1.5-3.2 pt. | x | x | 1.5-3.2 pt. | 1.5-3.2 pt. | x | 1.5-3.2 pt. | 12h | 12.8 pt. |
| malathion | | s | G | u | x | x | u | u | x | u | 3d | 4 |
| Nealta (1.67SC) | 25 | x | x | x | x | x | 13.7 fl. oz. | x | x | x | 12h | 27.4 fl. oz. |
| cyflumetofen | | x | x | x | x | x | G | x | x | x | 1d | 2 |
| Nexter (75WP) | 21 | x | x | x | x | x | 4.4-10.6 oz. | x | x | x | 12h | 21.34 oz. |
| pyridaben | | x | x | x | x | x | G | x | x | x | 1d | 2 |
| Oberon 2SC | 23 | x | x | x | x | x | 16 fl. oz. | x | x | x | 12h | 16 fl. oz. |
| spiromesifen | | x | x | x | x | x | G | x | x | x | 3d | 1 |
| Platinum 75SG | 4A | x | 1.7-4.01 oz. | x | x | x | x | x | x | x | 12h | 4.01 oz. |
| thiamethoxam | | x | E | x | x | x | x | x | x | x | 50d | NA |
| Portal XLO (0.4EC) | 21A | x | x | x | x | x | 2 pt. | x | x | x | 12h | 4 pt. |
| fenpyroximate | | x | x | x | x | x | E | x | x | x | 1d | 2 |
| Radiant SC (1SC) | 5 | 6-10 fl. oz. | x | 6-10 fl. oz. | x | x | x | x | 6-10 fl. oz. | x | 4h | 30 fl. oz. |
| spinetoram | | G | x | E | x | x | x | x | E | x | 1d | 3 |
| Rimon 0.83EC | 15 | 6-12 fl. oz. | x | x | 6-12 fl. oz. | x | x | x | x | 9-12 fl. oz. | 12h | 36 fl. oz. |
| novaluron | | u | x | x | E | x | x | x | x | E | 1d | NA |
| Savey 50DF | 10A | x | x | x | x | x | 6 oz. | x | x | x | 12h | 6 oz. |
| hexythiazox | | x | x | x | x | x | E | x | x | x | 3d | 1 |
| Sevin XLR Plus (4F) | 1A | x | 1-2 qt. | 1-2 qt. | x | x | x | 1-2 qt. | x | 1.5-2 qt. | 12h | 10 qt. |
| carbaryl | | x | G | F | x | x | x | x | x | G | 7d | 5 |
| Sluggo | UN | x | x | x | x | 20-44 lb. | x | x | x | x | 0d | UN |
| iron phosphate | | x | x | x | x | G | x | x | x | x | 0d | UN |

(Continued)

Table 9-6. Strawberry Insects - Postbloom Through Harvest¹ (continued)

| Product And Formulation Active Ingredient | IRAC Code ² | Eastern Flower Thrips | Leafhopper | Leafroller | Sap Beetle | Slugs | Spider Mite | Spittlebug | Spotted wing Drosophila | Tarnished Plant Bug | REI ³ PHI ⁴ | Max Amt ⁵ Max App ⁶ |
|---|------------------------|-----------------------|------------|----------------|------------|-------|-------------|------------|-------------------------|---------------------|-----------------------------------|---|
| Transform WG | 4C | 2.2 oz. | x | x | x | x | x | x | x | 1.5-2.2 oz. | 24h | 8.5 oz. |
| sulfoxaflor | | s | x | x | x | x | x | x | x | u | 1d | 4 |
| Vendex 50WP (RUP) | 12B | x | x | x | x | x | 1.5-2 lb. | x | x | x | 2d | 4 lb. |
| fenbutatin-oxide | | x | x | x | x | x | G | x | x | x | 1d | 2 |
| Verdepryn 100SL (0.83SL) | 28 | x | x | 8.2-11 fl. oz. | x | x | x | x | 8.2-11 fl. oz. | x | 4h | 33 fl. oz. |
| cyclaniliprole | | x | x | u | x | x | x | x | u | x | 1d | 3 |
| Zeal (72WP) | 10B | x | x | x | x | x | 2-3 oz. | x | x | x | 12h | 3 oz. |
| etoxazole | | x | x | x | x | x | E | x | x | x | 1d | 1 |

Footnotes: See how to read the spray schedule tables section at the beginning of the chapter.

Strawberry Postharvest And New Plantings - Disease

Apply every 10-14 days as needed.

Table 9-7. Strawberry Diseases - Postharvest And New Plantings¹

| Product And Dormulation Active Ingredient | FRAC Code ² | Anthrachnose Crown | Leaf Blight Phomopsis | Leaf Scorch (Diplocarpon) | Leaf Spot (Mycosphaerella) | Powdery Mildew | Fed Stele (Phytophthora) | REI ³ PHI ⁴ | Max Amt ⁵ Max App ⁶ |
|---|------------------------|--------------------|-----------------------|---------------------------|----------------------------|----------------|--|-----------------------------------|---|
| Aliette WDG | P07 (33) | x | x | x | x | x | 2.5-5 lb. | 12h | 30 lb. |
| aluminum tris | | x | x | x | x | x | E | 1d | NA |
| Captan (80WDG) | M | 1.8-3.75 lb. | 1.8-3.75 lb. | 1.8-3.75 lb. | 1.8-3.75 lb. | x | x | 24h | 30 lb. |
| captan | | E | G | G | G | x | x | 0d | NA |
| Cuproxat FL | M | x | 2.5-5 pt. | 2.5-5 pt. | 2.5-5 pt. | x | x | 48h | NA |
| tribasic copper sulfate | | x | F | F | F | x | x | NA | NA |
| Nordox 75G | M | x | 1.25-2.5 lb. | 1.25-2.5 lb. | 1.25-2.5 lb. | x | x | 12h | NA |
| cuprous oxide | | x | F | F | F | x | x | NA | NA |
| Orondis Gold | 49+4 | x | x | x | x | x | 20-62 fl. oz. Drench; 7-13.9 fl. oz. Foliar | 48h | 124 fl. oz. |
| oxathiapipralin+ mefenoxam | | x | x | x | x | x | E | 28d | 2 |

(Continued)

Table 9-7. Strawberry Diseases - Postharvest And New Plantings¹ (continued)

| Product And Dormulation Active Ingredient | FRAC Code ² | Anthraco- nose Crown | Leaf Blight Phomopsis | Leaf Scorch (Diplocarpon) | Leaf Spot (Mycosphaerella) | Powdery Mildew | Fed Stele (Phytophthora) | REI ³ PHI ⁴ | Max Amt ⁵ Max App ⁶ |
|--|---------------------------|----------------------------|--------------------------|------------------------------|-------------------------------|----------------|-----------------------------|--------------------------------------|--|
| Phostrol | 33 | x | x | x | x | x | 2.5-5 pt. | 4h | NA |
| phosphorous acid | | x | x | x | x | x | G | NA | NA |
| Prophyt | P07 (33) | x | x | x | x | x | 2-4 pt. | 4h | varies |
| phosphorous acid | | x | x | x | x | x | G | UL | 4 |

Footnotes: See how to read the spray schedule tables section at the beginning of the chapter.

Strawberry Postharvest And New Plantings - Insects

Table 9-8. Strawberry Insects - Postharvest and New Plantings¹

| Product And Formulation Active Ingredient | IRAC Code ² | Leafhopper | Leafroller | Root Weevils | Slugs | REI ³ PHI ⁴ | Max Amt ⁵ Max App ⁶ |
|--|------------------------|-----------------|----------------------------|-----------------|--------|--------------------------------------|--|
| Actara (25WDG) | 4A | 4 oz. | x | 4 oz. | x | 12h | 12 oz. |
| thiamethoxam | | E | x | E | x | 3d | NA |
| Assail 30SG | 4A | 4-6.9 oz. | 4-6.9 oz. | x | x | 12h | 13.8 oz. |
| acetamiprid | | G | G | x | x | 1d | 2 |
| Brigade eVo | 3A | x | 2.56-12.8 oz. | 3.2-12.8 oz. | x | 12h | 32 oz. |
| bifenthrin | | x | G | G | x | 0d | 3 |
| Danitol 2.4EC (RUP) | 3A | 16-21.3 fl. oz. | 10.66-21.33 fl. oz. | 16-21.3 fl. oz. | x | 24h | see label |
| fenpropathrin | | G | E | G | x | 3d | see label |
| Deadline MP's (4% bait) | UN | x | x | x | 25 lb. | 12h | 75 lb. |
| metaldehyde | | x | x | x | G | 0d | 3 |
| Diazinon AG600 WBC (RUP) | 1B | x | 12.75 fl. oz. /100 gal. | x | x | 3d | 64 fl. oz. |
| diazinon | | x | G | x | x | 5d | 2 |
| Dibrom 8E (RUP) | 1B | x | 1 pt. | x | x | 48h | 5 pt. |
| naled | | x | u | x | x | 1d | 5 |
| Entrust SC (2SC) | 5 | x | 4-6 fl. oz. | x | x | 4h | 18 fl. oz. |
| spinosad | | x | G | x | x | 1d | 3 |
| Grandevo WDG | UN | x | 1-3 lb. | x | x | 4h | NA |
| <i>Chromobacterium subtsugae</i> | | x | G | x | x | 0d | NA |
| Javelin WG | 11A | x | 0.25-1.5 lb. | x | x | 4h | NA |
| Bacillus thuringiensis | | x | G | x | x | 0d | NA |
| Malathion 5EC | 1B | 1.5-3.2 pt. | 1.5-3.2 pt. | 1.5-3.2 pt. | x | 12h | 8 pt. |
| malathion | | G | u | u | x | 3d | 4 |

(Continued)

Table 9-8. Strawberry Insects - Postharvest and New Plantings¹

| Product And Formulation Active Ingredient | IRAC Code ² | Leafhopper | Leafroller | Root Weevils | Slugs | REI ³ PHI ⁴ | Max Amt ⁵ Max App ⁶ |
|--|------------------------|--------------|--------------|--------------|-----------|--------------------------------------|--|
| Platinum 75SG | 4A | 1.7-4.01 oz. | x | 1.7-4.01 oz. | x | 12h | 4.01 oz. |
| thiamethoxam | | E | x | G | x | 50d | NA |
| Radiant SC (ISC) | 5 | x | 6-10 fl. oz. | x | x | 4h | 30 fl. oz. |
| spinetoram | | x | E | x | x | 1d | 3 |
| Sevin XLR Plus (4F) | 1A | 1-2 qt. | 1-2 qt. | x | x | 12h | 10 qt. |
| carbaryl | | G | F | x | x | 7d | 5 |
| Sluggo | UN | x | x | x | 20-44 lb. | 0d | NA |
| iron phosphate | | x | x | x | G | 0d | NA |

Footnotes: See how to read the spray schedule tables section at the beginning of the chapter.

Special Comments on the Strawberry Schedule

Annual plasticulture strawberry

Timely planting of healthy plugs is key to establishing a successful planting. Planting date influences runner and branch crown formation. Too early is better than too late. If planted too early, energy is directed toward runner formation; if planted too late, the 4-5 branch crowns that are desired at flowering may not develop in time. Research in Ohio and central Kentucky has identified early to mid-September as best for plugs in most years. Growers further north or at higher elevations in the Appalachian Mountains may be able to plant in late August, while growers further south or west may be able to plant slightly later. On-farm research over a number of years gives growers the best planting window for their particular location.

Weed management in the row is usually accomplished by applying a pre-emergent herbicide beneath the plastic prior to laying the plastic to control winter annuals. Growers use different strategies to manage weeds between rows. Many have had success by simply planting a cover crop of annual ryegrass or cereal rye to suppress weeds and then killing it with a graminicide in the spring. Insect management is usually not a significant issue. All growers should plant disease-free plugs; however, it's also a good idea to make a fungicide application for anthracnose crown rot to the plug trays or to the plants in the field after planting, especially since infections may be present but symptoms can be delayed or go unnoticed until they become more serious. To prevent the build-up of pathogens in the soil a second year of production with annual plants is not recommended.

Neopestalotiopsis disease

Neopestalotiopsis disease is an emerging fungal disease in the Midwest. The fungus can affect all parts of the plant (leaves, crown, roots, fruit) with leaf symptoms usually being the first to appear in the Midwest. The disease progresses rapidly under cool and wet conditions and can cause significant plant losses in a short amount of time. Fruit symptoms are uncommon in the Midwest because the plants typically die before the flowers develop into fruit.

Management is extremely difficult. The primary source of the fungus is contaminated nursery stock. Transplants should be inspected upon delivery for symptoms and plants confirmed to be infected should not be planted. There are currently no resistant cultivars and no fungicides specifically registered for Neopestalotiopsis disease. A few fungicides (Thiram and Switch), which are registered for other strawberry diseases have good to excellent efficacy but must be applied intensively throughout the production season.

Images and additional information on Neopestalotiopsis disease can be found at: <https://plantpathology.mgcafe.uky.edu/files/ppfs-fr-s-12.pdf>.

Table 9.9 Disease Resistance Of Strawberry Cultivars Commonly Grown In The Midwest¹

| Cultivar | Leaf Spot | Leaf Scorch | Powdery Mildew | Verticillium Wilt | Phytophthora Crown Rot | Red Stele (Phytophthora) |
|---------------------|-----------|-------------|----------------|-------------------|------------------------|--------------------------|
| June Bearing | | | | | | |
| AC Valley Sunset | R | R | U | R | U | R |
| AC Wendy | T | T | R | S | U | R |
| Allstar | T | T | R | R | S | R |
| Annapolis | S | S | S | T | U | R |
| Brunswick | U | U | U | U | U | R |
| Cabot | T | T | R | U | U | R |
| Camarosa | S | S | S | U | R | U |
| Cavendish | R | R | R | T | S | R |
| Chandler* | S | S | S | S | S | S |
| Clancy | R | R | R | U | U | R |
| DarSelect | T | T | S | U | U | U |
| Daroyal | U | U | U | U | U | U |
| Donna | U | U | U | U | U | U |
| Earliglow | R | R | T | R | S | R |
| Flavorfest | R | R | T | T | U | R |
| Galletta | U | U | U | U | U | U |
| Glooscap | T | T | T | S | U | S |
| Guardian | S | R | S | R | S | R |
| Herriot | T | T | U | R | U | U |
| Honeoye | R | R | T | S | S | S |
| Itasca | R | R | U | U | U | R |
| Jewel* | S | S | T | S | S | S |
| Kent | S | S | T | S | U | S |
| L'Amour | R | R | T | U | U | R |
| Lateglow | R | R | T | R | S | R |
| Mayflower | U | U | U | U | U | U |
| Mesabi | R | R | R | R | U | R |
| Mira | S | S | R | U | U | R |
| Northeast | T | T | S | R | U | R |
| Ovation | R | R | S | U | U | R |
| Redchief | S | R | R | R | S | R |
| Seneca | U | U | U | S | U | S |
| Sonata | U | U | T | U | U | S |
| Sparkle | S | S | S | S | S | S |
| Surecrop | R | R | R | R | S | R |
| Sweet Charlie | U | U | R | U | R | U |
| Winona | R | R | T | T | U | R |
| Day Neutral | | | | | | |
| Albion* | T | T | S | R | T | R |
| Tribute | R | R | R | R | S | R |
| Tristar | R | R | R | R | S | R |
| Seascape | S | S | R | U | U | R |
| San Andreas | S | S | R | T | U | U |

R=resistant. S=susceptible. T=tolerant. U=unknown.

*Recent research from Ohio and Indiana indicate that these varieties are very susceptible to aggressive isolates of *Neopestalotiopsis* sp. Transplants should be monitored for symptoms before and immediately after planting.

Vole Control in Fruit Plantings

Mice, also known as voles, can cause serious damage to tree fruit plantings. Frequently, damage occurs but growers do not notice it until trees become weak, die, or need to be removed.

You can anticipate vole damage each year, particularly from late summer to early spring, as mice eat bark from the base of small saplings. Such damage can girdle and kill a tree. Apple trees are most susceptible, but hungry voles will attack other fruit trees. Apple trees on dwarfing rootstocks are particularly palatable to them.

Many plantings are made in a hedgerow pattern, which does not permit cultivation between trees. Such plantings favor vole migration, as do mulches and vigorous sods. High populations also favor vole migrations.

Voles can be a problem in blueberry plantings but almost never feed on grapes, blackberries, raspberries, or strawberries.

General Orchard Management Practices

You can employ several general orchard management practices to reduce the risk of injury and improve control. No single material or technique is effective for complete control of voles. We therefore suggest you vary both the materials and methods of control during the season.

You can construct tree guards from “hardware cloth” or similar materials with a mesh no larger than 0.25 inch. These guards should enclose the tree and extend from several inches below soil surface — voles dig in the top 2 to 3 inches of soil — to several inches above maximum snow line (about 18 inches).

Placing pea-sized gravel or cinders around the trees in a circle 4 to 6 inches wide and at the same depth tends to discourage meadow voles from attacking crowns of trees, but does not discourage other mouse species.

Voles need abundant cover to proliferate. Maintaining a clean area 1 to 2 feet around the base of each tree discourages surface feeding and also regulates vole populations long term. Chemical weed control in early spring significantly reduces the amount of labor involved in keeping the area around the tree clean.

Mow short the orchard cover or sod in late August and again after harvest to reduce runway cover and aid baiting. Cleaning out drainage ditches and fencerows, and picking up or crushing all dropped fruit, discourages large vole populations.

Orchard Vole Control Program

Essential Knowledge

The first thing you must do to control voles is to determine the problem species. Use snap traps for this. The three common species are meadow vole (*Microtus pennsylvanicus*), prairie vole (*Microtus ochrogaster*), and pine vole (*Pitymys pinetorum*). While the control materials for these species may be the same, the control methods differ.

You can make quick field identifications of vole species (for both juveniles and adults) based on the length of their tails:

Pine vole: Tail is about as long as its hind foot.

Meadow and prairie vole: Tail is about twice as long as its hind foot.

Determine timing and site of infestations with snap traps. Knowing when and where mice are most abundant makes control easier.

Control

You can control voles in orchards by using either zinc phosphide or chlorophacinone baits. You must use both baits according to label directions.

Zinc phosphide, a restricted use pesticide, is an acutely toxic bait that kills mice within 24 hours. It is available either as a weather-resistant pellet bait or mixed with prepared grains such as oats and corn. Zinc phosphide is usually well accepted by mice. However, it is not effective if applied more than twice.

Chlorophacinone (e.g., RoZol) is an anticoagulant bait available as a weather-resistant, pellet-style bait. It is highly accepted by rodents but does not kill them for several days. For effective control, make a second application of chlorophacinone within 20 to 40 days.

Use caution: Baits can be attractive to other wildlife, including birds, and domestic pets. You must apply bait directly in runways or bait stations (see below) or broadcast. Pick up all spilled materials to avoid consumption by non-target animals.

Efficacy of Baits Against Meadow and Pine Voiles

Chlorophacinone is more effective against pine voles than meadow voles. Zinc phosphide is more effective against meadow voles than pine voles. Consistently using just one of these chemicals results in population shifts from one vole species to another. Therefore, alternate baiting using zinc phosphide in the first application, followed by chlorophacinone in the second application, to reduce populations of both species.

Baiting Techniques

There are three main baiting techniques.

1. **Machine baiting.** You expose bait in an artificial trail (Trail Builder).
2. **Trail baiting.** You expose bait only in natural, active runways.
3. **Broadcast baiting.** You broadcast bait by hand, cyclone-type seeder, or tractor-drawn equipment at recommended rates. *This technique is not recommended for pine vole control.* When using zinc phosphide baits, the 2% concentration is recommended.

Observe safety precautions: Zinc phosphide is a restricted use material. Read and follow all label directions and precautions.

Comparison of Baiting Techniques

| Baiting Technique | Meadow Or Prairie Vole Control (%) | Pine Vole Control (%) |
|-------------------|------------------------------------|-----------------------|
| Machine | 90-95 | 80-85 |
| Trail | 80-85 | 70-75 |
| Broadcast | 78 | Not Recommended |

Timing

Apply rodenticides on a sunny day in late fall when voles are active. Voles begin to build up in early August, but delay baiting as late as possible in the fall. The most effective application period is just before snow cover, after frost reduces the grass cover and the fruit is rotted. Spot treatment during the winter and into early spring is recommended. Treat marginal lands to prevent re-invasion.

Pre-harvest Baiting Is Not Recommended

Applying poisoned bait before harvest to prevent vole damage to fruit in cold storage is not a sound practice for the following reasons:

1. The recommended methods of orchard vole control do not always provide 100 percent control. Some voles survive the pre-harvest control and enter fruit boxes on the ground that are carried into cold storage.
2. The pre-harvest poison application reduces the population of voles in the orchard, which greatly reduces competition among survivors, making food and cover ample. Under these favorable conditions, survivors breed, with as many as eight young per litter. In a very short time, populations may recover to original levels.
3. The recommended control season for voles in orchard and winter storage facilities is just prior to freezing conditions. *Note: Check your control*

program with snap traps. Lack of visible damage does not indicate the efficacy of your program.

Control in Storage

To protect fruit in storage from rodents, pay attention to what you do before and during harvest.

Before harvest

- Poison rats and mice in storage one month before picking. Keep storage area baited and free of debris.
- Clean up outside debris one week before picking. Pay special attention near loading areas.
- Use rodent-proof storage. Seal all holes and cracks. Mice can fit through a hole the size of a dime.

During harvest

- Move filled boxes into storage quickly. Any box left overnight may have mice.
- As you load fruit into storage, bait the storage area. Place teaspoonful amounts in bait stations, on floor, along alleys, between rows of boxes, and under pallets. Do not place open baits on floors or any areas where contamination might occur. Commercial bait stations are available from agricultural supply companies. Always prevent contact with fruit.

Bait Stations in the Orchard

You can prepare bait stations in several ways and eliminate or reduce the opportunity for non-target animals to contact the bait. Squares of heavy roofing shingles (or other weather-resistant materials) placed out of traffic areas between trees can serve as bait stations to protect the bait and hiding of rodents.

Some growers have constructed bait stations that require less refilling by building inverted T-shaped stations from PVC tubing and fittings that provide bait storage and a protected feeding area. Place bait stations in the field two or three weeks before adding the bait.

Vole Control for Small Fruit

Prozap zinc phosphide pellets are a restricted use pesticide labeled for vole control in highbush blueberries, blackberries, and red and black raspberries. Apply this product only in the dormant season after final harvest and not later than the beginning of leaf emergence in the spring. The minimum pre-harvest interval is 70 days. Do not apply when the ground is snow-covered.

You may broadcast bait with a cyclone seeder or by hand. When applying by hand, throw a tablespoon (12 grams) into heavy cover along bushes, rocky outcrops, and fence lines. Make two applications at a rate of 6 to 10 lb per acre per application at a minimum interval of 21 days. The maximum application per growing season is 20 lb. per acre.

Growth Regulator PHIs and REIs

Growth Regulator Pre-harvest Intervals and Restricted Entry Intervals

| Trade Name | Common Name | Preharvest Interval (Days) | | | | | | REI (Hours) |
|---|------------------------|----------------------------|-----------------|-------|-----------------|-------------|------|-------------|
| | | Apple | Pear | Peach | Sweet Cherry | Tart Cherry | Plum | |
| Apogee, Cryova ² , Kudos, Pro-Hex ² | Prohexadione-calcium | 45 | — ¹ | — | 20 | — | — | 12 |
| Amid Thin-W | NAD | 2 | 2 | — | — | — | — | 48 |
| Blush | | 7 | — | — | — | — | — | 4 |
| Cytokin ² | kinetin | 0 | 0 | 0 | 0 | 0 | 0 | 4 |
| Ethrel, Motivate, Ethephon | ethephon | 7 | — | — | 7 | 7 | — | 48 |
| Fruitone N | NAA | 2 | 2 | — | — | — | — | 48 |
| K-Salt Fruit Fix 200 | NAA | 2 | 2 | — | — | — | — | 24 |
| K-Salt Fruit Fix 800 | NAA | 2 | 2 | — | — | — | — | 48 |
| MaxCel, Exilis | 6-benzlidenine | 86 | 86 | — | — | — | — | 12 |
| ProGibb | gibberellic acid (GA3) | — | — | — | 0 | 0 | 0 | 4 |
| ProVide | GA4 + 7 | — | — | — | — | — | — | 4 |
| Promalin, Typy, Cytoplex HMS, Perlan | 6BA + GA4 + 7 | 0 | NB ³ | 0 | NB ³ | — | — | 4/24 |
| ReTain | AVG | 7 | 7 | 7 | — | — | 7 | 12 |

¹ — = not registered or not recommended

² Check label for state registration

³ Non-bearing trees only.

Chemical Weed Control in Fruit Crops

Controlling weeds in fruit plantings is important. Weeds may reduce yields by competing for water and nutrients, harbor insects and other pests, and serve as alternate hosts for diseases. Herbicides can provide good weed management with less labor and frequently at a lower cost compared to manual weed control.

Proper Application

Herbicide effectiveness depends on the selection of the appropriate product and application of the product at the proper time, and the proper rate, with the proper equipment. The level of weed management depends largely on the operator's skill and attention to detail. In most cases, the given herbicide rates are for overall coverage (broadcast rates). For band treatment common in fruit plantings, reduce the amounts according to the portion of area treated. For example, to control weeds in a 4-foot-wide band beneath a crop planted in rows 10 feet apart, the amount of herbicide needed per

acre of crop is 4/10 of the broadcast amount per acre. Make sprayer adjustments and calibrations as precise as possible to assure accurate and uniform applications. Improper application can damage fruit plantings and may result in illegal residues on the fruit crop. Over- or under-application also can reduce the profitability of the planting. Use nozzles appropriate for herbicide application at low pressures (20-40 psi) on a fixed boom-type applicator, unless the label has specific recommendations. This type of sprayer is calibrated easily and, when designed properly, deposits herbicide uniformly.

Consider using one of the recently introduced low-drift nozzles such as the Turbo TeeJet Nozzle or TurboDrop Nozzle. They have been designed to provide similar performance to traditional flat fan nozzles while reducing the number of very small droplets that are highly subject to drift.

While backpack or hand sprayers may be suitable for spot treatment with post-emergence herbicides, do not use them to apply pre-emergent herbicides around fruit plants. The application rate is critical with pre-emergent herbicides, and hand sprayers cannot be

calibrated well enough for accurate application. Slight application rate errors can cause severe damage to fruit plants.

Calibrate each sprayer carefully and apply herbicides according to the suggested rates. Note that when applying many pre-emergence herbicides to the soil, you should adjust rates according to soil characteristics. Generally, use lower rates on sandy soils with low organic matter, and use higher rates on heavier textured soils and those high in organic matter. With some herbicides, no rate changes are suggested. If you are unsure about an herbicide's effectiveness or possible crop damage, test it on a small portion of the planting before using it extensively.

Herbicide Resistance Management

Continued use of the same herbicide can lead to the development of herbicide-resistant weeds or the establishment of tolerant weeds. Avoid using the same product or chemically related products for several consecutive years to avoid building up herbicide-resistant weed biotypes. The HRAC code on the label indicates what group the chemical is in and chemicals with the same number function the same way. We recommend that you rotate herbicides with different modes of action and include non-chemical controls whenever possible to avoid these problems and improve weed control.

Tank Mixes

Certain herbicides may be combined in suitable tank mixes. Consult product labels for approved combinations and recommended rates. Use caution when tank mixing herbicides that are not specifically listed on the label.

By using tank mixes, you can apply a pre-emergence herbicide together with a post-emergence herbicide to provide improved weed control, or you can apply two herbicides to gain better weed control. **Always follow label recommendations. Improper mixing can form chemical compounds that are not compatible and that may damage your sprayer.**

Timing of Applications

Weed management may require multiple applications each year. Timing is important for best results.

Growers often apply a post-emergence herbicide in early spring to control winter annuals and perennials before they flower. The timing of this application may be too early for maximum pre-emergence herbicide effectiveness. It is often wise to follow the first application with a second application of a tank mix

of post- and pre-emergence herbicides about three weeks after the first. This controls any weeds that have emerged since the first application and puts the pre-emergence in place at the right time, so it lasts through the main period of weed emergence.

Site Preparation Before Planting

Management of perennial weeds in perennial fruit crops can be challenging. Growers should strive to eradicate established perennial weeds during site preparation in the season prior to establishing the crop. Most perennial weeds cannot be controlled effectively in the spring before planting or once the crop is planted. Ideally, perennial broadleaf weeds should be approximately at the bud to early flowering stage at the time of treatment. Summer and early fall applications of glyphosate may be more effective against perennial broadleaf weeds than spring applications. Allow five to seven days for glyphosate to translocate throughout the root system before plowing under. This should be followed by repeated shallow cultivation as green "flushes" of weed seedlings appear. An alternative is to apply paraquat (Gramoxone) or glufosinate (Rely or generics) for contact non-selective weed control as flushes of weed seedlings appear.

Establishment of a grass crop on the site several years before planting will give the grower more options for control of perennial broadleaf weeds. Alternatively, cultivation combined with a non-selective herbicide can also be an effective strategy.

Trade Name and Active Ingredient (a.i.)

Herbicide labels list the chemical names of the active ingredients and the percentage or amount of the active ingredients as "a.i." Herbicides come in various formulations and under various trade names. For the sake of brevity, only the original trade name is listed in this guide. See the table on pages 255-260 for other trade names registered for use on fruit crops.

Always read each label carefully, as rates and labeled crops may differ between labels with similar active ingredients. Follow the recommended rates as they are listed on the label of the product you plan to use.

Follow the recommended rates as they are listed on the label of the product you plan to use.

Use Restrictions

Federal regulations control herbicide use and prescribe the crops the herbicides can be used on, as well as the timing and rates for which these materials are registered. Use only registered materials at the recommended rates for the crops listed. Herbicides are covered by

Worker Protection Standards where they apply. Labels include restricted-entry intervals (REI) and personal protective equipment (PPE) information. Product labels are the final authority — follow them carefully.

Good Rules to Remember

1. The rates recommended in this guide are mid-range rates applicable for medium to fine soils. Always refer to labels for full details about rates depending on soil type, organic matter content, age of plants, etc.
2. Applying post-emergence herbicides under stress conditions to weeds (such as high temperatures in midsummer, drought, cool temperatures in the spring, etc.) may result in poor weed control.
3. Use a fixed-spray boom, appropriate nozzles, and low pressure for even application without drift.
4. Spray only in little or no wind (less than 5 mph).
5. Adjust rates according to bandwidth.
6. Follow herbicide restrictions on new plantings. Allow plants to become well established and the soil well settled around plants before application.
7. Use herbicide sprayers for herbicides only.
8. Clean sprayers thoroughly when changing herbicides, especially when you have used 2,4-D, Chateau, or Prowl.
9. Store pesticides in locked storage. Do not allow liquid pesticides to freeze.
10. Protect the environment — avoid surface or ground water contamination. Dispose of excess spray material carefully and according to label directions. Do not allow grazing in treated areas.
11. **Read the label. Understand it thoroughly. Follow its directions.**

Herbicide Recommendations For Apple And Pear

For generic herbicides, see pages 281-284.

Herbicide Recommendations For Apple And Pear

| Weed Problem | Material And Rate Per Acre | Notes And Comments |
|---|--|---|
| Pre-emergence | | |
| Annual grasses and broadleaves | Alion (Indaziflam 1.67 lb. a.i./gal.) at 3.5-6.5 fl. oz. in minimum of 10 gal. water | Trees must be established at least 3 years after transplanting. Use lower rates as soil OM decreases. Avoid direct or indirect spray contact with crop foliage, green bark, roots, or fruit, as it may cause localized crop injury and death. Allow at least 30 days between applications. Do not exceed 10.3 fl. oz. per acre in a 12-month period. Do not apply to frozen ground. Do not apply within 25 feet of ponds, rivers, streams, or wetlands. Spot spraying is not recommended. Shake container well before use. |
| Annual broadleaves | Bellum (mesotrione 4 lbs. a.i./gal.) at 3 -6 fl. oz. | See Post-emergence section for details. |
| Annual and perennial grasses and broadleaves | Casoron 4G (dichlobenil 4% a.i./lb.) at 100-150 lb. | For perennial weed control, apply to untilled ground over old weed growth from November 15 to Feb 15; alternately apply late fall or very early spring before May 15 and incorporate immediately. For annual weed control, surface apply. Shallow incorporation or sprinkler irrigation is recommended when application is made during periods of high temperatures. Do not apply until 4 weeks after transplanting. Use higher rate for perennial weed control. Annual maximum rate 150 lbs./A. |
| Annual broadleaves and suppression of grasses | Chateau SW (flumioxazin 51% a.i.) at 6-12 oz. Chateau EZ (flumioxazin 41.4% a.i.) at 6-12 oz. | The preferred application timing is in the fall to maximize the potential for rainfall to activate and set the herbicide. Do not apply to trees established less than 1 year unless protected from spray contact by nonporous wraps. Make applications only to berms. Do not apply after pink bud on apples and bud break in pears. For non-bearing trees, do not apply during the period after flowering through leaf drop unless shielded application equipment ensures that spray drift will not contact crop foliage. Do not apply to fine textured soils. Do not exceed 2 applications in a growing season or make a sequential application within 30 days of the first application. Do not exceed 24 oz. per season Do not apply when plants are under stress. Do not apply within 300 yards of nondormant pears. Apply alone pre-emergence or tank mix with Gramoxone post-emergence with a crop oil 1% v/v or NIS 0.25% v/v. Do not incorporate. Do not allow drift to contact foliage or green bark. Trees may be transplanted 2 months after application. Minimum 30 days between applications. |

(Continued)

Herbicide Recommendations For Apple And Pear (*continued*)

| Weed Problem | Material And Rate Per Acre | Notes And Comments |
|--|---|--|
| Annual broadleaves and annual grasses | Chateau Complete (flumioxazin 30% a.i. + rimsulfuron 8.25%) at 12 oz. | Not registered in all states. The preferred application timing is in the fall to maximize the potential for rainfall to activate and set the herbicide. Do not apply to trees established less than 1 year. Do not apply after pink bud on apples and bud break in pears. Do not apply within 300 yards of nondormant pears. Make applications only to berms. Avoid direct or indirect spray contact to foliage and green bark (nonbarked trunk with the exception of undesirable suckers). Do not apply to powdery soils or soils that are susceptible to wind displacement unless irrigation can be applied immediately after application. Do not mow treated areas between bud break and final harvest. Do not incorporate. Max seasonal rate 12 oz./A. |
| Annual broadleaves and suppression of grasses | Goal 2XL (oxyfluorfen 2 lb. a.i./gal.) at 2-8 pt. in minimum of 20 gal. water | Dormant Application Only: Effective both pre-emergence (5-8 pt.) and post-emergence (2-8 pt.) as directed spray on weeds larger than 4 inches. Do not apply from bud swell until harvest completion. Can be mixed with other pre-emergence herbicides or with Roundup or Gramoxone. Do not exceed 8 pt. per year. |
| Annual broadleaves and suppression of grasses | GoalTender (oxyfluorfen 4 lb. a.i./gal.) at 2.5-4 pt. in minimum of 20 gal. water | Dormant Application Only: Effective both pre-emergence banded application (2.5-4 pt.) and post-emergence (1-4 pt.) as directed spray on weeds larger than 4 inches. Do not apply from bud swell until harvest completion. Can be mixed with other pre-emergence herbicides or with Roundup or Gramoxone. Do not exceed 3 pt. per year on a broadcast basis. |
| Annual grasses and broadleaves | Karmex DF (diuron 80% a.i.) at 4 lb. in 25-40 gal. water | Apple Only: Effective both pre-emergence and post-emergence (min. 70°F with high humidity). Apply under trees established at least 1 year. Do not treat trees grafted on full-dwarf rootstocks. Do not exceed 1 application per year. May be tank mixed with Sinbar (1.5-2 lb. each) in orchards established at least 2 years. Karmex/Sinbar can be applied in the spring before weeds emerge or after harvest in the fall. |
| Annual and perennial grasses and certain broadleaves | Kerb SC (pronamide 35.6% a.i.) at 2.5-9.5 pt. in 40-50 gal. water. Rate depends on weed pressure and soil type. See table on label. | Apply as a directed spray in the fall after harvest prior to soil freeze-up. Rainfall or irrigation are required to activate. Do not apply more than 5 pt./A/year or make more than 1 application per year. Age restriction: Kerb SC may not be applied to (1) trees less than 1 year old, (2) fall-transplanted stock transplanted less than 1 year old, or (3) spring-transplanted stock transplanted less than 6 months. Some Special Local Needs Labels (FIFRA 24(c)) are labelled as Restricted Use Pesticides (RUP), whereas the national label remains a general use pesticide. |
| Annual grasses and broadleaves | Matrix FNV (rimsulfuron 25% a.i.) at 4 oz. in minimum of 10 gal. water | Apply only to crops that have been established for 1 full growing season and are in good health and vigor. Weeds are controlled for 60-90 days after application. Matrix burns down small, actively growing weeds less than 1 inch tall. When weeds are present at application, a labeled burndown herbicide such as glyphosate, paraquat, or glufosinate with an appropriate adjuvant improves control. Avoid direct or indirect contact with crop foliage or fruit, except undesirable suckers. Do not use Matrix FNV in a spray solution with a pH below 4.0 or above 8.0. Best results are obtained when the soil is moist at the time of application and 0.5 inch of rainfall or sprinkler irrigation occurs within 2 weeks of application. |
| Annual grasses and broadleaves | Pindar GT (penoxsulam 0.083 lbs./gal. + oxyfluorfen 3.93 lbs./gal.) at 1.5-3 pt. | Apply only to trees established at least 4 years; see label for reset requirements. Use trunk guards to protect plants until adequate bark has developed. Application window beginning after harvest, up to bud swell. See label for tank partners for complete burndown. A single rainfall or sprinkler irrigation of 0.5 inches or more, or flood irrigation within 21 days after application, is necessary to activate. The addition of 1 quart per acre of crop oil concentrate or methylated seed oil, or 0.25% v/v of an 80% active nonionic surfactant labeled for application to growing food crops, is required for effective postemergence control of susceptible emerged weeds. Avoid direct contact with fruit trees. Single max application rate 3 pts./A; 4.5 pts./A max total for the year with 30 days between sequential applications at lower rates. |
| Annual grasses and broadleaves | Princep 4L (simazine 4 lb. a.i./ga) at 2-4 qt./A in min of 20 gal. water | Apply under trees established at least 1 year. Apply in spring before weeds emerge, avoiding contact with fruit, foliage, or stems. Avoid use on sandy soils with OM <1%. Limited to 1 application per calendar year, with a single max rate of 4 qt./acre. Check label for state-specific restrictions on tank mixes. |

(Continued)

Herbicide Recommendations For Apple And Pear (*continued*)

| Weed Problem | Material And Rate Per Acre | Notes And Comments |
|---|--|---|
| Broadleaves | Sandea (halosulfuron-methyl 75% a.i.) 0.5-1 oz. in minimum of 15 gal. water | Apple and Pear east of the Mississippi: Apply only to trees established at least 1 year, making sure to avoid spray or drift contact with tree foliage and fruit. Apply a single or sequential application to the orchard floor on either side of the row based on weed pressure. Apply to bare ground for best results. If small weeds are present, mix with a post-emergence broad-spectrum herbicide. See label for west of the Mississippi applications. |
| Annual grasses and broadleaves | Sinbar WDG (terbacil 80% a.i.) at 0.5-4 lb. in minimum of 20 gal. water | Apple Only: Apply either in the spring before weeds emerge or during early stages of seedling growth, or after harvest in the fall. Trees must be established at least 3 years. Do not contact foliage or fruit with spray or mist. Non-bearing: (young, newly planted) Apple: Apply at 0.5-1 lb. Make the first application after a significant rainfall or irrigation event that allows the ground to settle around the base of the trees. Make 1-2 applications per season. Do not exceed 1 lb. per year. Do not use on soils with <1% organic matter. |
| Annual grasses and broadleaves and suppression of yellow nutsedge | Solicam DF (norflurazon 78.6% a.i.) at 2.5-5 lb. in minimum of 20 gal. water | See label for rate based on soil type. Apply a directed spray to settled and firm soil from fall to early spring before weeds emerge. Soil should be settled and firm. Rainfall or irrigation of 0.5 inch is needed within 4 weeks. Do not contact fruit or foliage. Do not apply after bud break on sandy loam soils. Check label for maximum amount allowed per year depending on soil type. Apple: Can be applied immediately after planting. Pear: Minimum 12 months after planting east of the Mississippi River before first application; minimum 18 months after planting west of the Mississippi before first application. |
| Annual grasses and certain broadleaves | Surflan As Specialty Herbicide (oryzalin 4 lb. a.i./gal.) at 2-6 qt. in 20-40 gal. of water | No longer in production - supply extremely limited to not available. Make a single band or broadcast application to the ground beneath trees before weeds emerge. Apply alone to weed-free soil or post-emergence mixed with Roundup or Gramoxone. Rainfall or irrigation (0.5 inch) is required for activation. Minimum 2.5 months between applications. Do not exceed 12 qt. per year. |
| Annual and perennial grasses and broadleaves | Zeus Prime XC (carfentrazone-ethyl 3.5% and sulfentrazone 31.8% a.i.) at 7.7-15.2 fl. oz. per acre in minimum of 10 gal. water | Apple only: Apply as a broadcast or banded soil application directed to the base of the trunks. If weeds are present, tank mix with a post-emergence herbicide to eliminate emerged weeds. Apply a single broadcast application at 15.2 fl. oz. per acre (0.41 lb. a.i./acre). May be applied as a banded treatment twice per year. Do not exceed 15.2 fl. oz. (0.41 lb. a.i)/acre/year. Do not apply after petal fall except with a hooded shielded sprayer. Do not tank mix with Chateau or with other products containing sulfentrazone. Age Restriction: Apply to crops that have been growing for at least 2 years and are in good condition. |
| Post-emergence | | |
| Annual broadleaves | Aim EC Herbicide (carfentrazone 2 lb. a.i./gal.) at 2 fl. oz. in 20 gal. of water | Apply any time during the season. Always add NIS 0.25% v/v or crop oil 1% v/v. Mix with Roundup or Gramoxone for broader weed control. Do not exceed 7.9 fl. oz. per year. Minimum 14 days between applications. Sucker Control: Apply when suckers are green. Do not allow spray to contact fruit, foliage, or green bark. |
| Annual and some perennial broadleaves | 2,4-D (Amine) at 1-4 pt. | Apply as directed spray to annuals 1-2 inches high and to perennials up to early bud stage. Do not allow spray to contact leaves, fruit, limbs or exposed roots of tree. Use coarse spray and low pressure to avoid drift. Do not apply during windy periods, when there is a temperature inversion or at extremely high temperatures. Non-bearing trees must be established at least 1 year. On bearing trees, do not apply during bloom and only after irrigation. Do not apply to bare ground. Use higher rate on perennial weeds. Do not exceed 2 applications per year. Maximum 75 days between applications. |

(Continued)

Herbicide Recommendations For Apple And Pear (*continued*)

| Weed Problem | Material And Rate Per Acre | Notes And Comments |
|---|--|---|
| Annual and some perennial broadleaves | Embed (2,4-D Choline) at 1-4 pt. in a min of 10 gal. | Not registered in all states. Apply only to orchards that have been established for at least one year and are in vigorous growth condition. Apply to annual weeds when small and actively growing. Use higher rate on perennial weeds in bud to bloom stage. Do not allow spray to contact leaves, fruit, limbs or exposed roots of tree. Use coarse spray and low pressure to avoid drift. Do not exceed 2 applications per year. Maximum 75 days between applications, max total rate 8.42 pints/A. |
| Annual broadleaves | Bellum (mesotrione 4 lbs. a.i./gal.) at 3 -6 fl. oz. | Apply only to trees established for one full year and in good health and vigor. Use trunk guards to protect plants until adequate bark has developed. For pre-emergent residual control, apply before rainfall or irrigation. See label for tank mix partners. Apply in a spray volume of 10-40 gals./A. Total application rate in a 12-month period 12 fl. oz./A. Single maximum application rate 6 fl. oz./A. Three applications per year allowed in a 12-month period when using reduced rates, allowing at least 12 weeks between applications at the 6 fl. oz./A rate or 6 weeks between applications at the 6 fl. oz./A rate and subsequent applications of 3 fl. oz./A. For application to emerged weeds, the use of crop oil concentrate (COC) type adjuvant at 1% v/v or non-ionic surfactant (NIS) at 25% is advised. |
| Annual broadleaves and suppression of grasses | Chateau SW (flumioxazin 51% a.i.) at 6-12 oz. in 15-75 gal. water | The preferred application timing for Chateau SW Herbicide is in the fall to maximize the potential for rainfall to activate and set the herbicide. Do not apply to trees established less than 1 year unless protected from spray contact by nonporous wraps. Do not apply after pink bud on apples or bud break in pears. Do not apply to fine textured soils. Do not exceed 2 applications in a growing season or a sequential application within 30 days of the first application. Do not apply when plants are under stress. Do not apply within 300 yards of nondormant pears. Apply alone pre-emergence or tank mix with Roundup or Gramoxone post-emergence with a crop oil 1% v/v or NIS 0.25% v/v. Do not incorporate. Do not allow drift to contact foliage or green bark. Do not exceed 24 oz. per season. Minimum 30 days between applications. |
| Annual broadleaves and suppression of grasses | Chateau EZ (flumioxazin 41.4% a.i.) at 6-12 oz. | The preferred application timing for Chateau EZ Herbicide is in the fall to maximize the potential for rainfall to activate and set the herbicide. Do not apply to trees established less than 1 year unless protected from spray contact by nonporous wraps. Do not apply after silver bud on apples and bud break in pear. Do not apply to fine textured soils. Do not exceed 2 applications in a growing season or make a sequential application within 30 days of the first application. Do not apply when plants are under stress. Do not apply within 300 yards of nondormant pears. Apply alone pre-emergence or tank mix with Roundup or Gramoxone post-emergence with a crop oil 1% v/v or NIS 0.25% v/v. Do not incorporate. Do not allow drift to contact foliage or green bark. Do not exceed 24 oz. per season. Apple and pear trees can be transplanted 2 months after application. Minimum 30 days between applications. |
| Annual broadleaves and annual grasses | Chateau Complete (flumioxazin 30% a.i. + rimsulfuron 8.25%) at 6-12 oz. | Not registered in all states. See Pre-emergence section for details. If weeds are emerged at the time of application, apply with an adjuvant (0.25% v/v non-ionic surfactant or 1 qt./A crop oil concentrate). For complete control of emerged weeds, addition of a labelled burndown product required. |
| Annual broadleaves | Goal 2XL (oxyfluorfen 2 lb. a.i./gal.) at 2-8 pt. in minimum 40 gal. water | See Pre-emergence section for details. |
| Annual broadleaves | GoalTender (oxyfluorfen 4 lb. a.i./gal.) at 2.5-4 pt. in minimum 40 gal. water | See Pre-emergence section for details. |

(Continued)

Herbicide Recommendations For Apple And Pear (*continued*)

| Weed Problem | Material And Rate Per Acre | Notes And Comments |
|--|---|---|
| Most annual grasses and broadleaf weeds and top kill of perennial weeds | Gramoxone SL 3.0 (paraquat 3 lb. a.i./gal.) at 1.7-2.7 pt. in minimum of 10 gal. water | Apply as directed spray to actively growing weeds. Repeat applications are necessary to give sustained control. Apply as a coarse spray. Always add NIS 0.25% v/v (2 pt./100 gal.) or crop oil 1% v/v (1 gal./100 gal.). Do not allow spray to contact leaves, fruit, or green stems. May be used for root sucker control. Do not exceed 5 applications per year. If application made during harvest, pick all fruit off of the ground prior to application. Restricted use pesticide. Only certified applicators can mix, load and apply. Not to be used by uncertified persons working under the supervision of a certified applicator. Applicators must complete an EPA- approved paraquat training every 3 years https://www.epa.gov/pesticide-work-safety/paraquat-dichloride-training-certified-applicators Containers under 120 gallons will have "closed-system" packaging to be used with a closed-transfer system. Harvest crop at normal maturity. |
| Most annual grasses and broadleaf weeds and top kill of biennial and perennial weeds | Homeplate (caprylic acid 45.14% + Capric acid 34.74%) at 3-9% solution if used alone or 1% solution when tank mixed | OMRI listed. Use includes vegetation burndown, directed and shielded sprays, and sucker control. May be used any time during the year and works best during warm and dry conditions. Reapply if rain falls within 3 hours of application. Avoid contact with desirable foliage and green bark. Sucker control: Apply before suckers become woody. |
| Annual grasses and broadleaves | Karmex DF (diuron 80% a.i.) at 4 lb. in 25-40 gal. water | See Pre-emergence section for details. |
| Annual grasses and broadleaves | Pindar GT (penoxsulam 0.083 lbs./gal. + oxyfluorfen 3.93 lbs./gal.) at 1.5-3 pt. | See Pre-emergence section for details. |
| Annual and perennial grasses | Poast 1.5EC (sethoxydim 1.5 lb. a.i./gal.) at 1.5-2.5 pt. in 25 gal. water | Apply as a directed spray to actively growing grass before they exceed maximum recommended heights. Always add crop oil 1.25% v/v. Do not exceed 2.5 pt. per application or 7.5 pt. per season. |
| Annual and perennial grasses and broadleaves | Rely 280 (glufosinate 2.34 lb. a.i./gal.) at 48-82 fl. oz. in minimum of 20 gal. water | Apply as a directed spray to actively growing weeds. Avoid spray drift or mist contact with green bark, stems, or foliage, as injury may occur. Only trunks with callused, mature brown bark should be sprayed unless protected by nonporous wraps, grow tubes, or waxed containers. Maximum rate is 246 fl. oz. per acre in a 12-month period. Add AMS to the spray tank if spray water is hard. Do not make spot or directed spray applications to tree trunks or to apple suckers, as tree injury may occur. Do not apply more than 246 fl. oz. of product per acre per calendar year or make more than 3 applications at a maximum rate of 82 fl. oz. per acre per year. Applications must be a minimum of 14 days apart. Sucker Control: apply when suckers are young, green, uncallused, and less than 12 inches in length. Apply a split application approximately 4 weeks apart at 56 fl. oz./ac. Avoid all contact with desirable tree parts other than mature bark, or serious damage can occur. Counts towards maximum cumulative amount and application limits. |
| Annuals and some perennial grasses and broadleaves | Roundup WeatherMax, Roundup PowerMax 5.5EC (glyphosate 5.5 lb. a.i./gal.) at 11 fl. oz.-3.3 qt. in 10-40 gal. water | Rate depends on weed species and stage of growth. See label for details. Do not exceed 7 qt. per acre per year. Apply as preplant broadcast application or in fall for control of roots and rhizomes of perennial weeds or as a directed spray or wiper application (33-100% solution) to actively growing weeds in established plantings. Always add AMS 8.5-17 lb./100 gal. in hard water or drought conditions. Do not allow spray to contact any part other than mature bark. Avoid application to suckers and recent pruning wounds. Does not provide residual control. Can be mixed with labeled pre-emergence herbicides. |

(Continued)

Herbicide Recommendations For Apple And Pear (*continued*)

| Weed Problem | Material And Rate Per Acre | Notes And Comments |
|---|--|---|
| Broadleaves and nutsedge | Sandea (halosulfuron-methyl 75% a.i.) 0.5-1 oz. in minimum of 15 gal. water | Apple and Pear east of the Mississippi: For best results, use a NIS with post-emergence applications. Avoid spray drift on tree foliage and fruit, and do not apply when temperatures exceed 85°F. Do not apply to trees established less than 1 year. Do not exceed 2 oz. per 12-month period. May not control ALS-resistant weeds. Make a single application using a minimum of 0.75 oz. per acre of Sandea when nutsedge is fully emerged at the 3- to 5-leaf stage. A second application may be made later in the season for secondary nutsedge emergence. See label for west of the Mississippi applications. |
| Annuals and perennial grasses and broadleaves | Scythe 4.2E (pelargonic acid 4.2 lb. a.i./gal.) at 3-10% spray mix | For contact nonselective control or burndown of a broad spectrum of actively growing weeds. Use low rate for annual weed control and high rates for maximum vegetative burndown. Use as a directed or shielded spray. Can be mixed with Roundup. Sucker Control: Apply only to unwanted vegetative parts and before suckers become woody. |
| Most annual and perennial grasses | Select Max with Inside Technology (clethodim 0.97 lb. a.i./gal.) at 9-16 fl. oz. | Apply post-emergence as a directed spray to young, actively growing grasses. Do not exceed 16 fl. oz./A in a single application or per season. A minimum 14-day interval is required for repeat applications. Add NIS at 0.25% v/v or COC/MSO at 1 qt./A or 1% v/v. |
| Annual and perennial broadleaf weeds | Starane Ultra (fluroxypyr 2.8 lb. a.i./gal.) at 0.4 - 1.4 pt. | Trees must be established at least 4 years. Do not apply during bloom. Apply uniformly with ground equipment in a minimum of 10 gallons of water per acre during calm periods and when air temperatures are between 50°F and 80°F. Avoid contact with foliage. Do not make more than two treatments per crop per year. Maximum individual application rate and total yearly rate 1.4 pt./A. |
| Most annual and perennial broadleaves | Stinger (clopyralid 3 lb. a.i./gal.) at 1/3-2/3 pt. in 10 gal. or more of total spray volume per acre | Apple only: Make one or two applications per crop year. Apply only to trees 1 year or older. Avoid direct contact with foliage, fruit, or tree trunks. Do not apply during bloom east of the Rocky Mountains. Do not exceed a total of 2/3 pt. per acre per crop year. |
| Annual and perennial grasses | Targa (quizalofop p-ethyl 0.88 lb. a.i./gal.) at 12 oz. in 10-40 gal. | Apply as a directed spray in a band extending out a minimum of 3 feet on each side of the tree row in 10-40 gallons of water/acre to control labelled grass weeds. DO NOT apply by aerial application. Two applications with a max rate of 24 oz. per season with application intervals at least 14 days apart to allow regrowth to occur. DO NOT apply within 14 days of anticipated bloom of crop. |
| Annual and perennial broadleaves | Treevix Powered by Kixor Herbicide (saflufenacil 0.7 lb. a.i.) at 1 oz. in 20-40 gal. water | Trees must be established for 9 months prior to application. May be applied as a single application or up to 3 times per season with a separation of 21 days between sprays. Do not exceed 3.0 oz. per acre per cropping season. Trunk shields should be used until trees have been established for 2-3 years. For optimum burndown, use with methylated seed oil (MSO), ammonium sulfate (AMS), or urea ammonium nitrate (UAN) adjuvant. Do not use an NIS as a substitute for MSO. Apply only when wind is 10 mph or less and blowing away from nontarget areas. Rainfast in 1 hour. Do not use in tree nurseries. |
| Annual and perennial broadleaves | Venue (pyraflufen ethyl 0.17 lb. a.i./gal.) at 3.0-4.0 fl. oz. plus other labeled herbicides in minimum of 20 gal. water | Not registered in all states. Apply as a directed spray during dormant period, prior to bloom, and post-harvest. Avoid contact with foliage and green bark. More effective on weeds less than 4 inches tall and 3 inches in diameter. Use higher rate and spray volume for larger weeds. Do not exceed 3 applications or 6.8 fl. oz. per acre per season. Allow a minimum of 30 days between applications. Adding COC or NIS is recommended. May be used for sucker growth control on the basal portion of trunks and root sprouts when tissue is young, immature and not hardened off. Avoid contact with green uncallused bark of young trees less than one year old unless protected by nonporous wraps or grow tubes. Do not allow spray to drift onto desirable fruit or foliage as damage will occur. May be mixed with 2, 4-D, glyphosate, or grass herbicides for enhanced control. Spray water pH needs to be less than 7.5. |

(Continued)

Herbicide Recommendations For Apple And Pear (*continued*)

| Weed Problem | Material And Rate Per Acre | Notes And Comments |
|---|---|--|
| Annual and perennial grasses and broad-leaves | Zalo (quizalofop-p-ethyl .23 lbs./gal. + glufosinate-ammonium 2.29 lbs./gal.) at 46 fl. oz. in 10-40 gal. | Avoid contact with desirable foliage, green stems, or exposed non-woody roots. Do not apply to suckers. Do not apply within 14 days of anticipated bloom. Rainfall or irrigation within 4 hours of application may reduce weed control. Avoid disturbing treated areas for at least 5 days before and 7 days following application. In addition to spray grade ammonium sulfate (AMS) at 3 lbs./A, use one of the following: crop oil concentrate (COC) adjuvants at 1% v/v; nonionic surfactants (NIS) with at least 90% ai at 0.25 to 0.5% v/v; MSO (methylated seed oil) adjuvants at 1% v/v; or HSOC (high surfactant oil concentrate) adjuvants at 0.5% v/v. Maximum single application rate 46 fl. oz./A. Two applications per calendar year, with at least 14 days separating applications. |

Herbicide Recommendations For Peach, Nectarine, Plum, And Cherry

| Weed Problem | Material And Rate Per Acre | Notes And Comments |
|---|--|--|
| Pre-emergence | | |
| Annual grasses and broadleaves | Alion (Indaziflam 1.67 lb. a.i./gal.) at 5.0-6.5 fl. oz. in minimum of 10 gal. water | Trees must be established at least 3 years after transplanting. Use lower rates as soil OM decreases. Avoid direct or indirect spray contact with crop foliage, green bark, roots, or fruit, as it may cause localized crop injury and death. Allow at least 30 days between applications. Do not exceed 10.3 fl. oz. per acre in a 12-month period. Do not apply to frozen ground. Do not apply within 25 feet of ponds, rivers, streams, or wetlands. Spot spraying is not recommended. Shake container well before use. |
| Annual and perennial grasses and broad-leaves | Casoron 4G (dichlobenil 4% a.i./lb.) at 100-150 lb. | For perennial weed control, apply to untilled ground over old weed growth from November 15 to Feb 15; alternately apply late fall or very early spring before May 15 and incorporate immediately. For annual weed control, surface apply. Shallow incorporation or sprinkler irrigation is recommended when application is made during periods of high temperatures. Do not apply until 4 weeks after transplanting. Use higher rate for perennial weed control. Annual maximum rate 150 lbs./A. |
| Annual broadleaves and suppression of grasses | Chateau SW (flumioxazin 51% a.i.) at 6-12 oz. Chateau EZ (flumioxazin 41.4% a.i.) at 6-12 oz. | The preferred application timing in the fall to maximize the potential for rainfall to activate and set the herbicide. Do not apply to trees established less than 1 year unless protected from spray contact by nonporous wraps. Make applications only to berms. Do not apply after bud break. For non-bearing trees, do not apply during the period after flowering through leaf drop unless shielded application equipment ensures that spray drift will not contact crop foliage. Do not apply to fine-textured soils. Do not exceed 2 applications in a growing season or make a sequential application within 30 days of the first application. Do not exceed 24 oz. per season Do not apply when plants are under stress. Do not apply within 300 yards of nondormant pears. Apply alone pre-emergence or tank mix with Gramoxone post-emergence with a crop oil 1% v/v or NIS 0.25% v/v. Do not incorporate. Do not allow drift to contact foliage or green bark. Trees may be transplanted 2 months after application. Minimum 30 days between applications. |
| Annual broadleaves and annual grasses | Chateau Complete (flumioxazin 30% a.i. + rimsulfuron 8.25%) at 12 oz. | Not registered in all states. The preferred application timing is in the fall to maximize the potential for rainfall to activate and set the herbicide. Do not apply to trees established less than 1 year. Do not apply after bud break. Do not apply within 300 yards of nondormant pears. Make applications only to berms. Avoid direct or indirect spray contact to foliage and green bark (nonbarked trunk with the exception of undesirable suckers). Do not apply to powdery soils or soils that are susceptible to wind displacement unless irrigation can be applied immediately after application. Do not mow treated areas between bud break and final harvest. Do not incorporate. Max seasonal rate 12 oz./A. |

(Continued)

Herbicide Recommendations For Peach, Nectarine, Plum, And Cherry (*continued*)

| Weed Problem | Material And Rate Per Acre | Notes And Comments |
|--|---|--|
| Annual broadleaves and suppression of grasses | Goal 2XL (oxyfluorfen 2 lb. a.i./gal.) at 5-8 pt. in minimum of 20 gal. water | Dormant Application Only: Effective both pre-emergence (5-8 pt.) and post-emergence (2-8 pt.) as a directed spray on weeds larger than 4 inches. Do not apply from bud swell until harvest completion. Can be mixed with other pre-emergence herbicides or with Roundup or Gramoxone. Do not exceed 8 pt. per year. |
| Annual broadleaves and suppression of grasses | GoalTender (oxyfluorfen 4 lb. a.i./gal.) at 2.5-4 pt. in minimum of 20 gal. water | Dormant Application Only: Effective both pre-emergence banded application (2.5-4 pt.) and post-emergence (1-4 pt.) as directed spray on weeds larger than 4 inches. Do not apply from bud swell until harvest completion. Can be mixed with other pre-emergence herbicides or with Roundup or Gramoxone. Do not exceed 3 pt. per year on a broadcast basis. |
| Annual grasses and broadleaves | Karmex DF diuron (80% a.i.) at 2-5 lb. in 25-40 gal. water | Peach Only: Effective both pre-emergence and post-emergence (minimum 70°F with high humidity). Apply under trees established at least 3 years. May be tank mixed with Sinbar (2 lb. each) in orchards established at least 2 years. Karmex/Sinbar can be applied in the spring before weeds emerge or after harvest in the fall. Do not exceed 1 application per year. |
| Annual and perennial grasses and certain broadleaves | Kerb SC (pronamide 35.6% a.i.) at 2.5-9.5 pt. in 40-50 gal. water. Rate depends on weed pressure and soil type. See table on label. | Apply as a directed spray in the fall after harvest prior to soil freeze-up. Rainfall or irrigation are required to activate. Do not apply more than 5 pt./A/year or make more than 1 application per year. Age restriction: Kerb SC may not be applied to (1) trees less than 1 year old, (2) fall-transplanted stock transplanted less than 1 year old, or (3) spring-transplanted stock transplanted less than 6 months. Some Special Local Needs Labels (FIFRA 24(c)) are labelled as Restricted Use Pesticides (RUP), whereas the national label remains a general use pesticide. |
| Annual grasses and broadleaves | Matrix FNV (rimsulfuron 25% a.i.) at 4 oz. in minimum of 10 gal. water | Apply only to crops that have been established for 1 full growing season and are in good health and vigor. Weeds are controlled for 60-90 days after application. Matrix burns down small actively growing weeds less than 1 inch tall. When weeds are present at application, a labeled burndown herbicide such as paraquat with an appropriate adjuvant improves control. Avoid direct or indirect contact with crop foliage or fruit, except undesirable suckers. Do not use Matrix FNV in a spray solution with a pH below 4.0 or above 8.0. Best results are obtained when the soil is moist at the time of application, and 0.5 inch of rainfall or sprinkler irrigation occurs within 2 weeks of application. |
| Annual grasses and broadleaves | Pindar GT (penoxsulam 0.083 lbs./gal. + oxyfluorfen 3.93 lbs./gal.) at 1.5-3 pt. | Apply only to trees established at least 4 years; see label for reset requirements. Use trunk guards to protect plants until adequate bark has developed. Application window beginning after harvest, up to bud swell. See label for tank partners for complete burndown. A single rainfall or sprinkler irrigation of 0.5 inches or more, or flood irrigation within 21 days after application, is necessary to activate. The addition of 1 quart per acre of crop oil concentrate or methylated seed oil, or 0.25% v/v of an 80% active nonionic surfactant labeled for application to growing food crops, is required for effective postemergence control of susceptible emerged weeds. Avoid direct contact with fruit trees. Single max application rate 3 pts./A; 4.5 pts./A max total for the year with 30 days between sequential applications at lower rates. |
| Annual grasses and broadleaves | Princep 4L (simazine 4 lb. a.i./ gal.) at 2-4 qt./A in min of 20 gals. water | Apply under trees established at least 2 years. Apply in spring before weeds emerge avoiding contact with fruit, foliage, or stems. Avoid use on sandy soils with OM <1%. 1 application per calendar year, with a max single application rate of 4 qt./A. Check label for state-specific restrictions on tank mixes. Replant restriction of 12 months following application. Plum, Sweet Cherry Only: Use only in MO and states east of the Mississippi River. Nectarine: Use restricted to CA. |
| Annual grasses and broadleaves | Sinbar WDG (terbacil 80% a.i.) at 0.5-4 lb. in minimum of 20 gal. water | Peach Only: Apply either in the spring before weeds emerge or during early stages of seedling growth or after harvest in the fall. Trees must be established at least 3 years. Do not contact foliage or fruit with spray or mist. Non-bearing (young, newly planted) Stone Fruits: Apply at 0.5-1 lb. Make the first application after a significant rainfall or irrigation event that allows the ground to settle around the base of the trees. Make 1-2 applications per season. Do not exceed 1 lb. per year. Do not use on soils with <1% OM. |

(Continued)

Herbicide Recommendations For Peach, Nectarine, Plum, And Cherry (*continued*)

| Weed Problem | Material And Rate Per Acre | Notes And Comments |
|---|---|---|
| Annual grasses and broadleaves and suppression of yellow nutsedge | Solicam DF (noraflurazon 78.6% a.i.) at 2-5 lb./A in min of 20 gals. water | See label for rate based on soil type. Apply a directed spray from fall to early spring before weeds emerge. Soil should be settled and firm. Rainfall or irrigation of 0.5 inch is needed within 4 weeks. Do not contact fruit or foliage. Do not apply after bud break on sandy loam soils. Check label for maximum amount allowed per year depending on soil type. Peach, Nectarine: minimum 6 months after planting before first application. Plum: minimum 12 months after planting before first application. Cherry: minimum 18 months after planting before first application. |
| Annual grasses and certain broadleaves | Surflan 4AS As Special Herbicide (oryzalin 4 lb. a.i./gal.) at 2-6 qt. in 20-40 gal. of water | No longer in production - supply extremely limited to not available. Make a single band of broadcast application to the ground beneath trees before weeds emerge. Apply alone to weed-free soil or post-emergence mixed with Roundup or Gramoxone. Minimum 0.5 inch rainfall or irrigation required for activation. Minimum 2.5 months between applications. Do not exceed 12 qt. per year. |
| Annual grasses and broadleaves | Treflan HFP 4EC (trifluralin 4 lb. a.i./gal.) at 1.5-4 pt. in 5-40 gal. water | Peach, Plum Only: Incorporate within 24 hours to reduce loss of activity. New Plantings: Apply 1.25-2 pt. and incorporate before transplanting. Established Plantings: Apply 2-4 pt. and incorporate prior to period of weed germination or after removal of weeds with tillage of herbicides. |
| Post-emergence | | |
| Annual broadleaves | Aim EC (carfentrazone 2 lb. a.i./gal.) at 2 fl. oz. in 20 gal. water | Apply any time during the season. Add NIS (2 pt./100 gal.) or COC (1 gal./100 gal.). Mix with Roundup or Gramoxone for broader weed control. Do not exceed 7.9 fl. oz. per year. Minimum 14 days between applications. Sucker Management: Apply when suckers are green. Do not allow spray to contact fruit, foliage, or green bark. |
| Annual and some perennial broadleaves | 2,4-D (Amine) at 1-4 pt. in 5-25 gal. water | Apply as directed spray to annuals 1-2 inches high and to perennials up to early bud stage. Do not allow spray to contact leaves, fruit, limbs or exposed roots of tree. Use coarse spray and low pressure to avoid drift. Do not apply during windy periods, when there is a temperature inversion or at extremely high temperatures. Non-bearing trees must be established at least 1 year. On bearing trees, do not apply during bloom and only after irrigation. Do not apply to bare ground. Use higher rate on perennial weeds. Do not exceed 2 applications per year. Maximum 75 days between applications. |
| Annual and some perennial broadleaves | Embed (2,4-D Choline) at 1-4 pt. in a min. of 10 gal. | Not registered in all states. Apply only to orchards that have been established for at least one year and are in vigorous growth condition. Apply to annual weeds when small and actively growing. Use higher rate on perennial weeds in bud to bloom stage. Do not allow spray to contact leaves, fruit, limbs or exposed roots of tree. Use coarse spray and low pressure to avoid drift. Do not exceed 2 applications per year. Maximum 75 days between applications, max total rate 8.42 pints/A. |
| Annual broadleaves | Bellum (mesotrione 4 lbs. a.i./gal.) at 3 -6 fl. oz. | Apply only to trees established for one full year and in good health and vigor. Use trunk guards to protect plants until adequate bark has developed. For pre-emergent residual control, apply before rainfall or irrigation. See label for tank mix partners. Apply in a spray volume of 10-40 gals./A. Total application rate in a 12-month period 12 fl. oz./A. Single maximum application rate 6 fl. oz./A. Three applications per year allowed in a 12-month period when using reduced rates, allowing at least 12 weeks between applications at the 6 fl. oz./A rate or 6 weeks between applications at the 6 fl. oz./A rate and subsequent applications of 3 fl. oz./A. For application to emerged weeds, the use of crop oil concentrate (COC) type adjuvant at 1% v/v or non-ionic surfactant (NIS) at 25% is advised. |
| Annual broadleaves and annual grasses | Chateau Complete (flumioxazin 30% a.i. + rimsulfuron 8.25%) at 12 oz. | Not registered in all states. See Pre-emergence section (Page???) for details If weeds are emerged at the time of application, apply with an adjuvant (0.25% v/v non-ionic surfactant or 1 qt./A crop oil concentrate). For complete control of emerged weeds, addition of a labelled burndown product required. |

(Continued)

Herbicide Recommendations For Peach, Nectarine, Plum, And Cherry (*continued*)

| Weed Problem | Material And Rate Per Acre | Notes And Comments |
|--|---|--|
| Most annual and perennial grasses | Fusilade DX (Fluazifop-P-butyl 2 lb. a.i./gal.) at 16-24 fl. oz. in 20-40 gal. water | Apply post-emergence as a directed spray avoiding contact with tree foliage to young actively growing grasses. Add a COC at 1% v/v (1 gal./100 gal.) or NIS at 0.25-0.5% v/v (1-2 qt./100 gal.) in the finished spray volume. Do not exceed 72 fl. oz. per acre per year. Maintain a minimum of 21 days between applications. Do not exceed a maximum of 3 applications per year. |
| Annual broadleaves | Goal 2XL (oxyfluorfen 2 lb. a.i./gal.) at 5-8 pt. in minimum of 40 gal. water | See Pre-emergence section for details |
| Annual broadleaves | GoalTender (oxyfluorfen 4 lb. a.i./gal.) at 2.5-4 pt. in minimum of 40 gal. water | See Pre-emergence section for details |
| Most annual grasses and broadleaf weeds and top kill of perennial weeds | Gramoxone (paraquat 3 lb. a.i./gal.) at 1.7-2.7 pt. in minimum of 10 gal. water | Apply as directed spray to actively growing weeds. Repeat applications are necessary to give sustained control. Apply as a coarse spray. Always add NIS 0.25% v/v or crop oil 1% v/v. Do not allow spray to contact leaves, fruit, or green stems. Do not exceed 3 applications per year. Restricted use pesticide. Only certified applicators can mix, load and apply. Not to be used by uncertified persons working under the supervision of a certified applicator. Applicators must complete an EPA-approved paraquat training every 3 years https://www.epa.gov/pesticide-worker-safety/paraquat-dichloride-training-certified-applicators Containers under 120 gallons will have "closed-system" packaging to be used with a closed-transfer system. |
| Most annual grasses and broadleaf weeds and top kill of biennial and perennial weeds | Homeplate (caprylic acid 45.14% + Capric acid 34.74%) at 3-9% solution if used alone or 1% solution when tank mixed | OMRI listed. Use includes vegetation burndown, directed and shielded sprays, and sucker control. May be used any time during the year and works best during warm and dry conditions. Reapply if rain falls within 3 hours of application. Avoid contact with desirable foliage and green bark. Sucker control: Apply before suckers become woody. |
| Annual grasses and broadleaves | Karmex DF diuron (80% a.i.) at 2-5 lb. in 25-40 gal. water | See Pre-emergence section for details. |
| Annual grasses and broadleaves | Pindar GT (penoxsulam 0.083 lbs./gal. + oxyfluorfen 3.93 lbs./gal.) at 1.5-3 pt. | See Pre-emergence section for details. |
| Annual and perennial grasses | Poast 1.5E (sethoxydim 1.5 lb. a.i./gal.) at 1.5-2.5 pt. in 25 gal. water | Apply as a directed spray to actively growing grasses before they exceed maximum recommended heights. Always add crop oil 1.25% v/v. Do not exceed 2.5 pt. per application or 5 pt. per season. Peach, plum, and nectarine are very tolerant to Poast and may be applied over the top of small non-bearing trees. |
| Annual and perennial grasses and broadleaves | Rely 280 (glufosinate 2.34 lb. a.i./gal.) at 48-82 fl. oz. in minimum of 20 gal. water | Apply as a directed spray to actively growing weeds. Avoid spray drift or mist contact with green bark, stems, or foliage, as injury may occur. Only trunks with callused, mature brown bark should be sprayed unless protected by nonporous wraps, grow tubes, or waxed containers. Add AMS to the spray tank if spray water is hard. Maximum rate is 164 fl. oz. per acre in a 12-month period. Do not make more than 2 applications at a maximum rate of 82 fl. oz. per acre per year. Do not make spot or directed spray applications to tree trunks or to suckers as tree injury may occur. Applications must be a minimum of 28 days apart. |

(Continued)

Herbicide Recommendations For Peach, Nectarine, Plum, And Cherry (*continued*)

| Weed Problem | Material And Rate Per Acre | Notes And Comments |
|--|--|--|
| Annuals and some perennial grasses and broadleaves | Roundup 5.5EC (glyphosate 5.5 lb. a.i./gal.) at 11 fl. oz.-3.3 qt. in 10-40 gal. water (many other formulations) | Rate depends on weed species and growth stage. See label for details. Apply as preplant broadcast application or in fall for control of roots and rhizomes of perennial weeds or as a directed spray or wiper application (33-100% solution) to actively growing weeds in established plantings. Always add AMS 8.5-17 lb./100 gal in hard water or drought conditions. Do not allow spray to contact any part other than mature bark. Avoid application to suckers and recent pruning wounds. Use extreme care to ensure that no part of peach tree is contacted with spray. Apply only near trees that have been planted in the orchard for 2 or more years. Does not provide residual control; can be mixed with labeled pre-emergence herbicides. |
| Annual and perennial grasses and broadleaves | Scythe 4.2E (pelargonic acid 4.2 lb. a.i./gal.) at 3-10% solution | For contact nonselective control or burndown of a broad spectrum of actively growing weeds. Use low rate for annual weed control and high rates for maximum vegetative burndown. Use as a directed or shielded spray. Can be mixed with Roundup. |
| Most annual and perennial broadleaves | Select Max with Inside Technology (clethodim 0.97 lb. a.i./gal.) at 12-16 fl. oz. | Apply post-emergence as a directed spray to young, actively growing grasses. Do not exceed 16 fl. oz./A in a single application or per season. A minimum 14-day interval is required for repeat applications. Add NIS at 0.25% v/v or COC/MSO at 1 qt./A or 1% v/v. |
| Most annual and perennial broadleaves | Stinger (clopyralid 3 lb. a.i./ gal.) at 1/3-2/3 pt. in 10 gal. or more of total spray volume | Make one or two applications per crop year. Apply only to trees 1 year or older. Avoid direct contact with foliage, fruit, or tree trunks. Do not apply during bloom. Do not exceed 2/3 pt. per acre per crop year. |
| Annual and perennial grasses | Targa (quizalofop p-ethyl 0.88 lb. a.i./gal.) at 12 oz./A in 10-40 gal. | Apply as a directed spray in a band extending out a minimum of 3 feet on each side of the tree row in 10-40 gallons of water/acre to control labelled grass weeds. DO NOT apply by aerial application. Two applications with a max rate of 24 oz. per season with application intervals at least 14 days apart to allow regrowth to occur. DO NOT apply within 14 days of anticipated bloom of crop. |
| Annual and perennial broadleaves | Venue (Pyraflufen ethyl 2% a.i.) at 0.7-4.0 fl. oz. plus other labeled herbicides in minimum of 20 gal. water | Apply as a directed spray during dormant period and prior to bloom. Avoid contact with foliage and green bark. More effective on weeds less than 4 inches tall and 3 inches in diameter. Use higher rate and spray volume for larger weeds. Do not exceed 3 applications or 6.8 fl. oz. per acre per season. Allow a minimum of 30 days between applications. Adding COC or NIS is recommended. May be used for sucker growth control on the basal portion of trunks and root sprouts when tissue is young, immature and not hardened off. Avoid contact with green uncallused bark of young trees less than one year old unless protected by nonporous wraps or grow tubes. Do not allow spray to drift onto desirable fruit or foliage as damage will occur. May be mixed with 2, 4-D, glyphosate, or grass herbicides for enhanced control. Spray water pH needs to be less than 7.5. |
| Annual and perennial grasses and broadleaves | Zalo (quizalofop-p-ethyl .23 lbs./gal. + glufosinate-ammonium 2.29 lbs./gal.) at 46 fl. oz. in 10-40 gal. | Avoid contact with desirable foliage, green stems, or exposed non-woody roots. Do not apply to suckers. Do not apply within 14 days of anticipated bloom. Rainfall or irrigation within 4 hours of application may reduce weed control. Avoid disturbing treated areas for at least 5 days before and 7 days following application. In addition to spray grade ammonium sulfate (AMS) at 3 lbs./A, use one of the following: crop oil concentrate (COC) adjuvants at 1% v/v; nonionic surfactants (NIS) with at least 90% ai at 0.25 to 0.5% v/v; MSO (methylated seed oil) adjuvants at 1% v/v; or HSOC (high surfactant oil concentrate) adjuvants at 0.5% v/v. Maximum single application rate 46 fl. oz./A. Two applications per calendar year, with at least 28 days separating applications. |

(Continued)

Herbicide Recommendations For Non-Bearing Fruit Trees Only

| Weed Problem | Material And Rate Per Acre | Notes And Comments |
|--|---|--|
| Pre- and Post-emergence | | |
| Annual broadleaves and yellow nutsedge | Broadloom (bentazon 4 lb. a.i./gal.) at 1.5-2 pt. in minimum of 20 gal. water | Not labelled in all states. Apply as a directed post-emergence spray. Always add COC 1% v/v. Avoid spraying stems, bark, or foliage. Do not exceed 2 pt. per application or exceed 4 pt. per season. |
| | Crew (isoxaben 0.50% + dithiopyr 0.25%) at 150 lbs. | Prior to application, the bed or soil surface should be smooth and free of plant and weed debris. Apply using a drop or rotary-type spreader designed to apply granular fertilizer. A single rainfall or sprinkler irrigation of 0.5 inch is often required to activate this product. Maximum 2 application per year, with at least 3 months between applications. |
| Most annual and perennial grasses | Fusilade DX (Fluazifop-P-butyl 2 lb. a.i./gal.) at 16-24 fl. oz. in 20-40 gal. water | For non-bearing apple and pear that will not be harvested within 1 year after application. Apply post-emergence as a directed spray, avoiding contact with tree foliage to young actively growing grasses. Add COC at 1% v/v (1 gal./100 gal.) or NIS at 0.25-0.5% v/v (1-2 qt./100 gal.) in the finished spray volume. Do not exceed 72 fl. oz. per acre or 3 applications per year. Maintain a minimum of 14 days between applications. |
| Most broadleaves | Gallery 75DF (isoxaben 75% a.i.) at 0.66-1.33 lb. in minimum of 10 gal. water | Apply in late summer to early fall; or pre-emergence in early spring prior to seed germination or immediately after cultivation. Do not apply to new transplants until soil has settled with no cracks present. Rainfall or irrigation (1/2 inch) is needed within 21 days of application. Not effective on germinated weeds. Minimum 60 days between applications. Maximum rate is 4 lb. per acre. |
| Annual grasses and certain broadleaves | Prowl 3.3EC (pendimethalin 3.3 lb. a.i./gal.) Short-term weed control: at 2.4 qt. in minimum of 20 gal. water Long-term weed control: 4.8 qt. in minimum of 20 gal. water | Do not apply if buds have started to swell. May be applied preplant incorporated, preplant surface, or pre-emergence. For best results, rain or irrigation is needed within 21 days of application. Not effective on germinated weeds. Do not allow spray to contact leaves, shoots, or buds. For new plantings, do not apply until soil has settled and no cracks are present. |
| Annual grasses and broadleaves | Reglone (diquat 2 lb. a.i./gal.) at 1.5-2 pt. in minimum of 15 gal. water | Apply post-emergence as a directed spray using a shield for contact burn of weeds. Always use NIS at 0.5% v/v. Complete coverage is essential for good control, and best control is on weeds 1"-6" in height. Can be used during site preparations and up to within 1 year of harvest. Do not allow contact with green stems, foliage, or fruits. Apply when wind speeds are 3-10 mph. Do not use for food or feed for 1 year after application. |
| Annual grasses and certain broadleaves | Snapshot 2.5TG (isoxaben + trifluralin 2.5% a.i.) at 100- 200 lb. | Apply pre-emergence on weed-free clean soil. For best results 1/2 inch rain or irrigation is needed within 3 days of application. Not effective on germinated seeds. Minimum 60 days between applications. Do not exceed 600 lb. per year. |

(Continued)

Herbicide Recommendations For Grape

| Weed Problem | Material And Rate Per Acre | Notes And Comments |
|---|--|--|
| Pre-emergence | | |
| Annual and perennial grasses and broad-leaves | Alion (indaziflam (1.67 lb. a.i. /gal.) at 5 fl. oz. | Only use in established vineyards at least 5 years after planting and on vines that exhibit normal growth and good vigor. Do not use on sandy soil or soils with 20% or more gravel content. Ensure 12 inches of soil barrier between the surface and the major portion of the root system. Age Restriction: Do not apply to vines less than 5 years old. |
| Annual and perennial grasses and broad-leaves | Casoron 4G (dichlobenil 4% a.i./lb.) at 100-150 lb. | For perennial weed control, apply to untilled ground over old weed growth from November 15 to Feb 15; alternately apply late fall or very early spring before May 15 and incorporate immediately. For annual weed control, surface apply. Shallow incorporation or sprinkler irrigation is recommended when application is made during periods of high temperatures. Do not apply until 4 weeks after transplanting. Use higher rate for perennial weed control. Annual maximum rate 150 lbs./A. |
| Annual broadleaves and suppression of grasses | Chateau SW (flumioxazin 51% a.i.) at 6-12 oz. in 15 gal. water minimum Chateau EZ (flumioxazin 41.4% a.i.) at 6-12 oz. in 15 gal. water minimum | Do not apply after bloom unless with a hooded or shielded application. Apply alone pre-emergence or tank mix with Roundup or Gramoxone post-emergence. Do not incorporate. Do not allow drift to contact foliage or green bark. Do not exceed 24 oz. per season. Minimum 30 days between applications. Should be applied to weed free surface. Also has post-emergence activity. Age Restriction: Do not apply to vines established less than 2 years unless they are trellised at least 3 ft. from the ground or are protected by nonporous wraps, grow tubes, or waxed containers. New plantings of "own-rooted varieties," such as Concord, should be planted so that all roots are a minimum 8 inches below the soil surface to be treated. In some situations, this may require hilling soil around newly planted vines so that the settled depth of the hill will be 4 to 5 inches above the vineyard floor. |
| Annual broadleaves and annual grasses | Chateau Complete (flumioxazin 30% a.i. + rimsulfuron 8.25%) at 6-12 oz. | Not registered in all states. Do not apply more than 12 oz. per acre per year. Do not apply to crops established less than one year. If weeds are emerged at the time of application, apply with an adjuvant (0.25% v/v non-ionic surfactant or 1 qt./A crop oil concentrate). For complete control of emerged weeds, addition of a labelled burndown product required. |
| Annual grasses and broadleaves | Devrinol 2-XT (napropamide 2 lb. a.i./gal.) at 2 gal. | Apply from late fall (prior to soil freezing) to early spring (prior to weed emergence). Apply to a weed-free soil surface or tank mix with a suitable post-emergence herbicide. May be applied to newly planted and established crops. Do not exceed 2 gal per acre per crop cycle. |
| Broadleaves | Gallery 75 DF (isoxaben 0.75 lb. a.i./lb) at 0.66-1.33 lb. Gallery SC is for non-bearing only | Apply any time prior to germination of target weeds or immediately after cultivation. Specific Use Restrictions: <ul style="list-style-type: none"> Do not apply within 165 days of harvest. Do not apply Gallery 75 Dry Flowable more than twice per crop year (harvest to harvest) up to a maximum total of 1.33 lb. of product per acre per crop year. |
| Annual broadleaves and suppression of grasses | Goal 2XL (oxyfluorfen 2 lb. a.i./gal./) at 5-8 pt. in minimum of 20 gal. water | Dormant Application Only: Effective both pre-emergence (5-8 pt.) and post-emergence (2-8 pt.) as a directed spray on weeds less than 4 inches tall. Do not apply from bud swell to harvest. Can be mixed with other pre-emergence herbicides, or with Roundup or Gramoxone. Do not exceed 8 pt. per year. Age Restriction: Do not apply to grapes established less than 3 years unless vines are on a trellis wire a minimum of 3 feet above ground. |
| Annual broadleaves and suppression of grasses | GoalTender (oxyfluorfen 4 lb. a.i./gal.) at 2.5-4 pt. in minimum of 20 gal. water | Dormant Application Only: Effective both pre-emergence as a banded application (2.5-4 pt.) and post-emergence (1-4 pt.) as a directed spray on weeds less than 4 inches tall. Do not apply from bud swell to harvest. Can be mixed with other pre-emergence herbicides, or with Roundup or Gramoxone. Do not exceed 4 pt per year on a band application basis. Age Restriction: Do not apply to grapes established less than 3 years unless vines are on a trellis wire a minimum of 3 ft. above ground. |

(Continued)

Herbicide Recommendations For Grape (*continued*)

| Weed Problem | Material And Rate Per Acre | Notes And Comments |
|---|---|--|
| Annual grasses and broadleaves | Karmex DF (diuron 80% a.i.) at 2-6 lb. in 25-40 gal. water | Age Restriction: Use on vineyards established at least 3 years and trunks at least 1.5 inches in diameter. Apply as a directed spray to soil under trellis in early spring prior to weed germination. Do not exceed 1 application per year. On soils low in organic matter (1-2%), severe injury may result if heavy rainfall occurs soon after treatment. |
| Annual and perennial grasses and certain broadleaves | Kerb SC (pronamide 35.6% a.i.) at 2.5-9.5 pt. in 40-50 gal. water. Rate depends on weed pressure and soil type. See table on label. | Apply as a directed spray in the fall after harvest prior to freeze-up, or in early winter when temperatures are below 55° F. Rainfall or irrigation are required to activate. Do not exceed 1 application per year or exceed 9.5 pt./A/year. Some Special Local Needs Labels (FIFRA 24(c)) are labelled as Restricted Use Pesticides (RUP), whereas the national label remains a general use pesticide. Age restriction: Do not apply to vines less than 1 year old. |
| Annual grasses and broadleaves | Matrix FVN or SG (rimsulfuron 25% a.i.) at 4 oz. in a minimum of 10 gal. water | Apply as a banded application to the base of the vines. Best results are obtained when the soil is moist at the time of application, and 1/2 inch of rainfall or sprinkler irrigation occurs within 2 weeks after application. Age Restriction: Do not apply to vines established less than one year. |
| Annual and perennial grasses and broadleaves | Mission (flazasulfuron 25% a.i.) at 2.14-2.85 oz. in 15-50 gal. water | Apply as a directed spray to soil beneath vines to prevent injury to foliage and bark of young vines. You must use a protective for third year vines to minimize potential injury. Age Restriction: Apply to grapes established 3 years or more. |
| Annual grasses and broadleaves | Princep 4L (simazine 4 lb. a.i./ gal.) at 2-4 qt. in 25-40 gal. of water | Age Restriction: Use on vineyards established at least 3 years. Apply to soil under trellis between harvest and early spring before weeds emerge. Apply alone to weed-free soil or tank mix with Roundup or Gramoxone. Do not exceed 1 application per year. |
| Annual grasses and certain broadleaf weeds | Prowl 3.3 EC (pendimethalin 3.3 lb. a.i./ gal.) at 2.4-4.8 qt. | Non-Bearing Only. Use rates of this product vary by soil texture and organic matter. Most effective in controlling weeds mechanically incorporated or when incorporated into the weed germination zone by adequate rainfall or overhead irrigation after application. Apply only to dormant grapevines. - DO NOT apply if buds have started to swell. Application after buds have started to swell may result in leaf distortion. - DO NOT apply to newly transplanted trees or vines until ground has settled and no cracks are present. |
| Annual grasses and certain broadleaves | Prowl H2O (pendimethalin 3.8 lb. a.i./ gal.) at 3.2-6.3 qt. in minimum of 20 gal. water | In bearing vineyards, this product may be applied any time after fall harvest, during winter dormancy, and in the spring. In non-bearing vineyards this product may be applied preplant incorporated, pre-plant surface, or pre-emergence. For best results, rain or irrigation is needed within 21 days of application. Not effective on germinated weeds. Do not allow spray to contact leaves, shoots, or buds. For new plantings, do not apply until soil has settled and no cracks are present. |
| Annual grasses and certain broadleaves | Snapshot 2.5TG (isoxaben+trifluralin 2.5% a.i.) at 100-200 lb. | Non-bearing Only: May only be used on crops that will not be harvested within 1 year of application. Rainfall or irrigation of 0.5 inch is needed within 3 days of application. Not effective on germinated weeds. Minimum 60 days between applications. Do not exceed 600 lb. per 12-month period. Do not apply to new transplants until soil has settled and with no cracks. |
| Annual grasses and broadleaves and suppression of yellow nutsedge | Solicam DF (norflurazon 78.6% a.i.) at 1.25-5 lb. in minimum of 20 gal. water | Apply as a directed spray to settled and firm soil from fall to early spring before weeds emerge. Rainfall or irrigation is needed within 4 weeks of application. Do not contact fruit or foliage. Do not apply after bud break on sandy loam or other coarse-textured soils. Check label for maximum amount allowed per year depending on soil type. Age Restriction: Allow a minimum of 24 months after planting before first application. |

(Continued)

Herbicide Recommendations For Grape (*continued*)

| Weed Problem | Material And Rate Per Acre | Notes And Comments |
|---|--|---|
| Annual grasses and certain broadleaves | Surflan As Specialty Herbicide (oryzalin 4 lb. a.i./ gal.) at 2-6 qt. in 20-40 gal. of water | No longer in production - supply extremely limited to not available. Make a single band or broadcast application to the ground beneath vines before weeds emerge. Apply alone to weed-free soil or post-emergence mixed with Roundup or Gramoxone. Rainfall or irrigation (0.5 inch) is required for activation. Minimum 2.5 months between applications. Do not exceed 12 qt. per year. |
| Annual grasses and broadleaves | Treflan HFP 4EC (trifluralin 4 lb. a.i./ gal.) at 1-4 pt. in 5-40 gal. water | In a new planting, apply 1-4 pt. and incorporate within 24 hours. In an established planting, apply 2-4 pt. prior to weed germination or immediately after removal of weeds with tillage or other herbicides and incorporate within 24 hours. |
| Annual and perennial broadleaves | Trellis (isoxaben 75% a.i.) at 0.67-1.33 lb. in minimum of 10 gal. water Trellis SC (isoxaben 4.16 lb. a.i./gallon) at 16-31 oz. | Non-bearing: Apply any time before target weeds germinate or immediately after cultivation. Bearing: Apply before target weeds germinate or immediately after cultivation. Do not exceed 2 applications per crop year or exceed 1.33 lb. (1.0 lb isoxaben) per acre per crop year. Do not apply Trellis SC more than twice per crop year (harvest to harvest) up to a maximum total of 1.0 lb. a.i. (31 fl. oz.) per crop year. Do not apply Trellis SC to newly transplanted vines until soil has been settled and no cracks are present of plant injury may occur. |
| Annual and perennial grasses and broadleaves | Zeus Prime XC (carfentrazone- ethyl 3.5% and sulfentrazone 31.8% a.i.) at 7.7-15.2 fl. oz. per acre in minimum of 10 gal. water | Apply as a broadcast or banded soil application directed to the base of the vines. If weeds are present, tank mix with a post-emergence herbicide to eliminate emerged weeds. Apply a single broadcast application at 15.2 fl. oz. per acre (0.41 lb. a.i./acre). May be applied as a banded treatment twice per year. Do not exceed 15.2 fl. oz. (0.41 lb. a.i.)/acre/year. Minimum of 60 days between applications. Do not apply after bud break except with hooded or shielded sprayer. Age Restriction: Apply to crops that have been growing for at least 2 years and are in good condition. |
| Annual and perennial grasses and broadleaves | Zeus XC (sulfentrazone 39.6% a.i.) at 8-12 fl. oz. per acre in a minimum of 10 gal. water | Apply as a broadcast or banded soil application directed to the base of the vines. If weeds are present, tank mix with a post emergence herbicide to eliminate emerged weeds. Apply a single broadcast application at 8-12 fl. oz./acre (0.25-0.375 lb. a.i./acre). May be applied as a banded treatment twice per year. Minimum 60 days between applications. Do not exceed 12 fl. oz. (0.375 lb. a.i.)/acre/year. Do not apply after bud break except with hooded or shielded sprayer. Age Restriction: Apply to crops that have been growing for at least 3 years and are in good condition. |
| Post-emergence | | |
| Annual broadleaves | Aim EC (carfentrazone 2 lb. a.i./gal.) at 1-2 fl. oz. in 20 gal. water | Apply any time during the season as a post-emergence directed spray or as a hooded spray treatment. Always add NIS at 0.5% v/v or COC at 1% v/v. Mix with Roundup or Gramoxone or labeled pre-emergence herbicides for broader weed control. Do not exceed 7.9 fl. oz. per year. Minimum 14 days between applications. Sucker Management: Apply when suckers are green. Do not allow spray to contact desirable fruit, foliage, or green bark. |
| Annual broadleaves and suppression of grasses | Chateau SW (flumioxazin 51% a.i.) at 6-12 oz. in 15 gal. water minimum Chateau EZ (flumioxazin 41.4% a.i.) at 6-12 oz. in 15 gal. water minimum | Include an adjuvant (0.25% v/v non-ionic surfactant or 1 qt./A crop oil concentrate) for post-emergence use. See Pre-emergence section for other notes and restrictions. |

(Continued)

Herbicide Recommendations For Grape (*continued*)

| Weed Problem | Material And Rate Per Acre | Notes And Comments |
|--|---|---|
| Annual broadleaves and grasses | Chateau Complete (flumioxazin 30% a.i. + rimsulfuron 8.25%) at 6-12 oz. | Not registered in all states. Do not apply more than 12 oz. per acre per year. Do not apply to crops established less than one year. If weeds are emerged at the time of application, apply with an adjuvant (0.25% v/v non-ionic surfactant or 1 qt./A crop oil concentrate). For complete control of emerged weeds, addition of a labelled burndown product required. See Pre-emergence section for details. |
| Most annual and perennial grasses | Fusilade DX (fluazifop-p-butyl 2 lb. a.i./gal.) at 16-24 fl. oz. in 25 gal. water | Apply as a directed spray to actively growing grasses before tillering. Always add COC at 0.5-1% v/v or NIS at 0.25-0.5% v/v. Avoid contact with grape foliage. Rainfast in 1 hour. Do not exceed 24 fl. oz. per application per acre or exceed 72 fl. oz. per acre per year. Minimum 14 days between applications and a maximum of 3 applications per year. |
| Annual grasses and broadleaves | Gamma (tiafenacil 0.7 lb. a.i./lb.) 0.5-1.5 oz. in minimum of 10 gal. water | Not labelled in all states. Apply as a directed spray. Use a methylated seed oil (MSO) product that contains modified vegetable oil with at least 15% surfactant emulsifier or reduced performance can occur. MSO should be applied at a concentration equal to 1% v/v (1 gallon per 100 gallons spray carrier) of the final spray volume. Do not allow contact with green stems or foliage. Do not reapply within 14 days. Do not apply within 7 days of harvest. Age restriction: Do not apply to grapes established less than 2 years |
| Annual broadleaves | Goal 2XL (oxyfluorfen 2 lb. a.i./gal.) at 2-8 pt. in minimum of 20 gal. water | See Pre-emergence section for details. |
| Annual broadleaves | GoalTender (oxyfluorfen 4 lb. a.i./gal.) at 1-4 pt. in minimum of 40 gal. water | See Pre-emergence section for details. |
| Most annual grasses and broadleaves and top kill of perennial weeds | Gramoxone SL 3.0 (paraquat 3 lb. a.i./gal.) at 1.7-2.7 pt. in minimum of 10 gal. water | Apply as directed spray to actively growing weeds. Repeat applications are necessary to give sustained control. Avoid contact with desired new shoots, fruit, or foliage. Apply as a coarse spray. Always add NIS at 0.25% v/v or COC at 1% v/v. Best results with flat fan nozzles. Do not exceed 5 applications per year. Harvest at normal crop maturity. Sucker Management: Apply when suckers are less than 8 inches tall. Do not allow spray to contact desirable fruit, foliage, or green bark. Restricted use pesticide. Only certified applicators can mix, load and apply. Not to be used by uncertified persons working under the supervision of a certified applicator. Applicators must complete an EPA- approved paraquat training every 3 years https://www.epa.gov/pesticide-worker-safety/paraquat-dichloride-training-certified-applicators Containers under 120 gallons will have "closed-system" packaging to be used with a closed-transfer system. |
| Most annual grasses and broadleaf weeds and top kill of biennial and perennial weeds | Homeplate (caprylic acid 45.14% + Capric acid 34.74%) at 3-9% solution if used alone or 1% solution when tank mixed | OMRI listed. Use includes vegetation burndown, directed and shielded sprays, and sucker control. May be used any time during the year and works best during warm and dry conditions. Reapply if rain falls within 3 hours of application. Avoid contact with desirable foliage and green bark. Sucker control: Apply before suckers become woody. |
| Annual and perennial grasses and broadleaves | Mission (flazasulfuron 25% a.i.) at 2.14-2.85 oz. in 15-50 gal. water | Apply to weeds less than 4 inches tall and before tillering of grasses in sufficient volume to get thorough coverage. Always use an adjuvant. Do not exceed 2 applications at the 2.85 oz. rate per acre per year. |

(Continued)

Herbicide Recommendations For Grape (*continued*)

| Weed Problem | Material And Rate Per Acre | Notes And Comments |
|--|---|---|
| Annual and perennial grasses | Poast 1.5EC (sethoxydim 1.5 lb. a.i./gal.) at 1.5-2.5 pt. in minimum of 5 gal. water | Apply as a directed spray to actively growing grasses before tillering. Always add COC at 1% v/v. Do not exceed 2.5 pt. per application or exceed 5 pt. per season. |
| Annual grasses and broadleaves | Reglone (diquat 2 lb. a.i./gal.) at 1.5-2 pt. in minimum 15 gals. of water | Non-bearing Only: May only be used on crops that will not be harvest within 1 year of application. Apply as a directed spray using a shield for contact burn of weeds. Apply when wind speed is 3-10 miles per hour. Complete coverage is essential for good control. Always use NIS at 0.5% v/v. Can be used during site preparation and up to 1 year of harvest. Do not allow contact with green stems, foliage or fruits. |
| Annual and perennial grasses and broadleaves | Rely 280 (glufosinate 24.5% a.i. (2.34 lb./gal.)) at 48-82 fl. oz. in minimum of 15 gal. water | Spray only trunks with callused, mature, brown bark unless protected from spray contact by nonporous wraps, grow tubes, or waxed containers. Apply as a directed spray to actively growing weeds. Add AMS to the spray tank if spray water is hard. Do not exceed 246 fl. oz. per acre per year. Do not make more than 3 applications at a maximum rate of 82 fl. oz. per acre per year. For spot application, mix 1.7 fl. oz./gal. |
| Annuals and some perennial grasses and broadleaves | Roundup Weather-Max 5.5EC (glyphosate 5.5 lb. a.i./gal.) at 11 fl. oz. to 3.3 qt. in 10-40 gal. water | Apply as a directed spray or wiper application to actively growing weeds in established plantings. Rate depends on equipment used, weed species, and stage of growth. See label for details. Always add ammonium sulfate at 8.5-17 lb./100 gal. in hard water or drought conditions (see label). Do not allow spray to contact any part other than mature bark. Can be mixed with labeled pre-emergence herbicides. |
| Annual and perennial grasses and broadleaves | Scythe 4.2E (pelargonic acid 4.2 lb. a.i./gal.) at 3-10% spray solution | For contact nonselective control or burndown of a broad spectrum of actively growing weeds. Use low rate for annual weed control and high rate for maximum vegetative burndown. Use as a directed spray or shielded spray. Can be mixed with Roundup. |
| Most annual and perennial grasses | Select Max with Inside Technology (clethodim 0.97 lb. a.i./gal.) at 9-16 fl. oz. | Non-bearing Only: May only be used on crops that will not be harvested within 1 year of application. Apply as a directed spray to actively growing grasses before tillering. Do not use if rain is expected within 1 hour. Always add NIS at 0.25% v/v. Do not use COC. May be applied as a spot treatment at 0.32-0.64 fl. oz. per gal Do not exceed 32 fl. oz. per year. |
| Annual broadleaves | Venue (pyraflufen ethyl 0.17 lb. a.i./gal.) at 3.0-4.0 fl. oz. in minimum of 20 gal. water | Not registered in all states. Use as a directed spray from dormancy, prior to bloom and postharvest. Repeat if needed. Keep off green stems and foliage. The addition of COC at 1-2% is recommended. Do not exceed 6.8 fl. oz. per acre per year or 3 applications per growing season. |

Herbicide Recommendations For Highbush Blueberry (*continued*)

| Weed Problem | Material And Rate Per Acre | Notes And Comments |
|--------------------------------|---|---|
| Pre-emergence | | |
| Annual grasses and broadleaves | Alion (Indaziflam 1.67 lb. a.i./gal.) at 3.5 - 10.0 fl. oz. | Rate dependent on soil organic matter content. Do not apply more than a total of 7.0 fl. oz. product/A (0.09 lb. ai/A) per year on soils containing < 1 % organic matter content, or 10.0 fl. oz. product/A (0.13 lb. ai/A) per year on soils containing ≥ 1 % organic matter content in a 12 month period when used in any highbush blueberry planting. Do not allow spray to contact green stems, foliage, flowers, or berries or unacceptable injury may occur. Only apply to soil as a dormant application in late fall through early spring before bud swell. Apply as a directed application to the soil beneath the bushes. When making more than one application per year, allow a minimum of 90 days between applications. |

(Continued)

Herbicide Recommendations For Highbush Blueberry (*continued*)

| Weed Problem | Material And Rate Per Acre | Notes And Comments |
|--|--|--|
| Annual grasses and broadleaves | Callisto (mesotrione 4 lb. a.i./ gal.) at 3.0-6.0 fl. oz. | Apply pre-emergence or early post-emergence. For improved post-emergence control, apply split applications at 3.0 fl. oz. at least 14 days apart. Do not exceed 2 applications per year or exceed 6 fl. o.z per year. Do not apply after the onset of bloom. Include a COC tolerated by blueberries if applied post-emergence to weeds. |
| Annual and perennial grasses and broadleaves | Casoron 4G (dichlobenil 4% a.i.) at 100-150 lb. | For perennial weed control, apply to untilled ground over old weed growth from November 15 to Feb 15; alternately apply late fall or very early spring before May 15 and incorporate immediately. For annual weed control, surface apply. Shallow incorporation or sprinkler irrigation is recommended when application is made during periods of high temperatures. Do not apply until 4 weeks after transplanting. Use higher rate for perennial weed control. Annual maximum rate 150 lbs./A. |
| Annual broadleaves and suppression of grasses | Chateau SW (flumioxazin 51% a.i.) at 6-12 oz. Chateau EZ (flumioxazin 41.4% a.i.) at 6-12 oz. | Apply as a uniform band directed at the base of the bush. Avoid direct spray contact to foliage or green bark. Preferred application timing is in the fall. Do not exceed 6 oz. per acre per application. Do not make a sequential application within 30 days of the first application. Do not exceed 12 oz. per acre per 12-month period. Do not apply to farm alleys or roads where traffic may result in treated dust settling onto crops or other desirable vegetation. Do not mow treated areas between bud break and final harvest. Dust created by mowing may drift onto desirable vegetation resulting in injury. Do not apply within 300 yards of non-dormant pome or stone fruit. Age Restriction: Do not apply to plants less than 2 years old unless they are protected by nonporous wrap, grow tubes or waxed containers. |
| Annual broadleaves and suppression of grasses | Chateau Complete (flumioxazin 30% a.i. + rimsulfuron 8.25%) at 6-12 oz. | Not registered in all states. Do not apply more than 12 oz. per acre per year. Do not apply to crops established less than one year. Do not apply to plants less than 2 years old unless they are protected by nonporous wrap, grow tubes or waxed containers. If weeds are emerged at the time of application, apply with an adjuvant (0.25% v/v non-ionic surfactant or 1 qt./A crop oil concentrate). For complete control of emerged weeds, addition of a labelled burndown product required. See Pre-emergence section for details. |
| Annual grasses and broadleaves | Devrinol 2-XT (napropamide 2 lb. a.i./gal.) at 2 gal. per acre | Apply to a weed-free soil surface or tank mix with a suitable post-emergence herbicide. May be applied to newly planted and newly established crops. Do not exceed 2 gal per acre per crop cycle. |
| Most broadleaves | Gallery 75DF (isoxaben 75% a.i.) at 0.66-1.33 lb. in minimum of 10 gal. water Gallery SC (isoxaben 4.16 lb. a.i./gal.) at 16-31 oz. in minimum of 10 gal. water | Non-Bearing Only: May only be used on crops that will not be harvested within 1 year of application. Apply in late summer to early fall; or in early spring prior to weed germination or anytime immediately after cultivation. Do not apply to new transplants until soil has settled with no cracks present. Rainfall or irrigation of 1/2 inch is needed within 21 days of application. Minimum of 60 days between applications. Do not exceed 4 lb. per acre per 12-month period. Non-Bearing Only: Make a single application prior to germination of target weeds or immediately after cultivation. May only be used on crops that will not be harvested within 1 year of application. Do not exceed 1.0 lb. a.i./A/yr. |
| Annual grasses and broadleaves | Karmex DF (diuron 80% a.i.) at 1.5-4 lb. in 25-40 gal. water | Selected states only. Age Restriction: Use only in fields established at least 1 year. Apply as a band treatment at the base of bushes. Do not apply to exposed roots. For AR and MO Only: Apply 1.5-2 lb. in spring and repeat after harvest in the fall. Always add NIS at 0.25% v/v. For IN and OH Only: Apply 2-4 lb. in late spring. Alternatively, apply 2 lb. in fall and repeat in spring. |
| Annual and perennial grasses and certain broadleaves | Kerb SC (pronamide 35.6% a.i.) at 2.5-5.0 pt. in 20-50 gal. water. Rate depends on weed pressure. See table on label. | Apply as a directed spray in the fall after harvest prior to freeze-up, or in early winter when temperatures are below 55° F. Rainfall or irrigation are required to activate. Do not exceed 1 application per year or 5.0 pt./A/year. Some Special Local Needs Labels (FIFRA 24(c)) are labelled as Restricted Use Pesticides (RUP), whereas the national label remains a general use pesticide. Age restriction: Do not apply to newly transplanted blueberries until roots are well established. |
| | | <i>(Continued)</i> |

Herbicide Recommendations For Highbush Blueberry (*continued*)

| Weed Problem | Material And Rate Per Acre | Notes And Comments |
|---|---|--|
| Annual and perennial grasses and broad-leaves | Princep 4L (simazine 4 lb. a.i./ gal.) at 2-4 qt. in minimum of 40 gal. water | Apply in spring before weeds emerge and before canes leaf out or make a split application of 2 qt. in spring plus 2 qt. in fall. For plants established less than 6 months, apply half the above rate. Do not apply when fruit is present, or illegal residues may result. |
| Annual broadleaf weeds and nutsedge | Sandea (halosulfuron 75%) at 0.5-1 oz. in minimum of 15 gal. water | Apply with ground equipment as a broadcast application to the ground on either side of the row. Apply as a single or sequential application depending on weed pressure. Mix with a post-emergence broad-spectrum-type herbicide to maximize and enhance the spectrum of control If small weeds are present. For post-emergence nutsedge control, make a single application when nutsedge has fully emerged, or make 2 sequential applications. Apply the first to the initial nutsedge flush when it has reached the 3-5-leaf stage. If a second application is needed, it can be applied later in the season. Avoid contact with blueberry bushes. Minimum of 45 days between applications. Do not exceed 2 oz. per acre per year. Age Restriction: Do not apply to plants established less than 1 year. |
| Annual grasses and broadleaves | Sinbar WDG (terbacil 80% a.i.) at 2-3 lb. in minimum of 25 gal. water | Age Restriction: Use only on plantings established at least 1 year. Best results when applied shortly before or after weed emergence. Avoid contact of foliage or fruit with spray or mist. Apply either in the spring or after harvest in the fall before weeds emerge or during early stage of seedling regrowth. Do not use on soils where roots are exposed. Use rate varies by soil type. Do not use on sand or loamy sand with 1-3% organic matter. |
| Annual grasses and certain broadleaves | Snapshot 2.5TG (isoxaben+trifluralin 2.5% a.i.) at 100-200 lb. | Non-bearing Only: May only be used on crops that will not be harvested within 1 year of application. Do not apply to new transplants until soil has settled. For best results, 1/2 inch of rain or irrigation is needed within 3 days of application. Not effective on germinated weeds. Minimum 60 days between applications. Do not exceed 600 lb. per 12-month period. |
| Annual grasses and broadleaves and suppression of yellow nutsedge | Solicam DF (norflurazon 78.6% a.i.) at 2.5-5 lb. in minimum of 20 gal. water | Apply as a directed spray to settled and firm soil from fall to early spring before weeds emerge. Rainfall or irrigation of 1/2 inch is needed within 4 weeks. Do not contact fruit or foliage. Do not apply after bud break on sandy loam soils. Check label for maximum amount allowed per year depending on soil type. Minimum 6 months after planting before first application. |
| Annual grasses and broadleaves | Surflan As Specialty Herbicide (oryzalin 4 lb. a.i./gal.) at 2-6 qt. in 20-40 gal. water | No longer in production - supply extremely limited to not available. Make a single band or broadcast application to the ground beneath canes weeds emerge. Apply alone to weed-free soil or post-emergence mixed with Roundup or Gramoxone. Rainfall or irrigation of 1/2 inch is required for activation. Minimum 2.5 months between applications. Do not exceed 12 qt. per year. |
| Annual and perennial broadleaves | Trellis (isoxaben 75% a.i.) at 0.67-1.33 lb. in minimum of 10 gal. water Trellis SC (isoxaben 4.16 lb. a.i./gal.) at 16-31 oz. | Non-Bearing Only: Apply before target weeds germinate or immediately after cultivation. Do not apply Trellis SC more than twice per crop year (harvest to harvest) up to a maximum total of 1.0 lb. a.i. (31 fl. oz.) per crop year. Do not apply to newly transplanted bushes until soil has been settle and no cracks are present of plant injury may occur. |
| Annual grasses and broadleaves | Velpar 2L (hexazinone 2 lb. a.i./gal.) at 0.5-1 gal. in 20 gal. water | Apply to pruned blueberries in the spring before leaf emergence as a directed soil application. Some clones are susceptible to injury. Age Restriction: Use on plantings established at least 3 years. |

(Continued)

Herbicide Recommendations For Highbush Blueberry (*continued*)

| Weed Problem | Material And Rate Per Acre | Notes And Comments |
|---|---|---|
| Annual and perennial grasses and broad-leaves | Zeus Prime XC (carfentrazone- ethyl 3.5% and sulfentrazone 31.8% a.i.) at 7.7-15.2 fl. oz. per acre in minimum of 10 gal. water | Apply as a broadcast or banded soil application directed to the base of the trunks of bushes or vines. If weeds are present, tank mix with a post-emergence herbicide to eliminate emerged weeds. Apply a single broadcast application at 15.2 fl. oz. per acre (0.41 lb. a.i./acre). May be applied as a banded treatment twice per year. Do not exceed 15.2 fl. oz. (0.41 lb. a.i.)/acre/year. Minimum of 60 days between applications. Do not apply after bud break except with hooded or shielded sprayer. Age Restriction: Apply to crops that have been growing for at least 2 years and are in good condition. |
| Annual and perennial grasses and broad-leaves | Zeus XC (sulfentrazone 39.6% a.i.) at 8-12 fl. oz. per acre in a minimum of 10 gal. water | Apply as a broadcast or banded soil application directed to the base of the trunk of bushes. If weeds are present, tank mix with a post-emergence herbicide to eliminate emerged weeds. Apply a single broadcast application at 8-12 fl. oz. per acre (0.25-0.375 lb. a.i./acre). May be applied as a banded treatment twice per year. Minimum of 60 days between applications. Do not exceed 12 fl. oz. (0.375 lb. a.i.)/acre/year. Do not apply after bud break except with hooded or shielded sprayer. Age Restriction: Apply to crops that have been growing for at least 3 years and are in good condition. |
| Post-emergence | | |
| Annual broadleaves | Aim EC (carfentrazone 2 lb. a.i./gal.) at 1-2 fl. oz. in 20 gal. water | Apply broadcast at base of canes during dormant stage or with hooded shields between rows during growing season. Always add NIS at 0.25% v/v or COC at 1% v/v. Do not exceed 2 fl. oz. during dormant season or exceed 6.1 fl. oz. during growing season. Minimum 14 days between applications. |
| Annual broadleaves and suppression of grasses | Chateau SW (flumioxazin 51% a.i.) at 6-12 oz. Chateau EZ (flumioxazin 41.4% a.i.) at 6-12 oz. | Include an adjuvant (0.25% v/v non-ionic surfactant or 1 qt./A crop oil concentrate) for post-emergence use. See Pre-emergence section for other notes and restrictions. |
| Annual broadleaves and suppression of grasses | Chateau Complete (flumioxazin 30% a.i. + rimsulfuron 8.25%) at 6-12 oz. | Not registered in all states. Do not apply more than 12 oz. per acre per year. Do not apply to crops established less than one year. Do not apply to plants less than 2 years old unless they are protected by nonporous wrap, grow tubes or waxed containers. If weeds are emerged at the time of application, apply with an adjuvant (0.25% v/v non-ionic surfactant or 1 qt./A crop oil concentrate). For complete control of emerged weeds, addition of a labelled burndown product required. See Pre-emergence section for details. |
| Most annual and perennial grasses | Fusilade DX (fluazifop-p-butyl 2 lb. a.i./gal.) at 16-24 fl. oz. in 25 gal. water | Apply as a directed spray to actively growing grasses before tillering. Always add COC at 1% v/v or NIS at 0.25% v/v. Avoid contact with foliage. Rainfast in 1 hour. Do not exceed 48 fl. oz. in a maximum of two 24 oz. applications per year. Minimum 14 days between applications. |
| Most annual grasses and broadleaves and top kill of perennial weeds | Gramoxone SL 3.0 (paraquat 3 lb. a.i./gal.) at 1.3-2.7 pt. in minimum of 50 gal. water | Apply as directed spray to actively growing weeds before emergence of new canes or shoots. Repeat applications are necessary to give sustained control. Apply as a coarse spray to avoid drift injury. Avoid contact with desired new shoots, fruit, or foliage. Always add NIS at 0.25% v/v or COC at 1% v/v. Do not exceed 2 applications per year. Harvest fruit at normal maturity. Restricted use pesticide. Only certified applicators can mix, load and apply. Not to be used by uncertified persons working under the supervision of a certified applicator. Applicators must complete an EPA-approved paraquat training every 3 years https://www.epa.gov/pesticide-work-er-safety/paraquat-dichloride-training-certified-applicators Containers under 120 gallons will have "closed-system" packaging to be used with a closed-transfer system. |

(Continued)

Herbicide Recommendations For Highbush Blueberry (*continued*)

| Weed Problem | Material And Rate Per Acre | Notes And Comments |
|--|---|---|
| Most annual grasses and broadleaf weeds and top kill of biennial and perennial weeds | Homeplate (caprylic acid 45.14% + Capric acid 34.74%) at 3-9% solution if used alone or 1% solution when tank mixed | OMRI listed. Use includes vegetation burndown, directed and shielded sprays, and sucker control. May be used any time during the year and works best during warm and dry conditions. Reapply if rain falls within 3 hours of application. Avoid contact with desirable foliage and green bark. Sucker control: Apply before suckers become woody. |
| Annual and perennial grasses | Poast 1.5EC (sethoxydim 1.5 lb. a.i./gal.) at 1.5-2.5 pt. in minimum of 5 gal. water | Apply as a directed spray to actively growing grasses before tillering. Always add COC at 1% v/v. Do not exceed 2.5 pt. per application or exceed 5 pt. per season. |
| Annual grasses and broadleaves | Reglone (diquat 2 lb .a.i./gal.) at 1.5-2 pt. in minimum of 15 gal. water | Non-bearing Only: May only be used on crops that will not be harvested within 1 year of application. Apply as a directed spray using a shield for contact burn of weeds. Apply when wind speed is 3-10 miles per hour. Complete coverage is essential for good control. Always add NIS at 0.06-0.5% v/v. Can be used during site preparation, and up to 1 year of harvest. Do not allow contact with green stems, foliage, or fruits. |
| Annual and perennial grasses and broadleaves | Rely 280 (glufosinate 24.5% a.i. (2.34 lb./gal.)) at 48-82 fl. oz. in minimum of 15 gal. water | Apply as a directed spray to actively growing weeds. Do not apply on desirable foliage or drift on foliage, green, or uncalled bark. Coverage of all foliage is necessary for optimum control. Do not exceed 164 fl. oz. per acre per year. Do not make more than 2 applications at a maximum rate of 82 fl. oz. per acre per year. Add AMS to the spray tank if spray water is hard. |
| Annuals and some perennial grasses and broadleaves | Roundup WeatherMax 5.5EC (glyphosate 5.5 lb. a.i./gal.) at 0.5-5.3 qt. in 10-40 gal. water | Apply as a directed spray or wiper application to actively growing weeds in established plantings. Always add ammonium sulfate at 8.5-17 lb./100 gal. in hard water or drought conditions. Do not allow spray to contact any part other than mature bark. For applications within rows of berries, use only selective equipment (directed spray, hooded sprayer, shielded sprayer, or wiper application) to minimize the potential for overspray or drift onto the crop. For berry crops, hooded or shielded sprayers must be fully enclosed (including top, sides, front, and back). Only wiper applications or shielded sprayers capable of preventing all contact with the crop may be used. Rate depends on weed species and stage of growth. Can be mixed with labeled pre-emergence herbicides. |
| Annual and perennial grasses and broadleaves | Scythe 4.2E (pelargonic acid 4.2 lb. a.i./gal.) at 3-10% spray solution | For contact nonselective control or burndown of a broad spectrum of actively growing weeds. Use low rate for annual weed control and high rate for maximum vegetative burndown. Use as a directed spray or shielded spray. Can be mixed with Roundup. |
| Most annual and perennial grasses | Select Max with Inside Technology (clethodim 0.97 lb. a.i./gal.) at 9-16 fl. oz. | Apply as a directed spray to actively growing grasses before tillering. Minimum 14 days between applications. Always add NIS at 0.25% v/v. May be applied as a spot treatment at 0.32-0.64 fl. oz./gal. Rainfast in 1 hour. Do not exceed 64 fl. oz. per year. Do not use COC. |

(Continued)

Herbicide Recommendations For Brambles

| Material And Rate | | |
|---|--|--|
| Pre-emergence | | |
| Annual and perennial grasses and broadleaves | Casoron 4G (dichlorobenzil 4% a.i./lb.) at 100-150 lb. | For perennial weed control, apply to untilled ground over old weed growth from November 15 to Feb 15; alternately apply late fall or very early spring before May 15 and incorporate immediately. For annual weed control, surface apply no more than 100 lb./A. Shallow incorporation or sprinkler irrigation is recommended when application is made during periods of high temperatures. Do not apply until 4 weeks after transplanting. Use higher rate for perennial weed control. Annual maximum rate 150 lbs./A. |
| Annual broadleaves and suppression of grasses | Chateau EZ (flumioxazin 41.4% a.i.) at 6-12 oz. Chateau SW is not labelled for caneberrries in the Midwest | Apply as a uniform band directed at the base of the canes. Do not apply over the top of the crop or allow spray to come in contact with the crop as a result of application or drift. Preferred application timing is in the fall. Do not exceed 6 oz. per acre per application. Do not make a sequential application within 30 days of the first application Do not apply to farm alleys or roads where traffic may result in treated dust settling onto crops or other desirable vegetation. Do not mow treated areas between bud break and final harvest. Dust created by mowing may drift onto desirable vegetation resulting in injury. Do not apply within 300 yards of non-dormant pome or stone fruit. |
| Annual grasses and broadleaves | Devrinol 2-XT (napropamide 2 lb. a.i./gal.) at 2 gal./acre | Apply to a weed-free soil surface or tank mix with a suitable post emergent herbicide. May be applied to newly planted and newly established crops. Do not apply more than 2 gal. per acre per crop cycle. |
| Most broadleaves | Gallery 75DF (isoxaben 75% a.i.) at 0.66-1.33 lb. in minimum of 10 gal. water Gallery SC (isoxaben 4.16 lb. a.i./gal.) at 16-31 oz. in minimum of 10 gal. water | Non-bearing Only: May only be used on crops that will not be harvested within 1 year of application. Apply in late summer to early fall, or in early spring prior to weed germination, or anytime immediately after cultivation. Do not apply to new transplants until soil has settled with no cracks present. Rainfall or irrigation of 1/2 inch is needed within 21 days of application. Minimum 60 days between applications. Do not exceed 4 lb per acre per 12-month period. Non-Bearing Only: Make a single application prior to germination of target weeds or immediately after cultivation. May only be used on crops that will not be harvested within 1 year of application. Do not exceed 1.0 lb. a.i./A/yr. |
| Annual grasses and broadleaves | Karmex DF (diuron 80% a.i.) at 3 lb. in 25-40 gal. water | Selected states only. IN and OH only: Apply 3 lb in late spring for raspberries. If used post-emergence, avoid contact with foliage. Best results if temperature is at least 70°F with high humidity. Age Restriction: Apply in fields established at least 1 year. Do not exceed 1 application per year. Do not spray exposed roots to avoid injury. |
| Annual grasses and broadleaves | Princep 4L (simazine 4 lb. a.i./ gal.) at 2-4 qt. in minimum of 40 gal. water | Apply in spring before weeds emerge and before canes leaf out or make a split application of 2 qt. in spring plus 2 qt. in fall. On plants established less than 6 months, apply at half the rate. Do not apply when fruit is present, or illegal residues may result. |
| Annual grasses and broadleaves | Sinbar WDG (terbacil 80% a.i.) at 1-2 lb. in minimum of 20 gal. water | Make a single band or broadcast application as a directed spray to soil beneath the canes in the fall or early spring before fruit set and shortly before or after weed emergence. Avoid contact of foliage or fruit with spray or mist. Do not use on soils where roots are exposed. Age Restriction: Use only on plantings established at least 1 year. |
| Annual grasses and certain broadleaves | Snapshott 2.5TG (isoxaben+trifluralin 2.5% a.i.) at 100-200 lb. | Non-bearing Only: May only be used on crops that will not be harvested within 1 year of application. For best results, 1/2 inch of rain or irrigation is needed within 3 days of application. Not effective on germinated weeds Do not apply to new transplants until soil has settled. Minimum 60 days between applications. Do not exceed 600 lb. per 12-month period. |
| Annual grasses and broadleaves and suppression of yellow nutsedge | Solicam DF (norflurazon 78.6% a.i.) at 2.5-5 lb. in minimum of 20 gal. water | Apply as a directed spray to settled and firm soil from fall to early spring before weeds emerge. Rainfall or irrigation of 1/2 inch within 4 weeks to activate. Do not contact fruit or foliage. Do not apply after bud break on sandy loam soils. Check label for maximum amount allowed per year depending on soil type. Age Restriction: Minimum 12 months after planting before first application. |

(Continued)

Herbicide Recommendations For Brambles (*continued*)

| Weed Problem | Material And Rate Per Acre | Notes And Comments |
|---|---|--|
| Annual grasses and certain broadleaves | Surflan As Specialty Herbicide (oryzalin 4 lb. a.i./gal.) at 2-6 qt. in 20-40 gal. water | No longer in production - supply extremely limited to not available. Make a single band or broadcast application to the ground beneath vines before weeds emerge. Apply alone to weed-free soil or post-emergence mixed with Roundup or Gramoxone. Rainfall or irrigation of 1/2 inch is required for activation. Minimum 2.5 months between applications. Do not exceed 12 qt. per year. |
| Annual and perennial broadleaves | Trellis (isoxaben 75% a.i.) at 0.67-1.33 lb. in minimum of 10 gal. water Trellis SC (isoxaben 4.16 lb. a.i./gal.) at 16-31 oz. | Non-Bearing Only: Apply before target weeds germinate or immediately after cultivation. Do not apply Trellis SC more than twice per crop year (harvest to harvest) up to a maximum total of 1.0 lb. a.i. (31 fl. oz.) per crop year. Do not apply to newly transplanted canes until soil has been settled and no cracks are present of plant injury may occur. |
| Annual and perennial grasses and broadleaves | Zeus Prime XC (carfentrazone- ethyl 3.5% and sulfentrazone 31.8% a.i.) at 7.7-15.2 fl. oz./acre in minimum of 10 gal. water | Apply as a broadcast or banded soil application directed to the base of the trunks of bushes. If weeds are present, tank mix with a post-emergence herbicide to eliminate emerged weeds. Apply a single broadcast application at 15.2 fl. oz. (0.41 lb. a.i.) per acre. May be applied as a banded treatment twice per year. Do not exceed 15.2 fl. oz. (0.41 lb. a.i.) /acre/year. Minimum 60 days between applications. Do not apply after bud break except with hooded or shielded sprayer. Age Restriction: Apply to crops that have been growing for at least 2 years and are in good condition. |
| Annual and perennial grasses and broadleaves | Zeus XC (sulfentrazone 39.6% a.i.) at 8-12 fl. oz./acre in a minimum of 10 gal. water | Apply as a broadcast or banded soil application directed to the base of the trunk of bushes. If weeds are present, tank mix with a post-emergence herbicide to eliminate emerged weeds. Make a single broadcast application at 8-12 fl. oz. (0.25-0.375 lb. a.i.) per acre. May be applied as a banded treatment twice per year. Minimum of 60 days between applications. Do not exceed 12 fl. oz. (0.375 lb. a.i.)/acre/year. Do not apply after bud break except with hooded or shielded sprayer. Age Restriction: Apply to crops that have been growing for at least 3 years and are in good condition. |
| Post-emergence | | |
| Annual broadleaves | Aim EC (carfentrazone 2 lb. a.i./gal.) at 1-2 fl. oz. in 20 gal. water | Apply with hooded shields between rows during growing season. Always add NIS at 0.25% v/v or COC at 1% v/v. Do not exceed 25.6 fl. oz. per year. Minimum 14 days between applications. Primocane Control: Apply when primocanes are 6 inches at 6.4 fl. oz. in minimum of 20 gal. water at intervals of 14-21 days. Direct sprays to bottom 18 inches of canes. |
| Annual broadleaves and suppression of grasses | Chateau EZ (flumioxazin 41.4% a.i.) at 6-12 oz./A Chateau SW is not labelled for caneberries in the Midwest | Include an adjuvant (0.25% v/v non-ionic surfactant or 1 qt./A crop oil concentrate) for post-emergence use. See Pre-emergence section for other notes and restrictions. |
| Most annual and perennial grasses | Fusilade DX 2EC (fluazifop-p 2 lb. a.i./gal) at 16-24 fl. oz. in 25 gal. water | Apply as a directed spray to actively growing grasses before tillering. Always add COC at 1% v/v or NIS at 0.25% v/v. Avoid contact with foliage. Rainfast in 1 hour. Do not exceed 48 fl. oz. in a maximum of two 24 fl. oz. applications per year. Minimum 14 days between applications. |

(Continued)

Herbicide Recommendations For Brambles (*continued*)

| Weed Problem | Material And Rate Per Acre | Notes And Comments |
|--|---|---|
| Most annual grasses and broadleaves and top kill of perennial weeds | Gramoxone (paraquat 3 lb. a.i./gal.) at 1.3-2.7 pt. in minimum of 50 gal. water | Apply as directed spray to actively growing weeds before emergence of new canes or shoots. Repeat applications are necessary to give sustained control. Apply as a coarse spray to avoid drift injury. Avoid contact with desired new shoots, fruit, or foliage. Always add NIS at 0.25% v/v or COC at 1% v/v. Do not exceed 2 applications per year. Harvest crop at normal harvest maturity. Restricted use pesticide. Only certified applicators can mix, load and apply. Not to be used by uncertified persons working under the supervision of a certified applicator. Applicators must complete an EPA-approved paraquat training every 3 years https://www.epa.gov/pesticide-work-er-safety/paraquat-dichloride-training-certified-applicators Containers under 120 gallons will have "closed-system" packaging to be used with a closed-transfer system. |
| Most annual grasses and broadleaf weeds and top kill of biennial and perennial weeds | Homeplate (caprylic acid 45.14% + Capric acid 34.74%) at 3-9% solution if used alone or 1% solution when tank mixed | OMRI listed. Use includes vegetation burndown, directed and shielded sprays, and sucker control. May be used any time during the year and works best during warm and dry conditions. Reapply if rain falls within 3 hours of application. Avoid contact with desirable foliage and green bark. Primocane suppression: Apply before primocanes become woody. |
| Annual grasses and broadleaves | Karmex DF (diuron 80% a.i.) at 3 lb. in 25-40 gal. water | Selected states only. See Pre-emergence section for details. |
| Annual and perennial grasses | Poast 1.5EC (sethoxydim 1.5 lb. a.i./gal.) at 1.5-2.5 pt. in minimum of 5 gal. water | Apply as a directed spray to actively growing grasses before tillering. Always add COC at 1% v/v. Do not exceed 5 pt. per season. May be used as a spot treatment at 1-1.5% solution. |
| Annual grasses and broadleaves | Reglone (diquat 2 lb. a.i./gal.) at 1.5-2 pt. in minimum of 15 gal. water | Non-bearing Only: May only be used on crops that will not be harvested within 1 year of application. Apply as a directed spray using a shield for contact burn of weeds. Apply when wind speed is 3-10 miles per hour. Complete coverage is essential for good control. Always use NIS at 0.06-0.5% v/v. Can be used during site preparation, and up to 1 year of harvest. Do not allow contact with green stems, foliage or fruits. |
| Annuals and some perennial grasses and broadleaves | Roundup WeatherMax 5.5EC (glyphosate 5.5 lb. a.i./gal.) at 0.5-5.3 qt. in 10-40 gal. water | Apply as a directed spray or wiper application to actively growing weeds in established plantings. Always add ammonium sulfate at 8.5-17 lb./100 gal. in hard water or drought conditions. Do not allow spray to contact any part other than mature bark. For applications within rows of berries, use only selective equipment (directed spray, hooded sprayer, shielded sprayer, or wiper application) to minimize the potential for overspray or drift onto the crop. For berry crops, hooded or shielded sprayers must be fully enclosed (including top, sides, front, and back). Only wiper applications or shielded sprayers capable of preventing all contact with the crop may be used. Use with extreme care around raspberries as serious damage may occur if any part of the plant comes in contact with the product. Rate depends on weed species and stage of growth. Can be mixed with labeled pre-emergence herbicides. |
| Annual and perennial grasses and broadleaves | Scythe 4.2E (pelargonic acid 4.2 lb. a.i./gal.) at 3-10% spray solution | For contact nonselective control or burndown of a broad spectrum of actively growing weeds. Use low rate for annual weed control and high rate for maximum vegetative burndown. Use as a directed spray or shielded spray. Can be mixed with Roundup. |
| Most annual and perennial grasses | Select Max with Inside Technology (clethodim 0.97 lb. a.i./gal.) at 9-16 fl. oz. | Apply post-emergence as a directed spray to young actively growing grasses. Do not exceed 16 fl. oz./A in a single application or 64 fl. oz./A per season. A minimum 14-day interval is required for repeat applications. Always add NIS at 0.25% v/v. Do not use COC. Rainfast in 1 hour. |

(Continued)

Herbicide Recommendations For Strawberry

| Weed Problem | Material And Rate Per Acre | Notes And Comments |
|--|--|--|
| Pre-emergence | | |
| Annual broadleaves and suppression of grasses | Chateau SW (flumioxazin 51% a.i.) at 3 oz. Chateau EZ (flumioxazin 41.4% a.i.) at 3 oz. | Annual Strawberries Grown on Plastic Mulch: Apply a minimum of 30 days before transplanting and before laying plastic. Can be mixed with Gramoxone or Roundup. Pre-emergence on Dormant Plants: Can be applied over the top of established or newly planted dormant strawberries. Add COC at 1% v/v or NIS at 0.25% v/v to help control emerged broadleaf weeds. Do not apply to frozen ground. Shielded or Hooded Application in Row Middles: Do not apply after fruit set and not over strawberry plants. Apply prior to weed emergence. |
| Annual grasses and certain broadleaves | Devrinol 2-XT (napropamide 2 lb. a.i./gal.) at 2 gal. | Established Plantings (spring): Apply after removing straw mulch. Water into soil to a depth of 2-4 inches (by rainfall or irrigation) within 24-72 hours of application. Established Plantings (fall): Apply before putting winter protective mulch over plants. Water into soil to a depth of 2-4 inches (by rainfall or irrigation) within 24-72 hours of application. Do not apply to frozen ground. Do not exceed 2 gal. per acre per crop cycle. Strawberries Not Grown with Plastic: Apply to a weed-free soil surface. May be applied to newly transplanted crops. Delay application until the desired number of daughter plants has become established. Do not exceed 2 gal. per acre per crop cycle. Do not apply from bloom through harvest. Annual Strawberries Grown on Plastic Mulch: Apply to a weed-free soil before laying plastic mulch. Incorporate to a depth of 2 inches within 24-72 hours of application and before laying plastic. May also be applied to soil between beds. Do not exceed 2 gal. per acre per crop cycle. Do not apply from bloom through harvest. |
| Annual broadleaves, especially winter annuals | Goal 2XL (oxyfluorfen 2 lb. a.i./gal.) at 1-2 pt. in minimum of 40 gal. water | Fallow Bed Preparation Only: Apply with Roundup for control of winter annual broadleaves a minimum of 30 days before transplanting. Fallow bed should be worked thoroughly to a depth of 2.5 inches prior to planting. |
| Broadleaves and suppression of certain grasses | Optogen (bicyclopyrone 0.045 lb. a.i.) at 3.5 fl. oz. | Apply after strawberry emergence or transplanting as either a row middle or post-directed application. Avoid contacting the crop with direct or indirect spray or crop injury will occur. Using a hooded or shielded sprayer will minimize potential crop injury when applying as a post-directed application. Add a nonionic surfactant (NIS) at 1 qt./100 gal. of water (0.25% v/v) or a crop oil concentrate (COC) at the rate of 1 gal./100 gallons of water (1.0% v/v). In addition to NIS or COC, dry spray grade ammonium sulfate (AMS) may be added at a rate of 8.5 to 17 lb./gal. of water. Liquid AMS may be substituted at an equivalent rate. The use of AMS will improve the consistency level of weed control. For optimal control, make application to small (<2") weeds. |
| Annual grasses and certain broadleaves | Prowl H2O 3.8E (pendimethalin 3.8 lb. a.i./ gal.) at 1-2 pt. in minimum of 20 gal. water. | Rate depends on soil type. See label for details. Apply as a broadcast spray before transplanting or after transplanting but before growth starts. May not be used on beds that will be covered in plastic. A second application may be used in a band between rows up to 35 days before harvest. Do not allow the spray to contact strawberry foliage. May be applied to strawberries in fall or winter dormancy prior to the onset of new growth. May be applied to perennial strawberries during renovation after foliage has been mowed, but prior to the onset of new growth. Adequate rainfall or irrigation after application prior to weed emergence provides the most benefit. Do not exceed 3 pt. per application or exceed 6 pt. per season. |
| Annual grasses and broadleaves | Sinbar WDG (terbacil 80% a.i.) at 2-8 oz. in minimum of 20 gal. water | Planting Year: Apply 2-3 oz. immediately after transplanting but before runners start to root. Application of 2-6 oz. can also be made to dormant plants in late summer or early fall for control of winter annual weeds. If transplants have started to develop new foliage in the spring, or are not dormant in late summer or early fall at time of application, 1/2 to 1 inch of rain or irrigation is necessary to wash Sinbar off. Do not use on soils with less than 0.5% organic material, as plant injury can occur. Harvest Years: Apply 4-8 oz. after post-harvest renovation and before new growth begins in midsummer. An additional 4-8 oz. prior to mulching in late fall is recommended to extend weed control through harvest of the following year. Do not exceed 8 oz. per season. Note: Strawberry varieties differ in sensitivity to Sinbar, and significant plant injury is possible. Conduct a field test before adoption as a normal practice, particularly for new varieties. |

(Continued)

Herbicide Recommendations For Strawberry (*continued*)

| Weed Problem | Material And Rate Per Acre | Notes And Comments |
|---|--|--|
| Annual broadleaf weeds, grasses and nutsedge | Spartan 4F (sulfentrazone 39.6%) at 4-8 oz. in 20-40 gal. water | Rate depends on soil texture. Some cultivars may be sensitive. Apply prior to planting, post-transplant before new leaves emerge from dormant crowns. Do not exceed 8 fl. oz. per acre per application or exceed 12 oz. (0.375 lb. a.i.)/acre/year. Some states may have supplemental or Special Local Need labels. |
| Post-emergence | | |
| Annual broadleaves | Aim EC (carfentrazone 2 lb. a.i./gal.) at 0.5-2 fl. oz. in minimum of 10 gal. water | Apply with hooded shields between rows during growing season to actively growing weeds. Best results when weeds are <4 inches and rosettes <3 inches across. Always add NIS at 0.25% v/v or COC at 1% v/v. Do not exceed 6.1 fl. oz./year. Minimum 14 days between applications. |
| Annual and some perennial broadleaves | 2,4-D amine (4 lb. a.i./ gal.) at 2-3 pt. in 25-50 gal. water | Several 2,4-D amine products are available, but only a few are labeled for strawberry. For Established Plantings Only: Apply in early spring when strawberries are dormant or immediately after last picking. Do not apply unless possible injury to the crop is acceptable. Do not tank mix with Poast. Check label for specific use directions. |
| Most annual and perennial grasses | Fusilade DX (fluazifop-p-butyl 2 lb. a.i./gal.) at 16 fl. oz. in 25 gal. water | Apply as a directed spray to actively growing grasses before tillering. Always add COC at 1% v/v or NIS at 0.25% v/v. Rainfast in 1 hour. Do not exceed 16 fl. oz. per year. Do not exceed 1 application per year. |
| Annual broadleaves | Goal 2XL (oxyfluorfen 2 lb. a.i./gal.) at 1-2 pt. in minimum of 40 gal. water | See Pre-emergence section for details. |
| Most annual grasses and broadleaves and top kill of perennial weeds | Gramoxone SL 3.0 (paraquat 3 lb. a.i./gal.) at 1.3 pt. in minimum of 20 gal. water | Apply as a postemergence directed spray in a minimum of 20 gal. per acre. Apply by directed spray between rows, using shields to prevent contact with crop. Add NIS at 0.25% v/v or COC at 1% v/v. Do not exceed 3 applications per year. Restricted use pesticide. Only certified applicators can mix, load and apply. Not to be used by uncertified persons working under the supervision of a certified applicator. Applicators must complete an EPA-approved paraquat training every 3 years https://www.epa.gov/pesticide-worker-safety/paraquat-dichloride-training-certified-applicators Containers under 120 gallons will have "closed-system" packaging to be used with a closed-transfer system. |
| Broadleaves and suppression of certain grasses | Optogen (bicyclopyrone 0.045 lb.) at 3.5 fl. oz. | See Pre-emergence section for details. |
| Most annual and perennial grasses (post-emergence only) | Poast 1.5EC (sethoxydim 1.5 lb. a.i./gal.) at 1-2.5 pt. in 25 gal. water | Apply to actively growing grasses before tillering. Always add COC at 1% v/v. May be used as a spot treatment at 1-1.5% spray solution. Do not exceed 2.5 pt per application or exceed 2.5 pt. per season. Caution: Application of Poast up to six weeks after Sinbar application can occasionally cause strawberry leaf injury. |
| Annuals and some perennial grasses and broadleaves | Roundup WeatherMax 5.5EC (glyphosate 5.5 lb. a.i./gal.) at 0.5-5.3 qt. in 10-40 gal. water | Apply as pre-plant broadcast application or in fall prior to planting for control of roots and rhizomes of perennial weeds or as a hooded, shielded directed spray or wiper application (33-100% solution) to actively growing weeds between rows in established plantings. Always add ammonium sulfate 8.5-17 lb./100 gal in hard water or drought conditions. Do not allow spray to contact any desired plants. Can be mixed with labeled pre-emergence herbicides. Rate depends on weed species and stage of growth. |
| Annual and perennial grasses and broadleaves | Scythe 4.2E (pelargonic acid 4.2 lb. a.i./gal) at 3-10% spray solution | For contact nonselective control or burndown of a broad spectrum of actively growing weeds. Use low rate for annual weed control and high rate for maximum vegetative burndown. Use as a directed spray or shielded spray. Can be mixed with Roundup. |
| Most annual and perennial grasses | Select Max with Inside Technology (clethodim 0.97 lb. a.i./gal.) at 9-16 fl. oz. | Apply as a directed spray to actively growing grasses before tillering. Do not add COC. Rainfast in 1 hour. Do not exceed 64 fl. oz./year or exceed 16 fl. oz./application. Minimum 14 days between applications. May be applied as a spot treatment at 0.32-0.64 fl. oz./gal. |
| | | <i>(Continued)</i> |

Herbicide Recommendations For Strawberry (*continued*)

| Weed Problem | Material And Rate Per Acre | Notes And Comments |
|----------------------------------|--|--|
| Annual and perennial broadleaves | Spur (clopyralid 3 lb. a.i./gal.) at 2/3 pt. in minimum of 10 gal. water | Not registered in all states, but has 24(c) special local needs registration in several states. For perennial strawberries only. Make 1 application after harvest. Make only 1 application per crop year. Do not tank mix with other herbicides. |
| Annual grasses and broadleaves | Ultra Blazer 2E (acifluorfen 2 lb. a.i./gal.) at 1.5 pt. in minimum of 20 gal. water | <p>May be applied up to the maximum application rate of 1.5 pt. per acre per application using ground equipment. Make broadcast applications in 20 -40 gal. water per acre. Reduce rates proportionally for band or strip treatment. Do not apply more than 3 pt. per acre per season. Apply with NIS, COC, AMS or UAN solution additive.</p> <p>Annual Strawberries Grown on Plastic Mulch: Make 1 banded application before laying plastic and after final land preparation, and prior to transplanting the crop. For application between rows of plastic mulch, apply as a direct-shielded application between mulched beds. Do not allow contact with strawberry plants.</p> <p>Perennial Strawberry (matted row): Make 2 applications: the first can be made after the last harvest or following bed renovation. The second can be made when plants are dormant during late fall to early spring.</p> |

Relative Effectiveness Of Herbicides For Fruit Crops¹

| Herbicide | Grasses | | | | | Annual Broadleaves | | | | | | | | | | | | | | | | | Perennial Weeds | | | | |
|-----------------------|----------------|----------------|----------------|----------------|----------------|--------------------|----------------|-------------------|----------------|----------------|----------------|----------------------|----------------|----------------|-----------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|-----------------|----------------|------------------|-----------------|--------------------|
| | Barnyardgrass | Crabgrass | Foxtails | Goosegrass | Panicum, Fall | Chickweed | Cocklebur | Groundsel, Common | Henbit | lambquarters | maretail | Morningglory, Annual | Mustards | Nightshades | Palmer Amaranth | Pigweed | Purslane | Ragweed | Shepherdspurse | Smartweeds | Velvetleaf | Waterhemp | Dandelion | Johnsongrass | Nutsedge, Yellow | Thistle, Canada | Woodsorrel, Yellow |
| Pre-emergence | | | | | | | | | | | | | | | | | | | | | | | | | | | |
| Alion | G | G | G | G | G | G | N | G | F | F | G | F | G | N | N | G | G | F | G | G | G | N | G | N | N | N | F |
| Bellum | N | N | N | N | N | G | G | N | N | G | F | F | G | G | F | G | N | F | N | G | G | G | N | N | N | N | N |
| Broadloom | N | N | N | N | N | N | F | F ² | N | F | N | F | F | N | N | N | F | F ² | F | G | F | N | N | N | N | N | N |
| Callisto | N | N | N | N | N | G | G | N | N | G | F | F | N | G | F | G | N | G | N | G | G | G | N | N | F | N | N |
| Casoron | N | G | G | G | G | G | F | G | G | G | F | N | G | N | N | G | G | G | G | G | G | N | G | N | N | G | G |
| Chateau | N | N | N | N | N | F | F | N | N | G | G | F | N | G | F | G | G | F | G | F | F | F | N | N | N | N | N |
| Chateau Complete | G | G | G | G | G | F | F | N | N | G | G | G | N | G | G | G | G | G | G | F | F | F | N | N | N | N | N |
| Crew | G | G | G | G | N | G | N | F | G | G | G | G | G | N | N | G | G | N | G | N | N | N | N | N | N | N | |
| Dacthal | G | G | G | G | G | F | N | N | N | F | N | N | N | N | N | F | F | N | N | N | N | N | N | N | N | N | |
| Devrinol | G | G | G | G | G | G | F | N | N | F | N | N | N | N | N | G | G | N | N | N | N | N | N | N | N | N | |
| Gallery, Trellis | N | N | N | N | N | G | F | G | G | G | F | N | G | G | N | G | G | G | N | N | G | N | N | N | N | N | G |
| Goal | N | N | F | F | N | N | F | G | F | G | F | F | G | G | N | G | F | N | F | F | F | F | N | N | N | N | F |
| Karmex | G | G | F | G | F | G | F | G | G | G | F | F | G | G | N | G | G | G | G | N | P | N | N | N | N | N | N |
| Kerb | G | N | F | G | G | G | N | N | G | G | N | G | G | G | N | N | G | F | G | F | N | N | N | N | N | N | N |
| Matrix | G | G | G | N | G | N | F | G | G | F | G | N | G | F | N | F | G | F | F | F | F | N | G | N ² | F | F | N |
| Mission | N | N | G | N | N | G | N | G | G | G | F | N | F | N | N | G | G | G | G | N | N | N | G | N | G | N | N |
| Optogen | F | F | F | F | F | F | F | N | G | F | N | F | F | G | F | G | F | G | N | F | G | F | N | N | N | N | N |
| Pindar GT | F | N | N | N | N | N | G | N | N | G | G | N | N | G | N | G | F | N | N | F | G | N | N | N | N | N | N |
| Princep | G | G | G | G | G | G | N | G | G | G | N | G | G | G | N | G | G | G | G | N | F | N | N | N | F | N | N |
| Prowl | G | G | G | G | G | G | N | N | N | G | N | N | N | N | G | F | F | N | G | F | F | G | N | N ² | N | N | N |
| Sandea | N | N | N | N | N | N | F | G | N | G | F | N | G | N | N | G | F | G | G | G | G | N | N | N | G | N | N |
| Sinbar | G | G | G | N | G | G | N | F | G | G | N | N | G | G | N | G | G | G | G | G | N | N | G | F | F | N | N |
| Snapshot | G | G | F | G | G | G | F | G | G | G | F | F | G | F | N | G | N | N | G | F | G | N | G | F | N | N | G |
| Solicam | G | G | G | G | G | G | G | F | F | G | F | N | G | F | F | G | F | G | G | N | G | F | N | F | F | N | N |
| Spartan | N | F | N | N | G | G | N | N | N | F | N | G | F | G | G | G | G | N | F | F | N | G | N | N | F | N | N |
| Surflan | G | G | G | G | G | G | N | F | G | G | N | N | N | F | N | G | G | F | G | F | F | N | N | N ² | N | N | N |
| Treflan | G | G | G | G | G | N | N | N | G | F | N | N | F | N | F | G | G | N | N | N | N | N | N | F | N | N | F |
| Velpar | G | N | F | N | G | G | N | G | N | G | F | N | N | N | N | N | N | G | N | G | F | N | F ² | N | N | N | N |
| Zeus Prime XC | G | G | G | G | G | G | N | G | G | G | N | G | G | G | F | G | G | N | G | G | N | G | N | N | G | G | N |
| Zeus XC, Spartan | N | G | N | G | N | G | N | G | N | G | N | G | G | G | F | G | G | N | G | F | F | F | G | F | G | G | F |
| Post-emergence | | | | | | | | | | | | | | | | | | | | | | | | | | | |
| 2,4-D | N | N | N | N | N | F | F | G | N | F | G | G | G | F | F | N | G | G | F | F | F | G | N | N | F | N | |
| Aim | N | N | N | N | N | N | F | G | F | G | N | G | G | F | G | G | F | F | F | G | F | N | N | N | F | N | |
| Broadloom | N | N | N | N | N | N | F | F ² | N | F | N | F | F | N | N | N | F | F ² | F | G | F | N | N | N | N | N | N |
| Chateau | N | N | N | N | N | G | N | N | N | G | G | F | N | F | F | G | F | G | F | G | F | N | N | N | N | N | |
| Chateau Complete | G | G | G | G | G | F | F | N | N | G | G | G | N | G | G | G | G | G | G | F | F | F | N | N | N | N | N |
| Crew | G | G | G | G | N | G | N | F | G | G | G | G | G | N | N | G | G | N | G | N | N | N | N | N | N | N | |
| Embed | N | N | N | N | N | F | F | G | N | F | G | G | G | F | F | N | G | G | F | F | F | G | N | N | F | N | |
| Fusilade | G | G | G | G | G | N | N | N | N | N | N | N | N | N | N | N | N | N | N | N | N | N | F ² | N | N | N | |
| Gamma | F | F | N | N | N | F | N | N | G | N | F | N | N | N | N | N | N | N | F | N | G | F | F | G | N | N | N |
| Goal | N | F | F | F | N | N | F | G | G | G | F | F | G | G | N | G | F | N | F | F | F | F | N | N | N | N | F |
| Gramoxone | G | G | G | G | G | G | G | G | G | G | G | G | G | G | G | G | G | G | N | G | G | G | N | N | N | N | N |
| Homeplate | F ² | G | F ² | F ² | F ² | F ² | F ² | F ² | F ² | F ² | F ² | F ² | F ² | F ² | F ² | F ² | F ² | N ² | N | N | N | N ² | |

(Continued)

Relative Effectiveness Of Herbicides For Fruit Crops¹ (continued)

| Herbicide | Grasses | | | | | Annual Broadleaves | | | | | | | | | | | | | | | | Perennial Weeds | | | | | | | |
|-----------------------|----------------|----------------|----------------|----------------|----------------|--------------------|-----------|-------------------|----------------|----------------|----------------|----------------------|----------------|----------------|-----------------|----------------|----------------|----------------|----------------|----------------|----------------|-----------------|-----------|----------------|------------------|-----------------|--------------------|----------------|---|
| | Barnyardgrass | Crabgrass | Foxtails | Goosegrass | Panicum, Fall | Chickweed | Cocklebur | Groundsel, Common | Henbit | lambquarters | marestail | Morningglory, Annual | Mustards | Nightshades | Palmer Amaranth | Pigweed | Purslane | Ragweed | Shepherdspurse | Smartweeds | Velvetleaf | Waterhemp | Dandelion | Johnsongrass | Nutsedge, Yellow | Thistle, Canada | Woodsorrel, Yellow | | |
| Post-emergence | | | | | | | | | | | | | | | | | | | | | | | | | | | | | |
| Mission | N | G | G | N | N | | G | N | G | G | G | N | G | N | N | G | G | G | G | N | N | N | | F | N | G | G | N | |
| Optogen | F | F | F | F | F | | F | F | N | G | F | N | F | F | G | F | G | F | G | N | F | G | F | | N | N | N | N | N |
| Poast | G | G | G | G | G | | N | N | N | N | N | N | N | N | N | N | N | N | N | N | N | N | N | | N | F | N | N | N |
| Reglone | G | G | G | G | G | | G | G | G | G | G | G | G | G | G | G | G | G | G | G | G | G | G | | N | N | N | N | N |
| Rely | G | N | G | G | G | | G | G | N | N | G | G | G | G | F | G | G | G | G | G | G | F | F | | G | F ² | F | G | N |
| Roundup | G | G | G | G | G | | G | G | G | G | G | F | G | G | G | F | G | G | G | G | G | G | F | | G | F | F | G | G |
| Scythe | F ² | | G | F ² | F ² | F ² | F ² | F ² | F ² | F ² | F ² | F ² | F ² | F ² | F ² | F ² | F ² | F ² | | N ² | N | N | N | N ² | |
| Select | G | G | G | G | G | | N | N | N | N | N | N | N | N | N | N | N | N | N | N | N | N | N | | N | N | N | N | N |
| Starane Ultra | N | N | N | N | N | | G | G | N | N | N | N | F | F | F | N | N | N | F | N | N | G | N | | N | N | N | N | N |
| Stinger | N | N | N | N | N | | N | F | G | N | N | G | N | N | G | N | N | N | G | N | F | N | N | | G | N | N | G | N |
| Targa | G | G | G | G | G | | N | N | N | N | N | N | N | N | N | N | N | N | N | N | N | N | N | | N | F | N | N | N |
| Treevix | N | N | N | N | N | | N | G | F | N | G | G | G | G | N | G | G | G | G | G | G | G | G | | N | N | N | N | N |
| Trellis | N | N | N | N | N | | G | N | F | G | G | F | N | G | G | N | N | G | N | G | G | N | N | | N | N | N | N | N |
| Ultra Blazer | N | N | G | N | G | | N | F | N | N | G | N | G | G | G | G | G | G | N | G | G | G | G | | N | F ² | N | N | N |
| Velpar | G | N | F | N | G | | G | N | G | N | G | F | N | N | N | N | N | N | G | N | G | F | N | | F ² | N | N | N | N |
| Venue | N | N | N | N | N | | G | G | N | G | G | F | G | N | G | F | G | G | G | G | G | G | F | | G | N | N | N | N |
| Zalo | G | G | G | G | G | | G | G | N | N | G | F | G | G | F | G | G | G | G | G | G | G | G | | N | N | N | N | N |

G = good. F = fair. N = not listed, based on product labels.

²Provides partial control.

Tree Fruit Herbicide REI, PHI And Special Notes

| Trade Name | Common Name | WSSA | Risk of Resistance | REI | Apple | Pear | Peach | Nectarine | Plum | Cherry |
|------------------|-----------------------------|--------|--------------------|-----|--------------------------------|--------------------------------|--------------------------------|--------------------------------|--------------------------------|--------------------------------|
| Aim EC | carfentrazone-ethyl | 14 | medium | 12 | 3 | 3 | 3 | 3 | 3 | 3 |
| Alion | indaziflam | 21 | medium | 12 | 14 | 14 | 14 | 14 | 14 | 14 |
| Bellum | mesotrione | 27 | medium-high | 12 | 30 | 30 | 30 | 30 | 30 | 30 |
| Broadloom | bentazon | 6 | medium | 48 | NB/1 yr |
| Casoron 4G | dichlobenil | 20 | medium | 12 | N/A | N/A | -- | -- | -- | N/A |
| Chateau SW/EZ | flumioxazin | 14 | medium | 12 | Not after pink bud/60 | Not after bud break/60 |
| Chateau Complete | flumioxazin + rimsulfuron | 14 + 2 | Medium + high | 12 | Not after pink bud/60 | Not after bud break/60 |
| Crew | isoxaben + dithiopyr | 21 + 3 | Medium + low | 24 | NB/1 yr |
| Embed | 2,4-D choline | 4 | low | 48 | 14 | 14 | 40 | 40 | 40 | 40 |
| 2,4-D amine | 2,4-D amine | 4 | low | 48 | 14 | 14 | 40 | 40 | 40 | 40 |
| Fusilade DX | fluazifop | 1 | high | 12 | NB/1 yr | NB/1 yr | 14 | 14 | 14 | 14 |
| Gallery | isoxaben DF or SC | 21 | medium | 12 | NB/1 yr |
| GoalTender | oxyfluorfen 41% | 14 | medium | 24 | Fallow Pre-plant N-B Dormant |
| Goal 2XL | oxyfluorfen 22.3% | 14 | medium | 24 | Fallow Pre-plant N-B Dormant |
| Gramoxone SL 3.0 | paraquat | 22 | medium | 24 | N/A | N/A | 14 | 28 | 28 | 28 |
| Homeplate | caprylic acid + capric acid | 0 | very low | 24 | N/A | N/A | N/A | N/A | N/A | N/A |
| Karmex DF | diuron | 7 | medium | 12 | N/A | N/A | 20 | -- | -- | -- |
| Kerb SC | pronamide | 3 | low | 24 | N/A ¹ |
| Matrix FNV | rimsulfuron | 2 | medium | 4 | 7 | 7 | 14 | 14 | 14 | 14 |
| Pindar GT | penoxsulam | 2 | high | 24 | 60 | 60 | 60 | 60 | 60 | 60 |
| Poast 1.5 EC | sethoxydim | 1 | high | 12 | 14 | 14 | 25 | 25 | NB/1yr | 25 |
| Princep 4L | simazine | 5 | medium | 12 | 150 | 21 | 21 ⁷ | -- | 21 ⁷ | 21 ⁷ sweet |
| Prowl | pendimethalin | 3 | low | 12 | NB/1 yr |
| Prowl H20 | pendimethalin | 3 | low | 12 | 60 | 60 | 60 | 60 | 60 | 60 |
| Reglone | diquat | 22 | medium | 24 | NB/1 yr |
| Rely 280 | glufosinate | 10 | medium | 12 | 14 | 14 | 14 | 14 | 14 | 14 |
| Roundup | glyphosate | 9 | low | 12 | 14 | 14 | 17 | 17 | 17 | 17 |
| Sandea | halosulfuron | 2 | low | 12 | 14 | 14 | -- | -- | -- | -- |

(Continued)

Tree Fruit Herbicide REI, PHI And Special Notes (continued)

| Trade Name | Common Name | WSSA | Risk of Resistance | REI | Apple | Pear | Peach | Nectarine | Plum | Cherry |
|---------------|--|-----------|--------------------|-----|------------|---------|---------|-----------|---------|---------|
| Scythe | pelargonic acid | 26 | low | 12 | N/A | N/A | N/A | N/A | N/A | N/A |
| Select Max | clethodim | 1 | high | 12 | 14 | 14 | 14 | 14 | 14 | 14 |
| Showcase | trifluralin+ isoxaben+ oxyfluorfen | 3, 21, 14 | medium | 24 | -- | -- | NB/1 yr | NB/1 yr | NB/1 yr | NB/1 yr |
| Sinbar WDG | terbacil | 5 | medium | 12 | 60 | -- | 60 | -- | -- | -- |
| Snapshot | isoxaben+ trifluralin | 21+3 | medium | 12 | NB/1 yr | NB/1 yr | NB/1 yr | NB/1 yr | NB/1 yr | NB/1 yr |
| Solicam DF | norflurazon | 12 | medium | 12 | 60 | 60 | 60 | 60 | 60 | 60 |
| Starane Ultra | fluroxypyr | 4 | medium | 24 | 14 | 14 | -- | -- | -- | -- |
| Stinger | clopyralid | 4 | medium | 12 | 30 | -- | 30 | 30 | 30 | 30 |
| Surflan 4AS | oryzalin | 3 | low | 24 | N/A | N/A | N/A | N/A | N/A | N/A |
| Targa | quizalofop p-ethyl | 1 | high | 12 | 14 | 14 | 14 | 14 | 14 | 14 |
| Treevix | saflufenacil | 14 | low | 12 | 0 | 0 | -- | -- | -- | -- |
| Treflan | trifluralin | 3 | low | 12 | -- | -- | N/A | N/A | N/A | N/A |
| Trellis | isoxaben 75% | 21 | medium | 12 | NB/1 yr | NB/1 yr | NB/1 yr | NB/1 yr | NB/1 yr | NB/1 yr |
| Trellis SC | isoxaben 45.45% | 21 | medium | 12 | NB/1 yr 30 | NB/1 yr | NB/1 yr | NB/1 yr | NB/1 yr | NB/1 yr |
| Venue | pyraflufen ethyl | 14 | medium | 12 | 0 | 0 | 0 | 0 | 0 | 0 |
| Zalo | quizalofop p-ethyl + glu-fosinate-ammonium | 1 +10 | high + medium | 12 | 14 | 14 | 14 | 14 | 14 | 14 |
| Zeus Prime XC | carfentrazone-ethyl+ sulfentra-zone | 1 | high | 12 | 14 | -- | -- | -- | -- | -- |

-- = not labeled

DS/NCC = Directed Spray/No Crop Contact N/A - no PHI specified

1 = application must be in the fall, after the fruit is harvested, but prior to soil freeze-up

2 = application in the fall or early winter, but prior to soil freeze-up and snow cover

3 = apply before emergence of new canes or shoots

4 = PHI for Highbush Blueberry only, no PHI stated for Lowbush Blueberry

5 = See label

6 = do not apply when fruit is present or illegal residues may result

7 = apply late fall to early spring prior to weed emergence. Do not apply more than once per calendar year

8 = apply anytime between harvest and early spring. Do not apply more than once per calendar year

Small Fruit Herbicide REI, PHI And Special Notes

| Trade Name | Common Name | WSSA | Risk of Resistance | REI | Strawberry | Raspberry | Blackberry | Grape | Blueberry |
|------------------|---------------------------|--------|--------------------|-----|---|---------------------|---------------------|--------------------|---------------------|
| Aim EC | carfentrazone-ethyl | 14 | medium | 12 | 0 | 15 | 15 | 3 | 0 |
| Alion | indaziflam | 21 | medium | 12 | -- | 14 | 14 | 14 | 14 |
| Amine 4 | 2,4-D amine | 4 | low | 48 | N/A | -- | -- | -- | -- |
| Callisto | mesotrione | 27 | medium | 12 | -- | no bloom to harvest | no bloom to harvest | -- | no bloom to harvest |
| Casoron 4G | dichlobenil | 20 | medium | 12 | -- | N/A | N/A | N/A | N/A |
| Chateau SW/EZ | flumioxazin | 14 | medium | 12 | Row middle application - Do not apply after fruit set | 7 | 7 | 60 | 7 |
| Chateau Complete | flumioxazin + rimsulfuron | 14 + 2 | medium | 12 | -- | -- | -- | 60 | 21 |
| Devrinol DF-XT | napropamide | 15 | low | 24 | N/A ⁵ | N/A | N/A | 70 | N/A |
| Fusilade DX | fluazifop | 1 | high | 12 | 14 | 1 | 1 | 50 | NB/10 mo |
| Gallery DF or SC | isoxaben | 21 | medium | 12 | -- | NB/1 yr | NB/1 yr | NB/1 yr | NB/1 yr |
| Gama | tiafenacil | 14 | medium | 12 | | | | 7 | |
| GoalTender | oxyfluorfen 41% | 14 | medium | 24 | Fallow PP | -- | -- | Fallow PP Dormant | -- |
| Goal 2XL | oxyfluorfen 22.3% | 14 | medium | 24 | Fallow PP | -- | -- | Fallow PP Dormant | -- |
| Gramoxone SL 3.0 | paraquat | 22 | medium | 24 | 21 | N/A ³ | N/A ³ | N/A ⁵ | N/A ³ |
| Homeplate | Caprylic + capric acids | | low | 12 | N/A | N/A | N/A | N/A | N/A |
| Karmex DF | diuron | 7 | medium | 12 | -- | 0 | 0 | 0 | 0 |
| Kerb SC | pronamide | 3 | low | 24 | -- | -- | -- | N/A ¹ | N/A ² |
| Matrix FNV | rimsulfuron | 2 | medium | 4 | -- | | | 14 | -- |
| Mission | flazasulfuron | 2 | medium | 12 | -- | -- | -- | 75 | -- |
| Optogen | bicyclopyrone | 27 | medium | 24 | 30 | -- | -- | -- | -- |
| Poast 1.5 EC | sethoxydim | 1 | high | 12 | 7 | 45 | 45 | 50 | 1 HB/30 LB |
| Princep 4L | simazine | 5 | medium | 12 | -- | N/A ⁶ | N/A ⁶ | N/A ⁸ | N/A ⁶ |
| Prowl | pendimethalin | 3 | low | 12 | -- | -- | -- | NB/1 yr | -- |
| Prowl H2O | pendimethalin | 3 | low | 12 | 35 | -- | -- | 21 | -- |
| Reglone | diquat | 22 | medium | 24 | -- | NB/1 yr | NB/1 yr | NB/1 yr | NB/1 yr |
| Rely 280 | glufosinate | 10 | medium | 12 | -- | -- | -- | 14 | 14 |
| Roundup | glyphosate | 9 | low | 12 | 14 | 14 | 14 | 14 | 14 |
| Sandea | halosulfuron | 2 | low | 12 | -- | | | -- | 14 |
| Scythe | pelargonic acid | 26 | low | 12 | N/A | N/A | N/A | N/A | N/A |
| Select Max | clethodim | 1 | high | 12 | 4 | 7 | 7 | | 14 HB/45 LB |
| Sinbar WDG | terbacil | 5 | medium | 12 | 110 | 70 | 70 | -- | N/A |
| Sinbar WDG | terbacil | 5 | medium | 12 | 110 | 70 | 70 | -- | N/A |

(Continued)

Small Fruit Herbicide REI, PHI And Special Notes (continued)

| Trade Name | Common Name | WSSA | Risk of Resistance | REI | Strawberry | Raspberry | Blackberry | Grape | Blueberry |
|---------------|-------------------------------------|------|--------------------|-----|------------------|-----------|------------|------------|-------------------------|
| Snapshot | isoxaben+trifluralin | 21+3 | medium | 12 | -- | NB/1 yr | NB/1 yr | NB/1 yr | NB/1 yr |
| Solicam DF | norflurazon | 12 | medium | 12 | -- | Dormant | Dormant | 60 | 60 |
| Spartan | sulfentrazone | 14 | medium | 12 | preplant only | -- | -- | -- | -- |
| Spur, Stinger | clopyralid | 4 | medium | 12 | N/A ¹ | -- | -- | -- | -- |
| Surflan 4AS | oryzalin | 3 | low | 24 | -- | N/A | N/A | N/A | HB N/A |
| Treflan | trifluralin | 3 | low | 12 | -- | -- | -- | 60 | -- |
| Trellis | isoxaben 75% | 21 | medium | 12 | -- | NB/1 yr | NB/1 yr | NB/1 yr 60 | NB/1 yr |
| Trellis SC | isoxaben 45.45% | 21 | medium | 12 | -- | NB/1 yr | NB/1 yr | NB/1 yr 60 | NB/1 yr 60 ⁴ |
| Ultra Blazer | acifluorfen | 14 | medium | 48 | 60/120 | -- | -- | -- | -- |
| Velpar L CU | hexazinone | 5 | medium | 48 | -- | -- | -- | -- | HB 90/LB 450 |
| Venue | pyraflufen ethyl | 14 | medium | 12 | -- | -- | -- | 0 | -- |
| Zeus Prime XC | carfentrazone- ethyl+ sulfentrazone | 14 | medium | 12 | -- | 3 | 3 | 3 | 3 |
| Zeus XC | sulfentrazone | 14 | medium | 12 | -- | 3 | 3 | 3 | 3 |

-- = not labeled

DS/NCC = Directed Spray/No Crop Contact N/A - no PHI specified\

1 = application must be in the fall, after the fruit is harvested, but prior to soil freeze-up

2 = application in the fall or early winter, but prior to soil freeze-up and snow cover

3 = apply before emergence of new canes or shoots

4 = PHI for Highbush Blueberry only, no PHI stated for Lowbush Blueberry

5 = See label

6 = do not apply when fruit is present or illegal residues may result

7 = apply late fall to early spring prior to weed emergence. Do not apply more than once per calendar year

8 = apply anytime between harvest and early spring. Do not apply more than once per calendar year

Generic Pesticides

A generic agricultural chemical is manufactured and sold by a company other than the original manufacturer and patent holder, usually after the patent has expired. The generic pesticide contains the same active ingredient(s) (AI) and tend to be similar in performance to receive an EPA registration.

Generic products are not always identical, so be sure to carefully read the label, with special attention to rates and percent active ingredient.

Generic Fungicides

| Original Trade Name (Current Manufacturer) | Common Name | Other Trade Names (Manufacturers) |
|--|-------------------------------------|---|
| Abound (Syngenta) Quadris (Syngenta) | azoxystrobin | Aframe (Syngenta) Azaka (FMC) Azoxystar, Equation |
| Aliette 80WDG (Bayer) | fosetyl-Al | Legion 80WDG (Makhteshim) Linebacker WDG (NovaSource) |
| Apogee (BASF) | prohexadione-calcium (Pro-Ca) | Cryova PGR (Makhteshim) Kudos (Fine Americas) Pro Hex (Axil Solutions) Regalis PLUS (BASF) |
| Bravo Weather Stick (Syngenta) | chlorothalonil | Echo (Sipcam Agro) Equus DF (Sipcam Agro, AMVAC) Chlorothalonil 720 (Albaugh) |
| Captec 4L (Arysta LifeScience North America LLC) | captan | Has several formulations including 50W |
| Copper | copper hydroxide | Champ (Nufam) Kocide (Certis) KOP-Hydroxide (Drexel) Nu-Cop (Albaugh) |
| | copper octanoate | Camelot-O (SePRO) Cueva (Certis) |
| | copper oxychloride | COC (Albaugh) |
| | copper oxychloride+copper hydroxide | Badge (Gowan) |
| | copper (cuprous) oxide | Nordox (NOROX Industrier) |
| | copper sulfate (basic) | Basic Copper (Albaugh) Cuprofix (UPL) |
| | copper sulfate pentahydrate | KOP-5 (Drexel) Mastercop (ADAMA) Phyton 35 (Phyton Corp) |
| Dithane M45 (Dow AgriSciences) | mancozeb | Manzate Max (United Phosphorus, Inc) Penncozeb (several formulations) Roper (Loveland) Koverall (Cheminova) |
| Elite 45DF | tebuconazole | Orius 3.6F (Makhteshim) Orius 20AQ (Makhteshim) TebuStar 3.6 L (Albaugh) TebuStar 45WSP (Albaugh) Tebuzol 45DF (United Phosphorous, Inc.) |
| PH-D (United Phosphorous Inc.) | Polyoxin D zinc salt | Affirm (Nufarm) OSO (Certis Biologicals) |
| ProBlad Verde SymAgro | Banda Lupinus Albus Doce | Fracture (FMC) |
| Prophyt (Helena) | Phosphite (mono- and dibasic salts) | AgriFos (Monterey)/Agri-Fos (AgriChem) K-Phite (Plant Food Systems) Phostrol (Nufarm) Reliant (Quest Products) |
| Quilt Xcel (Syngenta) | azoxystrobin+ propiconazole | Aframe Plus (Syngenta) Cover XL (AgriStar) |

(Continued)

Generic Fungicides (continued)

| Original Trade Name (Current Manufacturer) | Common Name | Other Trade Names (Manufacturers) |
|---|--------------------|---|
| Rally 40WSP (Dow AgriSciences) | myclobutanil | Sonoma 40WSP (Albaugh) |
| Ridomil 2E (Syngenta) | metalaxyl | Metastar 2E (Arysta Life Science) |
| Ridomil Gold SL (Syngenta) | mefenoxam | Apron XL (Syngenta) ReCon Bold SL (Atticus) Thrive 4M (Albaugh) Ultra Flourish (Nufarm) |
| Rovral 4F (Bayer) | iprodione | Iprodione 4L AG (Arysta Life Science) Meteor 4L (United Phosphorus Inc.) Nevado 4F (Makhteshim) |
| Streptomycin 17 (Loveland Products Canada Inc.) | streptomycin | AG Streptomycin (ADAMA) FireWall (AgroSource) |
| Tilt (Syngenta) | propiconazole | Propimax 41.8L (Dow AgriSciences) Bumper 41.8L (Makhteshim) Orbit 41.8L (Syngenta) |
| Topsin-M 70WDG (United Phosphorous Inc.) | thiophanate methyl | Thiophanate Methyl 85WSB (Makhteshim) T-Methyl EAG 70WSB (Nufarm) T-Methyl 70WWSB (Arysta Life Science) |

Generic Insecticides

| Original Trade Name (Current Manufacturer) | Common Name | Other Trade Names (Manufacturers) |
|---|--------------|--|
| Acramite (UPL) | bifenazate | Actuate (Atticus) Bifenamite 2SC (Albaugh) Bizate (Loveland) Engulf (Nufarm) Floramite SC (OHP) Floramite SC/LS (UPL) Vigilant 4SC (UPL) |
| Admire Pro (Bayer) | imidacloprid | Acronyx 4F (Atticus) Advise Four (WinField) Alias 4F (ADAMA) Macho 2FL, 4 (Albaugh) Madari 4F (Avalaire) Malice 75WSP (Loveland) Midash 2SC, Forte 4F (Sharda) Montana 2F, 4F (Albaugh) Nuprid 4F Max (Nufarm) Prey 1.6F (Loveland) Provoke (Innvictis) Sherpa 1.6F (Loveland) Viloprid 4, FC 1.7 (Vive) Widow 2F (Loveland) Willowood 4SC (Generic) Wrangler 4F (Loveland) |

(Continued)

Generic Insecticides (continued)

| Original Trade Name (Current Manufacturer) | Common Name | Other Trade Names (Manufacturers) |
|---|-------------------------------|--|
| Agri-Mek SC (Syngenta) | abamectin | Abacus 0.15EC (Rotam) Abamex 0.15ED (Nufarm) Abba Ultra 0.30EC (Amvac) Averland 0.7FC (Vive) Borrada 0.15EC (ADAMA) Enterik 0.15LV, 0.7SC (Atticus) Reaper 0.15EC Advance 0.15EC Clearform 0.15EC (Loveland) Willowood Abamectin 0.15LV, 0.7SC (Generic) |
| Asana XL 0.66EC (Valent) | esfenvalerate | S-Fenvalostar 0.66EC (LG Life Sciences) |
| Assail 30SG, 30SC, 70WP (UPL) | acetamiprid | Afflict 30SG, 70WP (Aceto) Anarchy 30SG, 70WP (Loveland) ArVida 30SG (Atticus) Intruder Max 70WP (UPL) |
| Baythroid XL 1EC (Bayer) | Beta-cyfluthrin, cyfluthrin | Cryptoid XL 1EC (Atticus) Tombstone 2E (Loveland) Tombstone Helios 2E (Loveland) |
| Brigade 2EC, eVo (FMC) | bifenthrin | Banister 2EC (Avalaire) Batallion 2EC (Atticus) Bi-Dash 2EC (Sharda) Bifen 2AG Gold (WinField) Bifender 1.75FC (Vive) Bifenthrin 2EC (Aceto) Bifenture 2EC (UPL) Discipline 2EC (Amvac) Fanfare 2EC (ADAMA) GCS Bifenthrin 2EC, LFC 1.5EC (Generic) Lancer 2EC, FC 1.5EC (Albaugh) Reveal 2EC, Endurx 2EC (Innervictis) Sniper 2EC (Loveland) Tundra 2EC (Winfield) |
| Cygon 400EC (FMC) | dimethoate | Dimate 4EC (Winfield) Dimethoate 400, 400EC, 4EC (Loveland, FMC, Drexel) |
| Dimilin | diflubenzuron | Diflumas 2L (Helm Agro) Dimilin 2L, 25W (UPL) Durant 2L (Atticus) Micromite 2L (UPL) Unforgiven (Loveland) |
| Dipel (Valent) | <i>Bacillus thuringiensis</i> | Agree (Certis) Biobit (Valent) Bt Now (BioSafe) CryMax (Certis) Deliver (Certis) Javelin (Certis) Leptotec (Vestaron) Xentari (Valent) |
| Esteem 0.86EC, 35WP (Valent) Knack 0.83EC (Valent) | pyriproxyfen | Pitch (0.83EC), Pitch 35WP (ADAMA) |

(Continued)

Generic Insecticides (continued)

| Original Trade Name (Current Manufacturer) | Common Name | Other Trade Names (Manufacturers) |
|---|--------------------|---|
| Intrepid 2F (Corteva AgriSciences) | methoxyfenozide | GCS Methoxy 2F (Generic) Inspirato 2F (Atticus) Invertid 2F (Loveland) Invicar 2SC (Albaugh) Thwartex (Agsurf) Troubador 2F (Helena) Turnstyle 2F (UPL) Vexer (Innvictis) Zylo (UPL) |
| Mustang Maxx (FMC) | zeta-cypermethrin | Cortes Maxx (Atticus) |
| Pounce 25WP (FMC) | permethrin | Arctic 3.2EC (Winfield) Permethrin 3.2EC (Loveland) Perm-Up 3.2EC, 25DF (UPL) Permastar AG (LG Life Sciences) |
| Savey 50DF, Onager, Onager Optek, Hexy- gon, Hexygon IQ (Gowan) | hexythiazox | Hexamite (Albaugh) Hexcel EW (Atticus) |
| Sevin XLR Plus, 4L (Novasource) | carbaryl | Carbaryl 4L (Drexel, Loveland) |
| Warrior II 2.08CS (Syngenta) | lambda-cyhalothrin | Actylis Lambda-Cy (Aceto) Calvary II (Growmark) Crusader 1EC, 2ME (Albaugh) Firestone (Altitude) Grizzly Too, Z 1CS (Winfield) Kaiso 24WG (Nufarm) Kendo 22.8CS (Helm) L - C Insecticide (Drexel) Lambda-Cy AG (Winfield) Lambda-Cy 1EC (UPL) Lambdastar 1CS, Plus (FarmHannong) Lambda T 1EC (Helena) Lamcap II (Syngenta) Lunge (UPL) Nufarm Lambda-Cyhalothrin 1EC (Nufarm) Paradigm 1VC (Winfield) Province II (Tenkoz) Ravage 1EC (Innvictus) Serpent 1EC (Atticus) Silencer 1EC (ADAMA) Willowood Lambda-Cy 1EC (Generic) |
| Zeal (Valent) | etoxazole | Suremite SC (Aceto) Zara WSB, Zara SC (Atticus) |

Generic Herbicides'

| Original Trade Name (Current Manufacturer) | Common Name | Other Trade Names (Manufacturers) |
|---|---------------------|--|
| Aim® EC Herbicide (FMC Corporation) | carfentrazone-ethyl | Antik™ EC (Atticus Ag) Longbow™ EC (Nufarm Americas, Inc.) Maxunitech Carfentrazone 2 EC (Maxunitech North America, Inc.) Quinark™ EW (Atticus Ag) Shark® EW (FMC Corporation) |
| Alion® Herbicide (Bayer CropScience) | Indaziflam | Sage™ (Altamont) |
| Amine4 2,4-D (Tenkoz) | 2,4-D amine | 2,4-D Amine 4 (WinField United) 2,4-D/Amine 4 Herbicide (WinField United) Amine 4 2,4-D (Loveland Products, Inc.) Amine 6 (Loveland Products, Inc.) Base Camp® Amine 4 (Wilber-Ellis Company LLC) Clean Amine® (Loveland Products, Inc.) Cornbelt® 4Lb. Amine (Van Diest Supply Company) De-Amine® 4 (Drexel Chemical Company) Defy® Amine 4 (ADAMA) Embed® Extra (Corteva Agrisciences) Embed™ (Corteva Agriscience) Orchard Clean® (Nufarm Americas, Inc.) Orchard Master® Broadleaf Herbicide (PBI-Gordon Professional) Orchard Star® (Albaugh, LLC Agricultural Products) Rugged® Herbicide (WinField United) Saber® (Loveland Products, Inc.) Savage® Dry Soluble (Loveland Products, Inc.) Solution Water Soluble® (Nufarm Americas, Inc.) Usha 6 (Sharda USA LLC) Weedar® 64 (Nufarm Americas, Inc.) WeeDestroy® AM-40 Amine Salt (Nufarm Americas, Inc.) |
| Assure® II Herbicide (Ampac Chemical Corporation) | quizalofop p-ethyl | Targa® (Gowan Company, LLC) |
| Callisto (Syngenta Crop Protection, LLC) | mesotrione | Atticus Cavallo™ 4 SC (Atticus Ag) Bellum® (Albaugh, LLC Agricultural Products) Meso Star (Sharda USA LLC) Mesotrione 4SC (Albaugh, LLC Agricultural Products) MesoTryOne™ 4L (Drexel Chemical Company) Motif® Herbicide (UPL NA Inc.) Undercover™ (Innervictis Crop Care, LLC) |

(Continued)

Generic Herbicides' (continued)

| Original Trade Name (Current Manufacturer) | Common Name | Other Trade Names (Manufacturers) |
|---|-----------------------------|---|
| Chateau WDG (Valent USA) | flumioxazin | Chateau® EZ (Valent U.S.A. LLC) Chateau® Herbicide SW (Valent U.S.A. LLC) Flumi® 51 WDG (NuFarm Americas, Inc.) Flumi® SX Herbicide (Valent U.S.A. LLC) Flumigard Herbicide (Alligare, LLC) Flumigard SC (Alligare, LLC) Flumioxazin 51WDG Select (Alligare, LLC) Semera 51.0% WDG (Atticus EcoCore) Semera SC (Atticus EcoCore) SureGuard Herbicide (Nufarm Americas, Inc.) Tuscany® (Nufarm Americas, Inc.) Tuscany® SC Herbicide (Nufarm Americans, Inc.) Varsity® (Innvictis Crop Care, LLC) Varsity® SC (Innvictis Crop Care, LLC) Zaltus™ SC (Atticus Ag) |
| Gallery 75 DF / SC (Corteva Agrisciences) | isoxaben | Trellis and Trellis SC (Corteva AgriSciences) |
| Goal® 2XL Herbicide (Nufarm Americas, Inc.) | oxyfluorfen | Collide™ Herbicide (UPL NA Inc.) Galigan® 2E (ADAMA) Galigan® H2O Herbicide (ADAMA) GoalTender® Herbicide (Nufarm Americas, Inc.) Govee™ (Innvictis Crop Care, LLC) Oxystar® 2E (Albaugh, LLC Agricultural Products) Oxystar® 4L (Albaugh, LLC Agricultural Products) ScrollOVR™ (Atticus Ag) Scroll™ 2 XL (Atticus Ag) Willowood OxyFlo 2EC (Generic Crop Science LLC) |
| Gramoxone (Syngenta Crop Protection, LLC) | paraquat | Axill Solutions Paraquat 3SL (Axill Solutions, LLC) Devour™ (Innvictis Crop Care, LLC) Gramoxone® SL 3.0 (Syngenta Crop Protection, LLC) Helmquat 3SL (Helm Agro US, Inc.) Paraquat Concentrate (Solera Sources Dynamics, LLC) Para-Shot 3.0 (Sharda USA LLC) Parazone® 3SL (Amvac Chemical Corporation) Quik-Quat™ (Drexel Chemical Company) Willowood Paraquat 3SL (Generic Crop Science LLC) |
| HomePlate® (Certis USA, L.L.C.) | caprylic acid + capric acid | Fireworxx™ (OHP, Inc.) SUPPRESS® Herbicide EC (SAN Group Biotech USA Inc.) |
| Karmex DF (ADAMA) | diuron | Direx 4L (ADAMA) Diuron 4L/80 DF (Alligare, LLC) Diuron 4L/80 (Drexel, Chemical Company) Diuron 4L (ADAMA) Diuron 4L (Loveland Products, Inc.) Diuron 80 (Drexel Chemical Company) Diuron 80 DF (Alligare, LLC) Diuron 80 (WDG Weed Killer (Loveland Products, Inc.) Drill (Sharda USA LLC) |

(Continued)

Generic Herbicides (continued)

| Original Trade Name (Current Manufacturer) | Common Name | Other Trade Names (Manufacturers) |
|---|---------------|--|
| Kerb SC (Corteva AgriSciences) | pronamide | Kerb® 50-W Herbicide (Corteva Agriscience) Kerb® SC (RUP) (Corteva Agriscience) Willowood Pronamide 3.3SC (Generic Crop Science LLC) Willowood Pronamide 50WSP (Generic Crop Science LLC) |
| Matrix® FNV/SG (Corteva AgriSciences) | rimsulfuron | DuPont Matrix® SG (Corteva Agriscience) Grapple™ (Nufarm Americas, Inc.) Hinge™ (Albaugh, LLC Agricultural Products) Matrix® FNV (Corteva Agriscience) Pravin® Herbicide (ADAMA) Revolt™ (Innvictis Crop Care, LLC) Tetris™ SG (Atticus Ag) |
| Poast (BASF) | sethoxydim | Segment (BASF) |
| Princep® 4L (Syngenta Crop Protection, LLC) | simazine | Princep® Caliber 90° Herbicide (Syngenta Crop Protection, LLC) Simazine 4L (Drexel Chemical Company) Simazine 4L (WinField United) Simazine 4L Flowable (Loveland Products, Inc.) Simazine 90DF (Drexel Chemical Company) Simazine 90 WDG (Loveland Products, Inc.) Sim-Trol® 4L Simazine Flowable Herbicide (Sipcam Agro USA, Inc.) Sim-Trol® 90DF Simazine Dry Flowable Herbicide (Sipcam Agro USA, Inc.) |
| Prowl® 3.3 EC herbicide (BASF Ag Products) | pendimethalin | Acumen® Herbicide (TENKOZ, Inc.) Acumen® Microcap (TENKOZ, Inc.) Framework® 3.3 EC Herbicide (WinField United) Pendulum® 2G granule herbicide (BASF Professional and Specialty Solutions) Pendulum® 3.3 EC herbicide (BASF Professional and Specialty Solutions) Pendulum® AquaCap™ herbicide (BASF Professional and Specialty Solutions) Pin-Dee™ 3.3 EC (Drexel Chemical Company) Pin-Dee™ 3.3 T & O (Drexel Chemical Company) Prowl® H2O herbicide (BASF Ag Products) Satellite® 3.3 herbicide (UPL NA Inc.) Satellite® Flex (UPL NA Inc.) Satellite® HydroCap herbicide (UPL NA Inc.) Stealth® Herbicide (Loveland Products, Inc.) |
| Reglone (Syngenta Crop Protection, LLC) | diquat | Aceto Diquat 2L AG (Aceto Life Sciences, L.L.C. d/b/a Actylis) Capone™ Desiccant (Atticus Ag) Dessicash Ag (Sharda USA LLC) Nufarm Diquat 2 L (Nufarm Americas, Inc.) Nufarm Diquat SPC 2 L (Nufarm Americas, Inc.) Verdure-X-Herbicide (Helm Agro US, Inc.) |

(Continued)

Generic Herbicides (continued)

| Original Trade Name (Current Manufacturer) | Common Name | Other Trade Names (Manufacturers) |
|--|----------------------|--|
| Rely 280 (BASF Agricultural Solutions US LLC) | glufosinate-ammonium | Axill Solutions Glufosinate 280SL (Axill Solutions, LLC) Cheetah® Herbicide (Nufarm Americas, Inc.) Fever™ (Innactivis Crop Care, LLC) Forfeit® 280 (Loveland Products, Inc.) Inflammation™ 280 SL (Atticus Ag) Interline® Herbicide (UPL NA Inc.) Lifeline® (UPL NA Inc.) Opportunity (Sharda USA LLC) Reckon™ 280SL (Solera Source Dynamics, LLC) Refer® 280 SL Herbicide (Summit Agro USA, LLC) Rely® 280 Herbicide (BASF Ag Products) Scout™ Herbicide (Valent U.S.A. LLC) Surmise® (Albaugh, LLC Agricultural Products) Surmise® 5 (Albaugh, LLC Agricultural Products) Total TNV (WinField United) Total® 2.3 (WinField United) Total® Herbicide (WinField United) Willowood Glufosinate 280SL (Generic Crop Science LLC) |
| Roundup PowerMAX® Herbicide (Bayer Crop-Science) | glyphosate | Abundit® Edge (Corteva Agriscience) Aquamaster® (Bayer CropScience) Buccaneer® 5 Extra (TENKOZ, Inc.) Buccaneer® Plus (55467-9) (TENKOZ, Inc.) Clearout™ 5 Extra (Albaugh, LLC Agricultural Products) Cornerstone K Herbicide (WinField United) Cornerstone® 5 Plus (WinField United) Cornerstone® Plus (1381-192) (WinField United) Credit® 41 Extra (Nufarm Americas, Inc.) Credit® 5.4 Extra (Nufarm Americas, Inc.) Credit® Xtreme (Nufarm Americas, Inc.) Duramax® Herbicide (Corteva Agriscience) Durango® DMA® Herbicide (Albaugh, LLC Agricultural Products) Envy™ (Innactivis Crop Care, LLC) Envy™ Intense (Innactivis Crop Care, LLC) Envy™ Six Max (Innactivis Crop Care, LLC) Four Power Plus® (Loveland Products, Inc.) Gly Star® 5 Extra (Albaugh, LLC Agricultural Products) Gly Star® K-Plus (Albaugh, LLC Agricultural Products) Gly Star® Original (Albaugh, LLC Agricultural Products) Gly Star® Plus (Albaugh, LLC Agricultural Products) Honcho® K6 Herbicide (Bayer CropScience) Honcho® Plus (Bayer CropScience) Mad Dog® (Loveland Products, Inc.) Mad Dog® Plus (Loveland Products, Inc.) Makaze® Herbicide (Loveland Products, Inc.) |

(Continued)

Generic Herbicides (continued)

| Original Trade Name (Current Manufacturer) | Common Name | Other Trade Names (Manufacturers) |
|---|--------------|---|
| Roundup PowerMAX® Herbicide (Bayer Crop-Science) (Continued) | glyphosate | Roundup PowerMAX® 3 Herbicide (Bayer Crop-Science) Roundup PowerMAX® Herbicide (Bayer CropScience) Roundup PowerMAX® II (Bayer CropScience) Roundup Ultra® (Bayer CropScience) Roundup WeatherMAX® Herbicide (Bayer Crop-Science) Shar-Max Glyphosate 41% SL (Sharda USA LLC) Willowood Glypho 6 (Generic Crop Science LLC) Wynca USA™ Sunphosate 41% Herbicide (Wynca USA) Wynca USA™ Sunphosate® 5 MAX (Wynca USA) |
| Sandea® (Gowan Company, LLC) | halosulfuron | Herbivore® Herbicide (WinField United) |
| Select Max® Herbicide with Inside Technology™ (Valent U.S.A. LLC) | clethodim | Arrow® 2 EC (ADAMA) Avatar™ (Innvictis Crop Care, LLC) Avatar™ S2™ (Innvictis Crop Care, LLC) Axill Solutions Clethodim 2EC (Axill Solutions, LLC) Ceridian™ 2 EC (Atticus Ag) Cleanse® (WinField United) Cleanse® 2EC (WinField United) Clethodim 2E (Albaugh, LLC Agricultural Products) Dakota™ (Albaugh, LLC Agricultural Products)Envoy Plus™ Herbicide (Valent U.S.A. LLC Professional Products) Shadow® (UPL NA Inc.) Shadow® 3EC (UPL NA Inc.) Signature Clethodim (Albaugh, LLC Agricultural Products) Trizenta™ 3EC Herbicide (UPL NA Inc.) Trizenta™ Herbicide (UPL NA Inc.) Vaquero® (Wilbur-Ellis Company LLC) Volunteer® (TENKOZ, Inc.) Volunteer® Herbicide (42750-72-55467) (TENKOZ, Inc.) Volunteer® Herbicide (70506-484-55467) (TENKOZ, Inc.) Willowood Clethodim 2EC (Generic Crop Science LLC) |
| Stinger® Herbicide (Corteva AgriSciences) | clopyralid | Bite (Sharda USA LLC) Clean Slate (Nufarm Americas, Inc.) GCS Clopy 360SL (Generic Crop Science LLC) Spur® (Albaugh, LLC Agricultural Products) Stigmata™ (Atticus Ag) |
| Starane® Ultra Herbicide (Corteva Agriscience) | fluroxypyr | Comet® Selective Herbicide (Nufarm Americas, Inc.) Fancy (Sharda USA LLC) Stark™ Ultra (Atticus Ag) |
| Surflan AS | oryzalin | Fugitive (ADAMA) Oryzalin 4 AS (ADAMA) |
| Trellis® (Corteva Agriscience) | isoxaben | Trellis® SC (Corteva Agriscience) Gallery® 75 Dry Flowable Specialty Herbicide (Corteva Agriscience) Gallery® SC (Corteva Agriscience) |

(Continued)

Generic Herbicides (continued)

| Original Trade Name (Current Manufacturer) | Common Name | Other Trade Names (Manufacturers) |
|---|------------------|--|
| Treflan™ 4L Herbicide (Loveland Products, Inc.) | trifluralin | Treflan HFP Herbicide (Gowan Company, LLC) Treflan TR-10 Granular Herbicide (Gowan Company, LLC) Trifluralin 10G (Loveland Products, Inc.) Trifluralin 4 EC Herbicide (Aceto Life Sciences, L.L.C. d/b/a Actylis) Trifluralin 4EC (Albaugh, LLC Agricultural Products) Trifluralin 4EC (Drexel Chemical Company) Trifluralin HF (Loveland Products, Inc.) Trust (WinField United) |
| Ultra Blazer (United Phosphorus) | acifluorfen | Acifin 2 L (Summit) Acifluorfen 20.1% (Sharda) Avalanche Ultra (WinField) Derecho (Atticus) Levity (Innvictis) Uproar (WinField) |
| Velpar DF VU/L VU (Bayer) | hexazinone | Tide Hexazinone 2 SL/75 WDG (Tide Int'l) Velossa (Helena) Velpar L CU/DF CU (Tessenderlo) |
| Venue® (Nichino America, Inc.) | pyraflufen ethyl | Venue® Max Herbicide (Nichino America, Inc.) |
| Zeus /XC (FMC Corporation) | sulfentrazone | Maxunitech Sulfentrazone 4 SC Herbicide (Maxunitech North America, Inc.) Passage™ Herbicide (Alligare, LLC) Shutdown® Herbicide (UPL NA Inc) Willowood Sulfen 4SC (Generic Crop Science LLC) Zone 4F (Helm Agro US, Inc.) |

¹ Check label to make sure product is labeled for the crop that it is to be used on.

Fruit Grower Newsletters

Arkansas

University of Arkansas Division of Agriculture Cooperative Extension Service offers *Arkansas Fruit, Vegetable, and Nut Update*. It is published monthly or as needed to Arkansas growers at no cost. It provides timely information about fruit and nut production practices, disease and insect/mite activity, and upcoming meetings. To subscribe, go to: <https://www.uaex.uada.edu/farm-ranch/crops-commercial-horticulture/horticulture/ar-fruit-veg-nut-update-blog/>

Illinois

University of Illinois Extension publishes *Illinois Fruit & Vegetable News* (ipm.illinois.edu/ifvn). This newsletter covers production practices and insect and disease management. For more information, contact Local Food Systems and Small Farms Educator: Bronwyn Aly (1715 College Ave., Carmi, IL 62821, 618-395-2441, baly@illinois.edu); or Commercial Agriculture Educator: Nathan Johanning, 901 Illinois Avenue, PO Box 117, Waterloo, IL 62298, 618-939-3434). For disease and insect diagnostics and management recommendations, contact the University of Illinois Plant Clinic at S-417 Turner Hall 1102 S. Goodwin Ave., Urbana IL 61801, 217-333-0519; plantclinic@illinois.edu.

Indiana

Purdue Extension offers *Facts for Fancy Fruit*, a newsletter issued biweekly throughout the growing season, that provides timely information on diseases and insects throughout the state, cultural practices and announcements about upcoming events. Subscribe to the online version free of charge at fff.hort.purdue.edu or receive a printout via first class mail for \$15 a year.

For a hard copy, send your name, address, and current fruit interests along with a check for \$15, made out to Purdue University to: *Facts For Fancy Fruit*, Department of Horticulture and Landscape Architecture, 625 Agricultural Mall Drive, Purdue University, West Lafayette, IN 47907-2010.

Iowa

You can find general horticulture information and Iowa State University Plant and Insect Diagnostic Clinic updates at <https://yardandgarden.extension.iastate.edu/>

Kentucky

Cooperative Extension issues a monthly newsletter, *Kentucky Fruit Facts* (<https://horticulture.mgcafe.uky.edu/ky-fruit-facts>), to all Kentucky growers at no cost. This service supplies timely information on disease and insect activity throughout the state, as well as cultural information.

To subscribe, send an email message:

TO: listserv@lsv.uky.edu
SUBJECT: Fruit Facts
MESSAGE: subscribe KY-FRUITFACTS

Followed by a blank line

OR to unsubscribe, the lines:
signoff KY-FRUITFACTS
Followed by a blank line

You should receive confirmation by return email. If you have a problem, or if you wish to communicate with a person about "fruitfacts", the owner's address (the TO: line of the message) is: owner-ky-fruit-facts@lsv.uky.edu

Ohio

Ohio Fruit News (OFN) is published six times a year by the Department of Plant Pathology at The Ohio State University, CFAES-Wooster. The newsletter is available free of charge in electronic format at u.osu.edu/fruitpathology/fruit-news-2/, or as a printed copy by request. To subscribe, contact Melanie Lewis Ivey at ivey.14@osu.edu or 330-263-3849.

The Ohio State University Extension Specialty Crop Team maintains an online blog, *Fruit, Vegetable, and Specialty Crop News* (u.osu.edu/vegnetnews/). New posts are added daily and feature timely updates and information on all specialty food crops.

The *Ohio Grape-Wine Electronic Newsletter (OGEN)* is available at ohiograpeweb.cfaes.ohio-state.edu/news. To subscribe, email Maria Smith at smith.127203@osu.edu.

Pesticide Drift Communication Tools

Several states involved in this spray guide have web-based mapping tools that enable producers of pesticide sensitive crops avoid drift injury by communicating with agricultural chemical applicators.

DriftWatch.org serves Colorado, Delaware, Illinois, Indiana, Iowa, Kansas, Michigan, Minnesota, Missouri, Montana, Nebraska, Ohio, New Mexico, North Carolina, Wisconsin, and Saskatchewan, Canada.

The Ohio Sensitive Crop Registry is available at www.agri.ohio.gov/scr <https://agri.ohio.gov/divisions/plant-health/resources/oscr>.

Check with the state department of agriculture in your state about similar tools.

Network for Environment and Weather Applications (NEWA)

NEWA collects weather data through the Internet from weather stations primarily located on farms and generates real-time weather data summaries, crop production tools, and IPM forecasts.

Contact your NEWA State coordinators for additional information on purchasing a weather station and connecting to NEWA for IPM weather tools and IPM crop forecasts. If you don't have a state coordinator you can contact Dan Olmstead at newa@cornell-ipm.org.

Illinois: Elizabeth Wahle, University of Illinois at Urbana-Champaign; 618-344-4230; wahle@illinois.edu.

Kentucky: Brent Arnoldussen, University of Kentucky; 859-257-4721; bar244@uky.edu.

Ohio: Melanie Lewis Ivey, The Ohio State University, CFAES-Wooster; 330-263-3849; ivey.14@osu.edu

Wisconsin: Amaya Atucha, University of Wisconsin-Madison; 608-262-6452; atucha@wisc.edu

Using a Plant Diagnostic Lab

The best way to identify insects, plants, and plant diseases, or to diagnose plant and pest problems, is to send a sample to a diagnostic laboratory along with information and observations about the problem. The National Plant Diagnostic Network website (www.npdn.org) lists diagnostic laboratories by state and region. Check with your local County Extension Office or Diagnostic lab for any costs associated with this service.

To ensure an accurate diagnosis, it's important to collect and ship your specimens properly. Here are a few guidelines for collecting and shipping specimens to a diagnostic lab. *Your state has specific instructions for collecting and shipping samples; check your local clinic's website for details.*

1. Collect fresh specimens. Send a generous amount of material, if available.
2. Ship specimens in a crush-proof container immediately after collecting. If holdover periods are encountered, keep specimen cool. Ship packages early in the week so they to arrive on weekdays.
3. Incomplete information or poorly selected specimens may result in an inaccurate diagnosis or inappropriate control recommendations. Badly damaged specimens are often unidentifiable, and additional sample requests can cause delays.

Submitting Plant Specimens for Disease/Injury Diagnosis

Herbaceous Plants. For generally declining, wilting, or dying plants, send several whole plants showing a range of symptoms — early through more advanced — with roots and adjacent soil intact, if possible. Dig the plants carefully so the root system remains intact. Place roots and surrounding soil in a plastic bag and fasten it to the base of the stem with a twist tie or string. Wrap the plants in dry newspaper and place in a crush-proof container for shipment. Do not add water or moist paper towels.

Leaves/fruit/woody tissues. When localized infections (such as leaf spots, fruit rots, or cankers) are suspected, send specimens representing early and moderate stages of disease. Press leaves flat between heavy paper or cardboard — do not tape leaves to paper — and wrap fruits and woody tissue in dry paper. For large fruit, wrap each individually in newspaper. Do not place soft fruit (i.e., strawberry, raspberry, blackberry, etc.) in plastic bags. Pack firmly in a crush-proof container so that fruit is not bruised during shipping.

Submitting Insect Specimens

Package insects carefully so they aren't crushed when they arrive at the lab. Do not tape insects to paper or package them loosely in envelopes. Separate and label the specimens if you send more than one type in the same package. Provide the appropriate information for each specimen.

Tiny or Soft-bodied Specimens. Submit such specimens (aphid, mites, thrips, caterpillars, grubs, spiders) in a small, leak-proof bottle or vial that is 1 ounce or less filled with 70 percent rubbing (Isopropyl) alcohol or hand sanitizer. In Kansas, submit in vinegar. Do not submit insects in water or formaldehyde, or without rubbing alcohol; they will ferment and decompose.

Hard-bodied Specimens. Submit such specimens (flies, grasshoppers, cockroaches, wasps, butterflies, beetles) dry in a crush-proof container. As noted above, do not tape insects to paper or place them loose in envelopes.

Submitting Samples for Nematode Analysis

If you suspect a nematode problem, contact your clinic for state-specific submission information (see page 288).

In general nematode identification requires collection of at least one quart of soil from the root zone of affected plants. Include roots if the plants are actively growing.

Place the entire sample in a plastic bag. Do not add water or allow it to dry out. Protect the sample from extreme heat (for example, don't leave samples inside a parked vehicle in direct sunlight). It is often helpful to collect a second, similar sample from a nearby area where plant growth appears normal.

Attach a label, note, or tag identifying the sample to the outside of each bag or package.

Selected University Diagnostic Labs

Arkansas

Plant Health Clinic University of Arkansas

2601 N. Young Ave.

Fayetteville, AR 72704

479-502-9713

<https://www.uaex.uada.edu/yard-garden/plant-health>

Clinic: [https://www.facebook.com/UAEXPlant-](https://www.facebook.com/UAEXPlant-HealthClinic)

HealthClinic

Contact:

Taylor Klass, tklass@uada.edu

Illinois

University of Illinois Plant Clinic

S-417 Turner Hall

1102 S. Goodwin Ave.

University of Illinois

Urbana, IL 61801

217-333-0519

<https://extension.illinois.edu/plant-clinic>

plantclinic@illinois.edu

www.facebook.com/UofIPlantClinic

Contacts:

Diane Plewa, dplewa@illinois.edu, (217) 300-3441

Indiana

Plant and Pest Diagnostic Laboratory

Purdue University

LSPS 101

915 Mitch Daniels Blvd, LSPS 116

West Lafayette, IN 47907-2054

765-494-7071

Fax: 765-494-3958

<https://ag.purdue.edu/department/btny/ppdl/index.html>

ppdl-samples@purdue.edu

www.facebook.com/PurduePPDL

Contacts:

Tom Creswell, creswell@purdue.edu

John Bonkowski, jbonkows@purdue.edu

Iowa

Iowa State University Plant and Insect Diagnostic Clinic

2445 ATRB

2213 Pammel Dr

Ames, IA 50011

515-294-0581

Fax: 515-294-9420

<https://yardandgarden.extension.iastate.edu/pidc>

pidc@iastate.edu

www.facebook.com/ISUPIDC

Kentucky

Plant Disease Diagnostic Laboratory Agricultural

Science Building-North

1100 South Limestone Street

University of Kentucky

Lexington, KY 40546-0091

859-257-8949

Fax: 859-323-1961

<https://plantpathology.mgcafe.uky.edu/extension/diagnostic-laboratories>

Contact:

Julie Beale, jbeale@uky.edu

Ohio

C. Wayne Ellett

The Ohio State University, CFAES-Wooster

234 Selby Hall

1680 Madison Avenue

Wooster, OH 44691

Phone: 330-263-3650

Email: ppdc@osu.edu

ppdc.osu.edu

Contact:

Francesca Rotondo, rotondo.11@osu.edu

CFAES-Wooster, 330-263-3650

Wisconsin

Plant Disease Diagnostics Clinic Department of Plant

Pathology

1630 Linden Drive

University of Wisconsin-Madison

Madison, WI 53706-1598

pddc.wisc.edu

608-262-2863

Fax: 608-263-2626

Contact:

Brian Hudelson, hudelson@wisc.edu

Pesticide Applicator Safety Education Programs

Below are the state pesticide education programs that provide training and educational materials for becoming a certified pesticide applicator. Find other state pesticide safety education programs at [https://nifacontacts.ipmcenters.org/PSEPDirectory.cfm](https://nifacontacts.ipmcenters.org/PSEPDDirectory.cfm).

University of Arkansas

<https://uaex.uada.edu/farm-ranch/pest-management/pesticide-licensing/private-applicator-training.aspx>

University of Illinois

<https://extension.illinois.edu/psep>

Iowa State University

www.extension.iastate.edu/psep

University of Kentucky

<https://entomology.mgcafe.uky.edu/uk-pesticide-safety-education-program-psep>

Ohio State University

pested.osu.edu

Purdue University

<https://ag.purdue.edu/department/extension/ppp/>

University of Wisconsin

<https://fyi.extension.wisc.edu/pat>

Pesticide Emergency and Poison Control Centers

Nationwide phone numbers

Pesticide Poisoning: Call the **Poison Center**, 800-222-1222

This number automatically connects you to the poison center nearest you.

National Pesticide Information Retrieval System

(NPIRS): 765-494-5249

National Pesticide Information Center:

800-858-7378

CHEMTREC: (800) 424-9300

Arkansas

Arkansas Poison Center: 800-222-1222

Arkansas State Plant Board: 501-225-1595

Pesticide training, licensing, and education for applying restricted use pesticides.

Illinois

Illinois Poison Control Centers Emergency

Nationwide: 800-222-1222

Emergency TTY/TDD: 312-906-6185

Indiana

Indiana Poison Center: 800-222-1222

Pesticide Poisoning

Indiana Department of Environmental

Management:

888-233-7745 or 317-233-7745

Pesticide Spill Reporting

Purdue Pesticide Programs: 765-494-4566

General Information

Office of Indiana State Chemist: 765-494-1492

Pesticide Certification and Training

Environmental Protection Agency Region 5:

800-621-8431 or 312-353-2000

Iowa

Iowa Statewide Poison Control Center Emergency

Phone Number: 800-222-1222

Administrative Phone Number: 712-273-7757

Kentucky

Kentucky Regional Poison Control Center:

800-222-1222

KY Environmental Response: 800-928-2380 or

502-564-2380

Ohio

Ohio Poison Exposure Centers: 800-222-1222

TDD number: 800-253-7955

Wisconsin

Wisconsin Poison Center: 800-222-1222

Conversion Factors for Weights and Measures: Equivalents

| | Metric | U.S. |
|---------------|-------------------------------|------------------------|
| Length | 1 Millimeter | 0.039 inch |
| | 1 Centimeter (10 mm) | 0.39 inch |
| | 1 Meter (100 cm) | 39.4 inch |
| | 1 Kilometer (1,000 m) | 0.62 mile |
| Area | 1 Square Centimeter | 0.155 square inch |
| | 1 Square Meter | 1.2 square yards |
| | 1 Hectare (10,000 sq m) | 2.47 acres |
| | 1 Square Kilometer (100 ha) | 247 acres |
| Weight | 1 Gram | 0.035 ounces |
| | 1 Kilogram (1,000 g) | 2.2 pounds |
| | 1 Ton (metric) – 1,000 kg | 1.1 tons (U.S.) |
| Volume | 1 Milliliter | 0.034 fluid ounces |
| | 1 Liter (1,000 ml) | 1.056 quarts |
| | 1 Cubic Meter (1,000 l) | 264.17 gallons (U.S.) |
| | U.S. | Metric |
| Length | 1 Inch | 2.54 centimeters |
| | 1 Foot (12 in) | 30.5 centimeters |
| | 1 Yard (3 ft) | 0.91 meters |
| | 1 Mile (5,280 ft) | 1.6 kilometers |
| Area | 1 Square Inch | 6.5 square centimeters |
| | 1 Square Foot (144 sq in) | 930 square centimeters |
| | 1 Square Yard (9 sq ft) | 0.84 square meters |
| | 1 Acre (43,560 sq ft) | 0.405 hectares |
| | 1 Square Mile (640 acres) | 259 hectares |
| Weight | 1 Ounce | 28.3 grams |
| | 1 Pound (16 oz) | 0.454 kilograms |
| | 1 Ton (U.S.) – 2,000 lb | 0.907 tons (metric) |
| Volume | 1 Tablespoon (3 teaspoons) | 14.79 milliliters |
| | 1 Fluid ounce (2 tablespoons) | 29.6 milliliters |
| | 1 Cup (8 fl oz) | 0.237 liters |
| | 1 Pint (2 cups) | 0.473 liters |
| | 1 Quart (4 cups) | 0.946 liters |
| | 1 Gallon (U.S.) – 4 qts | 3.8 liters |
| | 1 Cubic Foot | 28.3 liters |

Metric Abbreviations: mm=millimeter; cm=centimeter; m=meter; km=kilometer; ha=hectare; mg=milligram; g=gram; kg=kilogram; ml=milliliter; l=liter.

NOTES

Midwest Fruit Pest Management Guide 2026-2027

The Midwest Fruit Pest Management Guide 2026-2027 was developed by the Midwest Fruit Workers Group.

Printed copies of this publication are available from the Purdue Extension Education Store, <https://edustore.purdue.edu>. A free PDF download also is available from the Education Store.

Citation

Long, E., Leach, A., Athey, K., Wimmer, M., Arnoldussen, B., Yadav, R., Meyer, S., Purcell, M., Santiago, J., Holland, L., Joshi, N., Whale, E., Babadoost, M., Vos, R., Slack, S., Cloyd, R., Bessin, R., Ivey, M., Wright, S., Beckerman, J., Leonberger, K., Gauthier, N. 2025. Midwest Fruit Pest Management Guide 2026-2027. Available online at: https://ag.purdue.edu/department/hla/extension/_docs/id-465.pdf

Additional Contributors

University of Arkansas

Entomology: Neelendra Joshi

University of Illinois

Entomology: Kacie Athey*

Horticulture: Elizabeth Wahle*, James Santiago*

Plant Pathology: Mohammed Babadoost*

Iowa State University

Horticulture: Randall Vos*, Suzanne Slack

Kansas State University

Entomology: Raymond Cloyd

University of Kentucky

Entomology: Ric Bessin*

Horticulture: Shawn Wright*, Brent Arnoldussen*, Elayna Stirn

Plant Pathology: Nicole Gauthier*, Kim Leonberger*

University of Minnesota

Horticulture: Madeline Wimmer*

Ohio State University

Entomology: Ashley Leach*

Horticulture: Ram Yadav*

Plant Pathology: Melanie Ivey*

Purdue University

Entomology: Elizabeth Long*

Horticulture: Stephen Meyer, Miranda Purcell

University of Wisconsin

Plant Pathology: Leslie Holland*

*Section Editor

Disclaimer

Reference in this publication to any specific commercial product, process, or service, or the use of any trade, firm, or corporation name is for general informational purposes only and does not constitute an endorsement, recommendation, or certification of any kind by Midwest Fruit Workers Group or their institutions. Individuals using such products assume responsibility for their use in accordance with current directions of the manufacturer.

Non-Discrimination Statement

The U.S. Department of Agriculture (USDA) prohibits discrimination in all its programs and activities on the basis of race, color, national origin, age, disability, and where applicable, sex, marital status, familial status, parental status, religion, sexual orientation, genetic information, political beliefs, reprisal, or because all or part of an individual's income is derived from any public assistance program. (Not all prohibited bases apply to all programs.) Persons with disabilities who require alternative means for communication of program information (Braille, large print, audiotape, etc.) should contact USDA's TARGET Center at 202-720-2600 (voice and TDD).

To file a complaint of discrimination, write USDA, Director, Office of Civil Rights, 1400 Independence Avenue, SW, Washington, DC 20250-9410 or call 800-795-3272 or 202-720-6382 (TDD).

USDA is an equal opportunity provider and employer.

Acknowledgements

Authors would like to sincerely thank Tammy Luck, Lead Administrative Assistant, Department of Entomology, Purdue University for her dedicated work in formatting this guide.

Authors would like to acknowledge Dr. Janna Beckerman for her years of devoted work to the development of this guide.